

US EPA ARCHIVE DOCUMENT



RADON: TEST, FIX, SAVE A LIFE.



## WINTER IS A GREAT TIME TO TEST YOUR HOME!



## Basic Radon Facts

### Radon is a cancer-causing, radioactive gas.

Radon is a naturally occurring radioactive gas released in rock, soil, and water from the natural decay of uranium. While levels in outdoor air pose a relatively low threat to human health, radon can accumulate to dangerous levels inside buildings. You can't see, smell, or taste it, but an elevated radon level in your home may be affecting the health of your family.

Exposure to radon is the second leading cause of lung cancer in the United States and the number one cause among non-smokers. The U.S. Environmental Protection Agency estimates that radon causes more than 20,000 lung cancer deaths in the country each year. Only smoking causes more lung cancer deaths. If you smoke and your home has radon, your risk of lung cancer can be higher.



### Radon is found all over the United States.

Radon has been found in elevated levels in homes in every state. No area of the country is free from risk. Indeed, two homes right next to each other can have vastly different radon levels. Just because your neighbor's house does not have an elevated level of radon does not mean that your house will have a low radon level. The only way to know if your home is under the EPA action level of 4 pCi/L is to test.

High levels of radon in homes usually come from the surrounding soil. Radon gas enters through cracks and openings—such as sump pump lids and plumbing features—on the lower levels of your home. Hot spots include basements, first-floor rooms, and garages, but radon can be found anywhere in your house.

### You should test for radon.

The U.S. Surgeon General recommends that all homes in the U.S. be tested for radon. Testing your house for radon is easy to do. If your house has a radon problem, you can take steps to fix it to protect yourself and your family.

## How to Obtain Radon Test Kits

To obtain an easy-to-use radon test kit, you can:

- Purchase a test kit from your local home improvement or hardware store. Many kits are priced under \$25.00.
- Use the radon test kit coupon found at [www.epa.gov/oncampus](http://www.epa.gov/oncampus). For more information on obtaining test kits and test kit coupons, visit [www.sosradon.org](http://www.sosradon.org).
- Contact your state radon program, details available at [www.epa.gov/iaq/whereyoulive.html](http://www.epa.gov/iaq/whereyoulive.html), for more information about obtaining test kits from your state or from a radon testing company or laboratory in your area.

## Basic Radon Facts (continued)

You can find out if your home has an elevated radon level by conducting a simple test. It's as easy as opening a package, placing a radon detector in a designated area, and, after a set number of days, sending the detector back to a lab for analysis. The lab will then inform you of your radon test results.

Radon test kits are available at your local home improvement or hardware store, or in some cases from your state radon office. You can also order them from Kansas State University (KSU) or radon testing companies. Another option is to hire a qualified tester to do a radon test for you. Contact your state radon office about obtaining a list of qualified testers. Information about testing your home for radon and finding a test kit is also available by calling 1-800-SOS-RADON.

Radon is measured in picocuries per liter of air (pCi/L), a measurement of radioactivity. EPA and the Centers for Disease Control and Prevention recommend that homes with radon levels at 4 pCi/L or higher should be fixed. EPA also recommends that Americans consider fixing their homes for radon levels between 2 pCi/L and 4 pCi/L. Based on a national residential radon survey completed in 1991, the average indoor radon level is about 1.3 pCi/L in the United States. The average outdoor level is about 0.4 pCi/L.

### You can fix a radon problem.

The cost of making repairs to reduce the radon level depends on several factors, including how your home was built. Most homes can be fixed for about the same cost as other common home repairs, like painting or having a new hot water heater installed. Look in your local phone book or call your state radon office to locate radon mitigators in your area if you find an elevated radon level in your home.

### New homes can be built with radon-resistant features.

Radon-resistant construction methods can be effective in reducing radon entry. When used properly, these simple and cost-effective techniques can help reduce the accumulation of radon gas in homes.

Every new home should be tested after occupancy, even if it was built using radon-resistant construction methods. If radon levels above EPA's action level at or above 4 pCi/L are detected, it is easier and less expensive to reduce radon levels in homes that have been built with radon-resistant construction techniques.

### For more information:

Contact your State Radon Office

<http://www.epa.gov/radon/wherelive.html>



*Radon presents a serious health risk, but it can be controlled easily and cost-effectively. Take action today. Encourage your friends and family members to do the same!*