

BINATIONAL FRAMEWORK FOR IDENTIFYING SUBSTANCES OF POTENTIAL THREAT TO THE GREAT LAKES BASIN

Test Case: Polybrominated Diphenyl Ethers (PBDEs)

I. FEEDERS FOR SUBSTANCE IDENTIFICATION

National Chemical Management Programs

Canada

PBDEs have been assessed CEPA toxic under the Canadian Environmental Protection Act (CEPA, 1999) and were added to CEPA 1999 Schedule 1- List of Toxic Substances.¹

Brominated flame retardants (including all PBDEs) are included in the monitoring plan for Chemicals Management Plan (CMP) (either Year 1 or Year 2).²

United States

Decabromodiphenylether (DecaBDE) is a High Production Volume (HPV) chemical.³

Great Lakes Monitoring and Surveillance

Yes, there are specific PBDEs (identified as PBDE congeners) substances common with the Great Lakes Screening Project. These include:⁴

<i>PBDE Congener</i>	<i>CAS No.</i>
tetrabromodiphenyl ether (tetraBDE)	40088-47-9
pentabromodiphenyl ether (pentaBDE)	32534-81-9
hexabromodiphenyl ether (hexaBDE)	36483-60-0
heptabromodiphenyl ether (heptaBDE)	68928-80-3
octabromodiphenyl ether (octaBDE)	32536-52-0
nonabromodiphenyl ether (nonaBDE)	63936-56-1
decabromodiphenyl ether (decaBDE)	1163-19-5

Other Sources of Information

United States

Several states, including California, Maine, Michigan, New York, Hawaii, Washington, Maryland and Illinois, have enacted or are considering legislation prohibiting products containing more than 0.1%/wt of PentaBDE or OctaBDE commercial mixtures.⁵

Europe

The European Community has adopted Directives to minimize PentaBDE, OctaBDE and DecaBDE commercial mixtures in electrical and electronic equipment (EEE) and to manage EEE waste. As of July 1, 2006 (July 1, 2008 for DecaBDE), specified EEE products are not allowed to be placed on the European Union (EU) market if a product contains more than 0.1%/wt of PBDEs.⁶

The European Community has also adopted a Directive, banning the manufacture or use of PentaBDE and OctaBDE commercial mixtures and the sale of products containing more than 0.1% PentaBDE or OctaBDE.⁷

Norway

In Norway, products containing more than 0.25% PentaBDE, OctaBDE or DecaBDE commercial mixtures are classified as hazardous waste when they are discarded. Norway has enacted a regulation whereby, as of April 1, 2008, all manufacture, import, export, sale, and use of DecaBDE as a substance or in preparations containing greater than 0.1%/wt of the compound is banned. Transport applications are the only exception.

Japan

Japan has a law concerning the reporting of releases of specific chemical substances and promoting improvements in their management. Under this law, yearly reports are required on volumes of DecaBDE commercial mixture imported and used and quantities released to the environment.⁸

China

China's Ministry of Information Industries issued the Management Methods for the Prevention and Control of Pollution Caused by Electronic Information Products. The Methods provide a broad regulatory framework for restrictions of "toxic and hazardous substances", including PBDEs.⁹ As of March 1, 2007, EEE products need to meet marking and disclosure requirements, including for PBDEs.

Sweden

Sweden reversed a ban on DecaBDE in textiles, furniture and some cables. Instead, the Swedish government is seeking a total ban on the use of DecaBDE at the EU level.

United Nations Economic Commission for Europe (UNECE) Long-Range Transboundary Air Pollution (LRTAP)

In December 2005, Parties agreed that PentaBDE commercial mixtures should be considered as a persistent organic pollutant (POP). As well, the European Commission in 2005 submitted a proposal to add OctaBDE commercial mixture to the Protocol.¹⁰

POP Review Committee under UN Stockholm Convention

In November 2006, the Persistent Organic Pollutants Review Committee under the United Nations Stockholm Convention on POPs decided that commercial PentaBDE warrants global action and is currently preparing a risk management evaluation prior to deciding whether to recommend addition of the substance to the Convention. The Committee also decided that commercial OctaBDE meets the criteria outlined in the Convention for consideration as a persistent organic pollutant, and a risk profile has been prepared to support a decision as to whether global action is warranted.¹¹

II. CONSIDERATIONS FOR SUBSTANCE SELECTION

Monitoring and Surveillance

Numerous monitoring and surveillance data are available which demonstrates the presence of PBDEs in the Great Lakes Basin. For example:

- Environment Canada 2006, “Canadian Environmental Protection Act, 1999- Ecological Screening Assessment Report on Polybrominated Diphenyl Ethers (PBDEs).¹²
 - Reports concentrations in Great Lake water, sediments, biota, ambient environment & sewage sludge.
 - Luckey *et al.* 2002 measured total (dissolved and particulate phases) PBDE (mono- to heptaBDE congeners) of approximately 9 pg/L in Lake Ontario surface waters in 1999. More than 60% of the total was composed of tetraBDE and pentaBDE, with pentaBDE and heptaBDE congeners each contributing approximately 5 to 8% of the total.
 - Stapleton and Baker 2001 analyzed water samples from Lake Michigan in 1997, 1998 and 1999 and found that total PBDE concentrations ranged from 31 to 158 pg/L.
 - Kolic *et al.* 2004 determined levels of PBDEs in sediments from Lake Ontario tributaries flowing to Lake Ontario. The total PBDEs (tri, tetra, penta, hexa, hepta, and decaBDEs) measured in sediment samples taken from 14 tributary sites (6 reported) ranged from approximately 12 to 430 µg/kg dw. Concentrations of tetra- to hexaBDEs ranged from approximately 5 to 49 µg/kg dw. Concentrations of BDE209 (decaBDE) ranged from 6.9 to 400 µg/kg dw. BDE 47 (hexaBDE), 99 (pentaBDE), and 209 were the predominant congeners measured in sediments.
 - Muir *et al.* (2003) measured concentrations of BDE209 along a north-south transect from southern Ontario/upper New York state to Ellesmere Island. The highest concentrations of BDE209 (up to 12 µg/kg dw) occurred in sediments collected from the western basin of Lake Ontario.

Table 4. Measured concentrations of PBDEs in the North American ambient environment and sewage sludge

Medium	Location; year	Total PBDEs	Reference
Air	Alert, Canada; 1994–1995	1–28 pg/m ³	Alaee et al. 2000
Air	Great Lakes; 1997–1999	5.5–52 pg/m ³	Strandberg et al. 2001
Air	Southern Ontario; 2000	10–1300 pg/m ³	Gouin et al. 2002
Air	Ontario; 2000	3.4–46 pg/m ³	Harner et al. 2002
Water	Lake Michigan; 1997–1999	31–158 pg/L	Stapleton and Baker 2001
Water	Lake Ontario; 1999	6 pg/L	Luckey et al. 2002
Sediment	Lake Michigan; 1998	4.2 µg/kg dw	Stapleton and Baker 2001
Sediment	British Columbia; 2001	2.7–91 µg/kg OC	Rayne et al. 2003a
Soil	United States; 2000	<0.1–76 µg/kg dw	Hale et al. 2002
Sewage sludge	Toronto, Canada United States	8280 µg/kg dw 730–24 900 µg/kg dw	La Guardia et al. 2001
Sewage sludge	United States; 2000	3005 µg/kg dw	Hale et al. 2002
Sewage sludge	Southern Ontario	1700–3500 µg/kg dw	Kolic et al. 2004

dw = dry weight; OC = organic carbon

Table 5. Measured concentrations of PBDEs in North American biota

Organism	Location; year	Total PBDEs	Reference
Dungeness crab hepatopancreas	West coast, Canada; 1993–1995	4.2–480 µg/kg lipid	Ikonomou et al. 2002b
Mountain whitefish (muscle)	Columbia River, British Columbia; 1992–2000	0.726–131 µg/kg ww	Rayne et al. 2003a
Heron egg	British Columbia; 1983–2000	1.308–288 µg/kg ww	Wakeford et al. 2002
Murre egg	Northern Canada; 1975–1998	0.442–2.93 µg/kg ww	
Fulmar egg	Northern Canada; 1975–1998	0.212–2.37 µg/kg ww	
Beluga whale blubber	Canadian Arctic	81.2–160 µg/kg lipid	Alaee et al. 1999
Herring gull egg	Great Lakes; 1981–2000	9.4–1544 µg/kg ww	Norstrom et al. 2002
Lake trout	Lake Ontario; 1997	95 µg/kg ww	Luross et al. 2002
	Lake Erie; 1997	27 µg/kg ww	
	Lake Superior; 1997	56 µg/kg ww	
	Lake Huron; 1997	50 µg/kg ww	
Rainbow trout	Spokane River, Washington, USA; 1999	297 µg/kg ww	Johnson and Olson 2001
Mountain whitefish		1250 µg/kg ww	
Largescale sucker		105 µg/kg ww	
Carp	Virginia, USA; 1998-1999	1140 µg/kg ww	Hale et al. 2001


- Report available at:
[http://www.ec.gc.ca/CEPARRegistry/documents/subs_list/PBDE_SAR/PBDEs_SAR_EC_June_2006_\(en\).pdf](http://www.ec.gc.ca/CEPARRegistry/documents/subs_list/PBDE_SAR/PBDEs_SAR_EC_June_2006_(en).pdf).

Environmental Levels and Trends

Some trends identified in Environment Canada's 2006 Ecological Screening Assessment Report on Polybrominated Diphenyl Ethers (PBDEs) are as follows:

- Concentrations of PBDEs in herring gull eggs have increased exponentially between 1981 and 2000 at Lake Ontario, Huron and Michigan sampling sites (see figure below).
- Concentrations of PBDEs (predominantly tetra- and pentaBDE congeners) have also increased exponentially between 1981 and 2000 in Arctic male ringed seals.¹³
- PBDEs have been detected in all environmental media as well as sewage sludge, and there is evidence that their levels in the North American environment are increasing.¹⁴

A summary of studies of PBDEs in the Great Lakes environment is presented in Appendix A.




PBDE in Herring Gull Eggs, Gull Island, Lake Michigan, 1981-2005. Source: Great Lakes Binational Toxics Strategy 2006 Annual Progress Report.

A spatial distribution of PBDEs in lake trout from the Great Lakes and temporal trends in lake trout in Lake Ontario along with biomagnification of PBDEs in Lake Ontario food web were accomplished under the supervision of Dave Sergeant and Mike Whittle. PBDE levels ranged between 22 ng/g-wet weight for Lake Erie and 84 ng/g-wet weight for Lake Ontario. The levels of PBDEs in lake trout from Lake Ontario increased from 0.12 ng/g-wet weight in 1978 to 148 ng/g-wet weight in 1998. Biomagnification factors for PBDEs in Lake Ontario food web ranged between 0.6 for transfer of BDE-99 from diporeia to sculpin, and 5.7 for transfer of BDE-100 from plankton to mysid.¹⁵

A more recent analysis indicates that concentrations of PBDEs in lake trout from Lake Ontario have increased exponentially from 0.54 ng/g in 1988 to 190 ng/g wet weight in whole fish samples collected in 2002 (Whittle et al., 2004). From archived lake trout samples from the five Great

Lakes, total PBDE concentrations increased exponentially with time, doubling every 3 to 4 years (see figure below).¹⁶



Temporal Trend of Total PBDE Concentrations in Fishes from the Great Lakes, 1975-2005.
Source: Great Lakes Binational Toxics Strategy 2006 Annual Progress Report.

In 2008, Venier and Hites reported that concentrations of PBDEs in air at IADN sites on the Great Lakes during 2005-2006 were highest at the urban sites in Chicago and Cleveland (65 ± 4 and 87 ± 8 pg/m^3 , respectively), and lowest at the remote site in Eagle Harbor (5.8 ± 0.4 pg/m^3). Concentrations of BDE-209 (decaBDE) were not decreasing at any of the five IADN sites analyzed. Concentrations of BDE-47 and BDE-99 in the atmosphere were decreasing rapidly, except at the Chicago site, with half-lives of ~ 2 years.¹⁷

Venier and Hites also estimated the atmospheric deposition of PBDEs into the Great Lakes during 2005-2006. Total PBDEs measured in precipitation in this study ranged from 94 ± 19 ng L^{-1} at Chicago to 0.65 ± 0.14 ng L^{-1} for the rural Sturgeon Point site. The study findings also suggested that Chicago may be a source of BDE-47 to Lake Michigan and that Cleveland may be a source of BDE-209 to Lake Erie.¹⁸

Source/Use/Release/Exposure Information

Source/Use

PBDEs are not manufactured in Canada. The DecaBDE commercial mixture is imported into Canada for addition to various intermediate and finished products, such as computer housings, household appliances, furniture, automotive/aircraft seating and interiors, and a variety of electrical and electronic components.¹⁹

The only United States manufacturer of PentaBDE and OctaBDE commercial mixtures voluntarily ceased production in December 2004.²⁰

Release

PBDEs are potentially released to the environment throughout their lifecycle, from the chemicals themselves as well as from products containing them. PBDEs may enter the environment through treated or untreated municipal or industrial wastewater discharges to surface water and also through leachate from landfills and municipal incineration when products and materials containing these substances are sent for final disposal. PBDEs may also be released directly to air, land and surface water when products containing PBDEs are manufactured and during their use.²¹

It is estimated that releases from the commercial mixture PentaBDE are primarily associated with in-service use of polyurethane foam and to a lesser extent disposal of the foam at the end of its lifecycle. The majority of releases from the use of the OctaBDE commercial mixture are estimated to be associated with product disposal and materials handling before resin compounding. Releases of the DecaBDE commercial mixture from plastics are mainly associated with product disposal; whereas releases of the DecaBDE commercial mixture associated with textile applications appear to be spread relatively evenly throughout the lifecycle – with most releases being associated with textile processing/finishing and releases during product service life. Releases of DecaBDE commercial mixture during service life of textiles would occur during washing and would be to wastewater.²²

Exposure:

It is difficult to derive meaningful estimates of exposure to individual congeners or congener groups. However, exposure is related to releases.²³

Pathways of exposure include pathways of exposure food, house dust, and indoor air. Research has suggested that house dust may be a substantial contributor to PBDE exposure. To further examine exposure pathway issues, EPA and the USDA are collaborating on a study of the bioavailability of PBDEs in house dust.²⁴

In a study of U.S. exposures to PBDEs, an adult intake dose of total PBDEs was estimated to be 7.7 ng/kg body weight/day, and children's estimated intakes were higher at 49.3 ng/kg/day for ages 1–5, 14.4 ng/kg/day for 6–11, and 9.1 ng/kg/day for 12–19. The much higher dose for the child aged 1–5 was due to the doubling of dust ingestion from 50 to 100 mg/day. Exposures to PBDEs in house dust accounted for 82% of the overall estimated intakes.²⁵ Huwe et al. 2008 found that PBDEs in household dust are readily bioavailable and that ingestion of PBDE-contaminated dust by rats leads to bioconcentration in the liver.²⁶

Concentrations of PBDEs in the blood of the general U.S. population were measured as part of the National Health and Nutrition Examination Survey (NHANES) 2003–2004. A total of 2,062

serum samples, from participants in NHANES aged 12 years and older, were analyzed for PBDEs. The highest serum concentration was found for 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) with a geometric mean of 20.5 ng/g lipid, followed by 2,2',4,4',5,5'-hexaBDE (BDE-153) [5.7 ng/g lipid], 2,2',4,4',5-pentaBDE (BDE-99) [5.0 ng/g lipid; a value equal to the highest limit of detection for an individual sample], 2,2',4,4',6-pentaBDE (BDE-100) [3.9 ng/g lipid], BB-153 [2.3 ng/g lipid], and 2,4,4'-triBDE (BDE-28) [1.2 ng/g lipid]. In examining those with the highest serum concentrations (above the 95th percentile) of BDE-47, age, but not sex and race/ethnicity, was significantly associated. Senior adults (60 years and older) were two times more likely to have serum concentrations of BDE-47 above the 95th percentile than adults aged 20–59 years and 54% more likely than adolescents to have serum BDE-47 concentrations above the 95th percentile.²⁷

Anderson et al. 2008 investigated serum concentrations of PBDEs in a cohort of 508 consumers of sport-caught fish living in five Great Lake states. BDE-47 was found in 98% of samples, and BDE-99 in 62% of samples. Total PBDE levels were positively associated with age, hours spent outdoors, years of sportfish consumption, and catfish and shellfish intake; other dietary components were not predictive of PBDE levels.²⁸

Spliethoff et al. 2008 examined PBDE body burdens among 36 sportfish consumers in New York State. The authors found that the number of years of Lake Ontario sportfish consumption between 1980 and 1990, after adjusting for plasma lipids, was a statistically significant predictor of total PBDE blood plasma levels. In addition, the number of meals eaten in the year prior to the study, of Lake Ontario sportfish species known to have high levels of other persistent organic pollutants (POPs), was correlated with total PBDE blood plasma levels.²⁹

Concentrations of PBDEs in human breast milk from across Canada ranged between 19.08 ng/g for samples collected from the Maritimes and 2.57 ng/g for samples collected in Ontario. Levels of PBDEs in breast milk rose from a mean of 0.2 ng/g for samples collected across Canada in 1982 to a mean of 42.5 ng/g (25.4 ng/g median) lipid weight for samples collected in the Vancouver area in 2001-2.³⁰ Schechter et al. 2005 analyzed PBDEs in breast milk in the U.S. The highest 5% of breast milk levels in the U.S. were 6 to 21 ng/g wet weight.³¹

Concentrations of PBDEs in a number of food basket items ranged between 0.04 ng/g for pasta and 1.2 ng/g for wieners; these values resulted in an estimated daily dietary intake of 44 ng, which was higher than that for PCDD/Fs (2.4 ng), and lower than that for PCBs (285 ng).³²

Environmental Benchmarks

There are no environmental quality benchmark criteria available from Canada and the United States.

Environmental and Health Data

Information on health data can be found from Health Canada's State of the Science Report for a Screening Health Assessment for PBDEs [Tetra, Penta, Hexa, Hepta, Octa, Nona, and Deca] (December 2004).³³

Chronic toxicity/carcinogenicity studies of BDE-99 (a component of commercial pentaBDE) are not available. There is “inadequate information to assess the carcinogenic potential” of BDE-99. However, the available animal data indicate that the nervous system is a sensitive target. Neurobehavioral developmental toxicity has been identified as the critical endpoint of concern in mice following pre- and neonatal oral exposure to BDE-99.³⁴

Herbstman found that BDEs 47, 99, and 100 were associated with lower mental development scores at age 4.³⁵

Johansson et al. 2008 reported that neonatal exposure to DecaBDE (BDE-209) causes dose-response changes in spontaneous behaviour and cholinergic susceptibility in adult mice, and that BDE-209 can be as potent as the lower brominated PBDEs in causing developmental neurotoxic defects.³⁶

An RfD of 0.1 µg/kg-day was calculated from a BMDL1SD of 0.32 mg/kg-day for effects on spontaneous motor behavior in mice. The database for BDE-99 is sparse for the derivation of an RfD: there are no standard reproductive, developmental, subchronic, or chronic studies in rats or mice nor a much needed developmental neurotoxicity study. In addition, there are several concerns regarding the experimental design of the Viberg et al. (2004a) study used in proposing an RfD. The overall confidence in the RfD assessment is low.³⁷

No information is available on the carcinogenicity of decaBDE in humans.³⁸ Chronic dietary studies of decaBDE have been conducted in rats and mice. The weight of evidence suggests that decaBDE shows “suggestive evidence of carcinogenic potential.”³⁹

In Europe the FIRE project (Flame retardants Integrated Risk assessment for Endocrine effects; see www.rivm.nl/fire) completed a 28-day toxicity study of decaBDE in rats. Van der Ven reported results of the 28-day toxicity study, which suggested that the endocrine effects of decaBDE may present a reproductive human health hazard. The study also reported that body burdens in children, with upper values reported as high as “233 ng/g lipid (Fischer et al., 2006) and 23 ng/g blood (Watson, 2005),” suggest that the need for further risk reduction measures be considered.⁴⁰

In addition, a separate study of decaBDE developmental neurotoxicity in mice has been published (Rice et al., *Neurotoxicology and Teratology* 2007).⁴¹

Other Reasons for Concern

Evidence suggests that PBDEs have endocrine disrupting properties.⁴² Toxicity data indicate potential for thyroid hormone effects (inhibition of tail resorption in tadpoles exposed to a commercial PBDE mixture).⁴³ Turyk et al. 2008 linked PBDE exposure with changes in thyroid antibodies and thyroid hormone homeostasis in men, reporting that “PBDE exposure, at levels comparable to the general US population, was associated with increased thyroglobulin antibodies and increased thyroxine in adult males.”⁴⁴

The animal model indicates a potential for concern for early lifetime exposure (i.e., fetal or infant exposure) to the chemical. The identification of BDE-99 in human maternal and cord serum, milk,

and children's serum (Mazdai et al., 2003; Schechter et al., 2003; Thomsen et al., 2002) implies humans are exposed to BDE-99 during a period of rapid development of the brain, a critical window of development, indicating a potential for susceptibility. Whether such exposure constitutes a health risk for adverse neurodevelopmental effects in children is not known at this time because of the limited toxicological data base for BDE-99. An association between prenatal or neonatal exposures to BDE-99 and neurobehavioral dysfunction in humans has not been established.⁴⁵

Riu et al. 2008 found that DecaBDE in late gestation rats is absorbed and reaches the fetus, where it debrominates into OctDBE and other metabolites.⁴⁶ These findings suggest potential concern for human fetal exposures to DecaBDE.

III. PRESENT MANAGEMENT STATUS

Canada

Polybrominated Diphenyl Ethers Regulations (PBDE Regulations) were published in the *Canada Gazette*, Part II, on July 9, 2008. The Regulations prohibit the manufacture of PBDEs (i.e. tetra-, penta-, hexa-, hepta-, octa-, nona-, and decaPBDE congener groups) in Canada. These regulations also prohibit the manufacture, use, sale, offer for sale and import of mixtures, polymers and resins containing those PBDEs that meet the criteria for virtual elimination (tetra-, penta- and hexaBDE congener groups) under the *Canadian Environmental Protection Act, 1999 (CEPA, 1999)*. More information on this regulation can be found at:

http://www.ec.gc.ca/ceparegistry/documents/regs/g2-14214_r1.pdf.

United States

The EPA has issued a Significant New Use Rule (SNUR) that will require the notification of, and evaluation by, the USEPA of any new use of PentaBDE or OctaBDE commercial mixtures (71 FR 34015 Available at: <http://www.epa.gov/oppt/pbde/>).⁴⁷

Trends in releases of decabromodiphenyl oxide (decaBDE) are being followed by the US through the Toxics Release Inventory.

In March 2006, EPA prepared a Project Plan for PBDEs, which outlined EPA's activities regarding PBDEs. Since 2006, EPA has prepared an updated status report on PBDEs annually. EPA's Project Plan and latest status report (March 2008) are available at <http://www.epa.gov/opptintr/pbde/>. Highlights of the March 2008 status report are presented below.

EPA has formed a Furniture Flame Retardancy Partnership with several industry associations and other stakeholders. The partnership has conducted a screening level hazard assessment of flame retardant chemicals that may be suitable substitutes for pentaBDE. The summary assessment is available at <http://www.epa.gov/opptintr/dfe/pubs/projects/flameret/index.htm>.

The U.S. Consumer Product Safety Commission (CPSC) has proposed a new mandatory standard to address residential upholstered furniture fires. Under the proposal, manufacturers could meet the performance standard by using smolder-resistant cover fabrics or interior fire resistant barriers

to protect the furniture's internal filling material, without requiring the use of fire retardant chemicals.

EPA is considering proposing a SNUR under the Toxic Substances Control Act (TSCA) for flame retardants, which would include decaBDE among 16 chemical substances/categories. The proposed SNUR is on hold at EPA pending the CPSC's decision on a new mandatory standard for residential upholstered furniture flammability.

The industry sponsor of decaBDE in the Voluntary Children's Chemical Evaluation Program (VCCEP) committed to conducting a Tier 2 VCCEP assessment for decaBDE in December 2005. New documents will be posted at <http://www.epa.gov/oppt/vccep/pubs/chem21.htm>.⁴⁸

A program called the Electronic Product Environmental Assessment Tool (EPEAT) was created by a multi-stakeholder group in the US. EPEAT is a procurement tool used to identify desktop computers, notebooks and monitors that satisfy established environmental criteria, including conformance to the European RoHS (i.e. products that may have PBDEs).⁴⁹

RECOMMENDATION

To be determined.

APPENDIX A

Brominated Flame Retardants in the Great Lakes Environment - Published Studies

1. "Concentrations and spatial variations of polybrominated diphenyl ethers and other organohalogen compounds in Great Lakes air." B. Strandberg, N. G. Dodder, I. Basu and R. A. Hites. *Environmental Science & Technology*. 2001; 35(6): 1078-1083. PBDEs\Strandberg et. al. 2001.pdf

Summary: This study measured PBDEs in air around the Great Lakes and compares it to well studied compounds such as PCBs and organochlorine pesticides. PBDEs were found in every air sample indicating these compounds are prevalent and well distributed and dispersed in Great Lakes air. Concentrations of PBDEs were similar to that of pesticides like DDT. Spatial trends were similar to PCBs, indicating that gas-particle partitioning is important in understanding the fate of PBDEs.

2. "Geographical Distribution (2000) and Temporal Trends (1981-2000) of Brominated Diphenyl Ethers in Great Lakes Herring Gull Eggs." R.J. Norstrom, M. Simon, J. Moisey, B. Wakeford, and D.V. Chip Weseloh. *Environmental Science & Technology*. 2002; 36: 4786-4789. PBDEs \Norstrom et.al. 2002.pdf

Summary: This study measured the concentrations of PBDEs in Herring Gull eggs from 1981-2000, and showed distribution of PBDEs in the Great Lakes for the year 2000. Highest concentrations were found in northern Lake Michigan and Toronto harbor (1000-1400 mg/kg) and

lowest in Lake Huron and Lake Erie (192-340 mg/kg). The distribution suggested that input from large urban/industrial areas through air or water emissions contributes local contamination to the herring gull food web. Temporal trends for pentaPBDEs showed an exponential increase. Doubling times for pentaPBDEs were calculated to be 2.6 years in Lake Michigan, 3.1 years in Lake Huron and 2.8 years in Lake Ontario.

3. "Time Trends (1983-1999) for Organochlorines and Polybrominated Diphenyl Ethers in Rainbow Smelt (*Osmerus Mordax*) from Lakes Michigan, Huron, and Superior, USA". S.M. Chernyak, C.P. Rice, R.T. Quintal, L.J. Begnoche, J.P. Hickney, and B.T. Vunyard. *Environmental Toxicology and Chemistry*. 2005; 24,(7):1632-1641. PBDEs\Chernyak et al 2005.pdf

Summary: The USGS Great Lakes Center collected fish from the early 1980's until today, this study measured organochlorines and PBDEs in the fish to understand contaminant trends in the Great Lakes. Concentrations of PBDEs were traced from its entry into the ecosystem in 1980 until 1999, and trends showed PBDE concentrations increasing exponentially at every site with doubling times estimated at 1.58 to 2.94 years.

4. "Dramatic Changes in the Temporal Trends of Polybrominated Diphenyl Ethers (PBDEs) in Herring Gull Eggs from the Laurentian Great Lakes: 1982-2006." L.T. Gauthier, C.E. Herbert, D.V. Chip Weseloh, and R.J. Letcher. *Environmental Science & Technology*. 2008; 42(5); 1524-1530. PBDEs\Gauthier et.al. 2008.pdf

Summary: This study measured 43 PBDEs, specifically decaPBDE (209) in Herring Gull eggs from all five Great Lakes. The total PBDE concentration for lower Brominated congeners in the gulls was in the same range as the concentrations for octa and nona PBDEs, indicating that the higher Brominated congeners are accumulating at higher rates than lower brominated ones. PBDE-209 and its products showed increases in concentration over time. Lower Brominated congeners showed rapid increase in the eggs until 2000. Over all trends in pentaPBDEs show a decrease in Lakes Superior, Michigan, and Ontario from 2000-2006, while measurements at Lake Erie showed an increase. Lake Huron had two sites, one showed an overall increase of pentaPBDEs, and the other showed an overall decrease in the concentrations over time. Measurements of PBDE 207 and 197 were used as indicators of metabolic debromination from parent birds to the eggs.

5. "Spatial distribution and temporal trends in PBBs and PBDEs in lake trout from the Great Lakes." J. M. Luross, C. M. Cannon, M. Alae, D. B. Sergeant, D. M. Whittle and K. R. Solomon. *International Association for Great Lakes Research 44th Conference*, Green Bay, WI. 2001.

6. "Comparison of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) in Lake Michigan salmonids." J. B. Manchester-Neesvig, K. Valters and W. C. Sonzogni. *Environmental Science & Technology*. 2001; 35(6): 1072-1077. PBDEs \Manchester-Neesvig et. al. 2001.pdf

Summary: PCBs and PBDEs were measured in 21 coho and Chinook salmon taken from Lake Michigan tributaries in 1996 for this study. The study found PBDEs to be found in fish at one-

third the concentration of PCBs, but still at the highest reported level for fish worldwide. Implications are that PBDEs are as prevalent in the Lake Michigan water column as PCBs.

7. "Brominated flame retardants in the Great Lakes" Persistent Organic Pollutants in the Great Lakes, Handbook in Environmental Chemistry, Vol. 5, Part N, pp. 328-366, Springer-Verlag: Berlin & Heidelberg, Germany (2006), R. A. Hites, editor.

8. "Identification of brominated carbazoles in sediment cores from Lake Michigan." L. Y. Zhu and R.A. Hites. Environmental Science & Technology. 2005; 39, 9446-9451. Brominated carbazoles\Zhu 2005.pdf

Summary: This study measured a possible naturally occurring brominated organic a brominated carbazole. The time trend determined by measuring these compounds as a function of depth in the Lake Michigan sediment cores indicated that the concentration of 1,3,6,8-tetrabromocarbazole peaked around 1920-1935. This was the first time that bromocarbazoles have been found in the environment.

9. "Deposition versus Photochemical Removal of PBDEs from Lake Superior Air." J.D. Raff and R.A. Hites. Environmental Science & Technology. 2007; 41: 6725-6731. PBDEs\Raff & Hites 2007.pdf

Summary: This study measured PDBEs from sediment on a remote Lake Superior island as well PBDEs in Lake Superior air. The study show decrease in PBDEs concentrations from air to sediment, except PBDE 209. Study concludes that photolysis is an important process for PBDE degradation in air and could be one reason that PBDE congener profiles in sediment are highly variable.

10. "Brominated flame retardants in sediment cores from Lakes Michigan and Erie" L. Y. Zhu and R.A. Hites. Environmental Science & Technology. 2005; 39, 3488-3494; erratum 39, 5904 (2005). PBDEs\Zhu & Hites 2005 Lake Erie and Michigan.pdf

Summary: Dated sediment cores were studied to see trends in PBDEs and PBBs (Polybrominated Biphenyls). The study showed higher concentrations in Lake Michigan than Lake Erie, and Lake Superior. The study also showed rapid increase in both PBDEs and PBBs in sediments, concurrent with market increase in demand for these flame retardants. The total burdens of these compounds in the sediment of Lakes Michigan and Erie were 110 and 10 metric tons, respectively. The estimated total burden of these compounds in all of the Great Lakes approximately 200 tons.

11. "Temporal trends and spatial distributions of brominated flame retardants in archived fishes from the Great Lakes." L. Y. Zhu and R.A. Hites. Environmental Science & Technology. 2004; 38: 2779-2784. PBDEs\Zhu & Hites 2004.pdf

Summary: PBDEs and PBBs were measured in fish from all five Great Lakes. Spatial trends showed Lake Michigan and Ontario had the highest total concentrations of both PBDEs.

Concentrations in all fish increased exponentially over time with a calculated doubling rate of 3-4 years. Concentrations of PBBs, banned in the 1970's, remained at a constant with only a slight decrease in fish from Lake Huron.

12. "Polybrominated diphenyl ethers in the sediments of the Great Lakes. 3. Lakes Ontario and Erie." W. Song, J.C. Ford, A. Li, N.C. Sturchio, K.J. Rockne, D.R. Buckley, and W.J. Mills. *Environmental Science & Technology*. 2005; 39,(15):5600-5. PBDEs\Song et. al. 2005 Lakes Ontario and Erie.pdf,

Summary: Sediment cores from both Lake Ontario and Erie were analyzed for PBDE and PCB concentrations Lake Ontario. Results showed a dramatic increase in PDBE concentrations, while PCB concentrations peaked and then showed slow decrease. Lake Erie sediments showed no temporal trends due to high rates of sediment mixing. Overall congener profiles of PBDEs showed that PBDE-209 is the highest single congener concentration making up 90-95% of total PBDE concentration.

13. "Polybrominated diphenyl ethers in the sediments of the Great Lakes. 2. Lakes Michigan and Huron." W. Song, A. Li, J.C. Ford, N.C. Sturchio, K.J. Rockne, D.R. Buckley, and W.J. Mills.. *Environmental Science & Technology*. 2005; 39, (10):3474-9. PBDEs\Song et.al. 2005 Lakes Michigan & Huron.pdf

Summary: PBDEs and PCBs were measured in sediments from both Lake Huron and Lake Michigan. PBDE concentrations increase over time at all sampling sites. Spatial distribution and PBDE congener profiles implied that both lakes concentrations appeared to be dependent upon latitude and the sites proximity to populated areas. It was suggested that north-bound air plumes from urban areas are the major sources of PBDEs found in the lake sediments at locations away from the shores.

14. "Polybrominated diphenyl ethers in the sediments of the Great Lakes. 1. Lake Superior." W. Song, J.C. Ford, A. Li, W.J. Mills, K.J. Rockne, and D.R. Buckley. *Environmental Science & Technology*, 2004; 38(12):3286-93. PBDEs\Song et. al. 2004 Lake Superior.pdf

Summary: Measurements of PBDE and PCB concentrations in Lake Superior showed that PCBs concentrations in sediments are slowly declining, while PBDEs are increasing with time. The load of total PBDEs to Lake Superior was estimated to be 2-6 metric tons, and the calculated loading rate was 80-160 kg per year.

15. "Polybrominated diphenyl ethers in the sediments of the Great Lakes. 4. Influencing factors, trends, and implications." A. Li, W. Song, J.C. Ford, N.C. Sturchio, K.J. Rockne, D.R. Buckley, and W.J. Mills. *Environmental Science & Technology*. 2006 Dec 15; 40 (24):7528-34. PBDEs\Li et. al. 2006.pdf

Summary: Sediment cores from all five Great Lakes were analyzed for PBDEs. Spatial trends indicated that major individual PBDE congeners showed strong dependence on the latitude of sampling sites. The study suggested that data reflected both the influence of urbanization, which showed south-to-north gradient in the region, and the general direction of long-range transport of

airborne pollutants in the northern hemisphere. Doubling times of PBDEs in sediment showed a longer time than reported doubling times for humans or fish, indicating a slower response in sediments to emissions of PBDEs. Regressions applied to the data showed that latitude, year of deposition, and organic matter were all independent variables to PBDE concentration.

16. "Partitioning and Bioaccumulation of PBDEs and PCBs in Lake Michigan". S. Streets, S. Henderson, A. Stoner, D. Carlson, M. Simcik, and D. Swackhamer. *Environmental Science & Technology*. 2006; 40: 7263-7269. PBDEs\Streets et al 2006.pdf

Summary: Water samples from Lake Michigan were analyzed for PBDEs and PCBs. Concentrations of PBDEs were similar to PCBs in the water column, and data suggested that partitioning for PBDEs between particulates and dissolved phase behaves similar to PCBs partitioning. Fish from all the Great Lakes were also analyzed for PCBs and PBDEs, specifically BDE-99. PCBs and PBDEs did not follow the same trend of accumulation in fish, and this paper suggested that biotransformation of BDE-99 are occurring.

17. "Polybrominated Diphenyl Ethers and Hydroxylated and Methoxylated Brominated and Chlorinated Analogues in the Plasma of Fish from the Detroit River." K. Valters, H. Li, M. Alaei, I. D'Sa, G. Marsh, A. Bergman, and R.J. Letcher. *Environmental Science & Technology*. 2005; 39: 5612-5619. PBDEs\Valters et.al 2005.pdf

Summary: This study measured concentrations of PBDEs and their metabolites-hydroxylated PBDEs (OH-PBDEs), methoxylated-PBDEs (MeO-PBDEs), as well as the antimicrobial triclosan and its metabolites-OH-trichlorodiphenyl ether, and methylated (MeO) triclosan in fish from the Detroit River. The study found that the sum of PBDE to sum of OH-PBDE concentration ratios ranging from 0.0005 to 0.02. OH-PBDEs are likely derived in these freshwater species as metabolites of precursor PBDEs and are subsequently retained in the blood, although they could have external origins and be picked up and accumulated by fish. Anthropogenic triclosan concentrations ranged from 750 to >10,000 pg/g wet weight and was clearly a bioaccumulative halogenated phenolic compound in these fish. MeO-triclosan concentrations were considerably lower.

18. "Spatial distribution of polybrominated diphenyl ethers and polybrominated biphenyls in lake trout from the Laurentian Great Lakes." J. M. Luross, M. Alaei, D. B. Sergeant, C. M. Cannon, D. D.M. Whittle, K. R. Solomon and D. C. G. Muir. *Chemosphere*. 2002; 46, (5):665-672. PBDEs\Luross et.al. 2002.pdf

Summary: This study measured PBDE and PBB concentrations in trout from all five Great Lakes. Concentrations were highest at Lake Ontario and lowest at Erie for PBDEs. Lake Huron had the highest concentrations of PBBs, with the lowest concentrations being at Lake Superior. This study also found that the three most prominent PBDEs (47, 99,100) in fish are all components of the commonly used penta-BDE found in flame retardants.

Note: Appendix A compiled by Carolyn Peerson, University of Iowa, for International Joint Commission Chemicals of Emerging Concern Workgroup, July, 2008.

REFERENCES

- ¹ Environment Canada. *Management of Toxic Substances- Substance Detail- Polybrominated Diphenyl Ethers (PBDEs)*. Available online at: http://www.ec.gc.ca/TOXICS/EN/detail.cfm?par_substanceID=201&par_actn=s1.
- ² *Information from Suzanne Easton* (2008).
- ³ <http://www.epa.gov/HPV/pubs/update/spnchems.htm>
- ⁴ Environment Canada- Chemicals Sector Division, Environmental Stewardship Branch. December 2006. *Risk Management Strategy for Polybrominated Diphenyl Ethers (PBDEs) – tetra, penta, hexa, hepta, octa, nona, deca*. Available online at: http://www.ec.gc.ca/Toxics/docs/substances/PBDE/PBDE_RMS/EN/toc.cfm. To: *Excel file: EPA Report Table_SRC* (Muir, D.C., Howard, P.H. and Meylan, W. 2007. Project: Screening Chemicals in Commerce to Identify Possible Persistent and Bioaccumulative (P&B) Chemicals in the Great Lakes.)
- ⁵ Environment Canada - Chemicals Sector Division, Environmental Stewardship Branch. December 2006. *Risk Management Strategy for Polybrominated Diphenyl Ethers (PBDEs) – tetra, penta, hexa, hepta, octa, nona, deca*. Available online at: http://www.ec.gc.ca/Toxics/docs/substances/PBDE/PBDE_RMS/EN/toc.cfm.
- ⁶ Ibid.
- ⁷ Ibid.
- ⁸ Ibid.
- ⁹ Ibid.
- ¹⁰ Ibid.
- ¹¹ Ibid.
- ¹² Environment Canada. June 2006. *Canadian Environmental Protection Act, 1999 - Ecological Screening Assessment Report on Polybrominated Diphenyl Ethers (PBDEs)*. Available online at: [http://www.ec.gc.ca/CEPARRegistry/documents/subs_list/PBDE_SAR/PBDEs_SAR_EC_June_2006_\(en\).pdf](http://www.ec.gc.ca/CEPARRegistry/documents/subs_list/PBDE_SAR/PBDEs_SAR_EC_June_2006_(en).pdf).
- ¹³ Ibid.
- ¹⁴ Ibid.
- ¹⁵ Health Canada. 2007. *Impact of Polybrominated Diphenyl Ethers on Canadian Environment and Health of Canadians*. Available at <http://www.hc-sc.gc.ca/sr-sr/finance/tsri-irst/proj/persist-org/tsri-237-eng.php>.
- ¹⁶ Great Lakes Binational Toxics Strategy 2006 Annual Progress Report. Available at www.binational.net.
- ¹⁷ Venier, M. and R.A. Hites. In press (a). *Environ. Sci. Technol.*
- ¹⁸ Venier, M. and R.A. Hites. In press (b). *Environ. Sci. Technol.*
- ¹⁹ Environment Canada. *Management of Toxic Substances- Substance Detail- Polybrominated Diphenyl Ethers (PBDEs)*. Available online at: http://www.ec.gc.ca/TOXICS/EN/detail.cfm?par_substanceID=201&par_actn=s1.
- ²⁰ Health Canada. December 2004. *State of the Science Report for a Screening Health Assessment- Polybrominated Diphenyl Ethers (PBDEs) [Tetra-, Penta- Hexa-, Hepta- Octa-, Nona- and Deca- Congeners]*. Available online at: http://www.ec.gc.ca/CEPARRegistry/documents/subs_list/PBDE_SAR/HC_SOS_PBDE_e.pdf.

-
- ²¹ Environment Canada- Chemicals Sector Division, Environmental Stewardship Branch. December 2006. *Risk Management Strategy for Polybrominated Diphenyl Ethers (PBDEs) – tetra, penta, hexa, hepta, octa, nona, deca*. Available online at: http://www.ec.gc.ca/Toxics/docs/substances/PBDE/PBDE_RMS/EN/toc.cfm.
- ²² Ibid.
- ²³ Ibid.
- ²⁴ United States Environmental Protection Agency Polybrominated Diphenyl Ethers Workgroup. 2008. Tracking Progress on U.S. EPA’s Polybrominated Diphenyl Ethers (PBDEs) Project Plan: Status Report on Key Activities. Available at <http://www.epa.gov/opptintr/pbde/pubs/pbdestatusreport.pdf>.
- ²⁵ Lorber, M. *Journal of Exposure Science and Environmental Epidemiology* (2008) 18, 2–19; doi:10.1038/sj.jes.7500572; published online 11 April 2007
- ²⁶ Huwe, J.K., Hakk, H., Smith, D.J., DiLiberto, J.J., Richardson, V., Stapleton, H.M., Birnbaum, L.S. 2008. *Environ. Sci. Technol.* 42, 2694–2700.
- ²⁷ Sjöden, A., Wong, L-Y., Jones, R.S., Park, A., Zhang, Y., Hodge, C., DiPietro, E., McClure, C., Turner, W., Needham, L., Patterson, D.G. 2008. *Environ. Sci. Technol.* 42, 1377–1384.
- ²⁸ Anderson, H.A., Imm, P., Knobeloch, L., Turyk, M., Mathew, J., Buelow, C., Persky, V. 2008. *Chemosphere* 73, 187–194.
- ²⁹ Spliethoff, H.M., Bloom, M.S., Vena, J., Sorce, J., Aldous, K.M., Eadon, G. In press. Exploratory assessment of sportfish consumption and polybrominated diphenyl ether exposure in New York State anglers. *Environmental Research*.
- ³⁰ Health Canada. 2007. Impact of Polybrominated Diphenyl Ethers on Canadian Environment and Health of Canadians. Available at <http://www.hc-sc.gc.ca/sr-sr/finance/tsri-irst/proj/persist-org/tsri-237-eng.php>.
- ³¹ Schechter, A.; Paepke, O.; Tung, K. C.; Joseph, J.; Harris, T. R.; Dahlgren, J. Polybrominated diphenyl ether flame retardants in the U.S. population: Current levels, temporal trends, and comparison with dioxins, dibenzofurans, and polychlorinated biphenyls. *J. Occup. Environ. Med.* 2005, 47, 199–211.
- ³² Ibid.
- ³³ Health Canada. December 2004. *State of the Science Report for a Screening Health Assessment- Polybrominated Diphenyl Ethers (PBDEs) [Tetra-, Penta- Hexa-, Hepta- Octa-, Nona- and Deca- Congeners]*. Available online at: http://www.ec.gc.ca/CEPARRegistry/documents/subs_list/PBDE_SAR/HC_SOS_PBDE_e.pdf.
- ³⁴ U.S. EPA. IRIS Toxicological Review of Pentabromodiphenyl Ether (External Review Draft). U.S. Environmental Protection Agency, Washington, D.C.
- ³⁵ Herbstman, unpublished.
- ³⁶ Johansson, N., Viberg, H., Fredriksson, A., Eriksson, P. *Neurotoxicology*. 2008 Sep 27.
- ³⁷ Ibid.
- ³⁸ U.S. EPA. IRIS Toxicological Review of Decabromodiphenyl Ether (External Review Draft).
- ³⁹ Ibid.

⁴⁰ Van der Ven, L.T.M., van de Kuil, T., Leonards, P.E.G., Slob, W., Cantón, R.F., Germer, S., Visser, T.J., Litens, S., Håkansson, H., Shrenk, D., van den Berg, M., Piersma, A.H., Vos, J.G., Opperhuizen, A. 2008. Toxicology Letters 179, 6–14.

⁴¹ United States Environmental Protection Agency Polybrominated Diphenyl Ethers Workgroup. 2008. Tracking Progress on U.S. EPA's Polybrominated Diphenyl Ethers (PBDEs) Project Plan: Status Report on Key Activities. Available at <http://www.epa.gov/opptintr/pbde/pubs/pbdestatusreport.pdf>.

⁴² Canadian Cancer Society. *Polybrominated diphenyl ethers (PBDEs)*. Available online at: http://www.cancer.ca/ccs/internet/standard/0,3182,3172_1438644643_langId-en,00.html.

⁴³ Health Canada. 2007. Impact of Polybrominated Diphenyl Ethers on Canadian Environment and Health of Canadians. Available at <http://www.hc-sc.gc.ca/sr-sr/finance/tsri-irst/proj/persist-org/tsri-237-eng.php>.

⁴⁴ Turyk, M.E., Persky, V.W., Imm, P., Knobeloch, L., Chatterton, R., Anderson, H.A. 2008. Hormone Disruption by PBDEs in Adult Male Sport Fish Consumers. Environ Health Perspect. doi:10.1289/ehp.11707. Available at <http://dx.doi.org/> [Online 24 July 2008].

⁴⁵ U.S. EPA. IRIS Toxicological Review of Pentabromodiphenyl Ether (External Review Draft). U.S. Environmental Protection Agency, Washington, D.C.

⁴⁶ Riu, A., Cravedi, J-P., Debrauwer, L., Garcia, A., Canlet, C., Jouanin, I., Zalko, D. Disposition and metabolic profiling of [14C]-Decabromodiphenyl ether in pregnant Wistar rats. Environment International. Volume 34, Issue 3, April 2008, Pages 318-329. Available at http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6V7X-4NMV02P-2&_user=573921&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_acct=C000029198&_version=1&_urlVersion=0&_userid=573921&md5=07153aa680ad5a0d10a299d4e8f670.

⁴⁷ Environment Canada- Chemicals Sector Division, Environmental Stewardship Branch. December 2006. *Risk Management Strategy for Polybrominated Diphenyl Ethers (PBDEs) – tetra, penta, hexa, hepta, octa, nona, deca*. Available online at: http://www.ec.gc.ca/Toxics/docs/substances/PBDE/PBDE_RMS/EN/toc.cfm.

⁴⁸ United States Environmental Protection Agency Polybrominated Diphenyl Ethers Workgroup. 2008. Tracking Progress on U.S. EPA's Polybrominated Diphenyl Ethers (PBDEs) Project Plan: Status Report on Key Activities. Available at <http://www.epa.gov/opptintr/pbde/pubs/pbdestatusreport.pdf>.

⁴⁹ Environment Canada- Chemicals Sector Division, Environmental Stewardship Branch. December 2006. *Risk Management Strategy for Polybrominated Diphenyl Ethers (PBDEs) – tetra, penta, hexa, hepta, octa, nona, deca*. Available online at: http://www.ec.gc.ca/Toxics/docs/substances/PBDE/PBDE_RMS/EN/toc.cfm.