

US EPA ARCHIVE DOCUMENT

Presented at

Great Rivers Reference Condition Workshop

January 10-11, Cincinnati, OH

Sponsored by

The U.S. Environmental Protection Agency and The Council of State Governments



EMAP
Great River Ecosystems

Potential Uncertainties in Great River Biocriteria: An example from the rehabilitation of the Kennebec River (Maine)



Andrew F. Casper , James H. Thorp, Susan P. Davies and David L. Courtemanch.




Workshop Goals:

- developing reference biocriteria
- defining reference conditions / sites

Goals for this Presentation:

- Do not focus on impairments alone (e.g. rehabilitation)
- Assessment / Management are situation specific (i.e. they influences choice of reference site/conditions as well as biocriteria)

A River Reborn



Benefits for People and
Wildlife of the Kennebec River
Following the Removal
of Edwards Dam

THE KENNEBEC COALITION



1 Backhoe scoops out gravel packed behind the dam

gravel gone, the water flows

ENVIRONMENT

A River Runs Through It

75,000 dams were built. Now a few are coming down.

salmon migrating to sea. As a result, the Federal Energy Regulatory Commission (FERC) is refusing to relicense dams where the environmental costs outweigh the value of the hydropower, or demanding that a dam be retrofitted with fish ladders. That's often so expensive that the owner opts to

Dam being destroyed as a preservation act

By Glenn Adams
ASSOCIATED PRESS

AUGUSTA, Maine—As church bells pealed Thursday, a torrent of water was unleashed through a manmade

Gov. Angus King started the process by ringing a bell. More bells ringing from a church signaled a big backhoe to rip into a temporary dam.



THE NEW YORK TIMES NATION

National News Briefs

Dam Removal Project Is Successful in Maine

AUGUSTA, Me., Nov. 6 (AP) — To the delight of conservation groups and fishermen, Atlantic salmon have returned to the waters above the site of a 162-year-old dam that was torn down last month to allow fish to swim upstream.

Striped bass have come back, too, in such numbers that fly fishermen are having a field day on the

upper Kennebec River. The striped bass have been feeding on another species that swims upstream from the sea, alewives.

The removal of the 24-foot-high Edwards Dam has completed the rebirth of what was historically such a rich fishery that early settlers grew weary of eating fish.

The dam was built in 1837 to supply power to mills that rose along the river's banks. It was about 40 miles upstream from the Atlantic and

stretched 917 feet across the Kennebec, blocking salmon, shad, herring and other fish from reaching their spawning grounds upstream.

In 1997, the Federal Energy Regulatory Commission decided the dam should be torn down for the good of the environment. It became the first hydroelectric dam in the country ordered removed by the federal government against its owners' wishes.

Fearing His Father Dead, Boy Shoots Himself

YUCATAN, Minn., Nov. 6 (AP) — A teenager who mistakenly thought he had killed his father in a hunting

acc...
of f...
7...
his...
hos...
crit...
sta...
M...
a gap from...
m...
get the or...
though the...
The Feder...
Commission...
Dam's own...
removal and...
1...
But in a...
some enviro...
Works and...
agreed to s...
ting up m...

Local Resource Management Situation:

Hydropower usefulness declines

Maintenance/Safety expense rises

+ Sport-fishing/boating/tourism rising

Expectations from [Edwards] Dam Removal:

Increased biodiversity & productivity

Does dam removal work ?

Conditional Yes,

[based on benthic biomonitoring in smaller rivers]

- **Upstream Migration–Connectivity of fish improves**
- **Downstream near-term sedimentation increases**




Photos USFWS & Stanley & Doyle




Edwards Dam / Kennebec River Assessment:

Three levels of resolution:



- Community similarity
- Dominant taxa
- “Most responsive” taxa




Community Similarity (Pre-Removal)



Community Similarity (Post-Removal)



Community Similarity (more resolution):



Dominant Taxa (high relative abundance):

Pre-Removal

Restored
Reach

Reference
Flowing

Hyallela

Hyallela

Stenonema

Helopelopia

Chimarra

Microtendipes

Neureclipsis

Tribelos

Cheumatopsyche

Chimarra

Dominant Taxa (high relative abundance):

Pre-Removal

Restored
Reach

Reference
Flowing

Hyallela

Hyallela

Stenonema

Helopelopia

Chimarra

Microtendipes

Neureclipsis

Tribelos

Cheumatopsyche

Chimarra

Post-Removal

Restored
Reach

Reference
Flowing

Chimarra

Chimarra

Neureclipsis

Rheotanytarsus

Rheotanytarsus

Neureclipsis

Microtendipes

Hydropysche

Stenonema

Polypedilum

“Most Responsive” Taxa (low relative abundance):

Compare Restored Reach...

Pre- (1999) versus Gained 8 unique to restored reach
Post-Removal (2000) (6 lotic specialists or facultative), Lost 12

Take Home Message for Assessing Dam Removal:

“Pollution” sampling frameworks works


BUT it would be nice to have better Dam[n] biocriteria

(i.e. are biocriteria *AND* reference

impairment/improvement specific?)

Environmental History & biocriteria development?

-Many catchment parameters have changed since the times of historic reference



For Example:

- there has been considerable climate variability,
there will be more climate variability
- The presence of alien or non-native species in most river systems has increased over time.

