US ERA ARCHIVE DOCUMENT

Estimating the Condition of Gulf of Mexico Estuaries: National Coastal Assessment and National Estuary Program

Virginia Engle
USEPA/ORD/NHEERL
Gulf Ecology Division
EMAP Symposium 2004

Program Goals

- National Coastal Assessment
 - Assess condition of U.S. coastal resources
 - Create an integrated, comprehensive coastal monitoring program among the coastal states.
- National Estuary Program
 - Improve quality of estuaries of national importance
 - Attain or maintain water quality in estuaries by developing Comprehensive Conservation and Management Plans.












Estuarine Assessments






Surface Area of Gulf of Mexico Estuaries





Gulf of Mexico Estuarine Drainage Areas Land Cover Estimates





Methods

- Probability-based survey designs
- Data collected by states using standardized field, lab, & QA protocols
- Data analysis
 - Cumulative distribution functions (CDFs) with confidence intervals
 - Maps and pie charts showing distribution of good, fair, poor conditions



Number of Stations

NCA	2000
Florida	23
Alabama	50
Mississippi	35
Louisiana	38
Texas	45

NEP	2000	2001	2002
Charlotte Harbor			30
Sarasota Bay		20	
Tampa Bay	25		
Mobile Bay	33	33	
Barataria-Terrebonne	11	14	
Galveston Bay	12	16	
Coastal Bend Bays	13	14	



Indicators

Water Quality

Sediment Quality

DIN

DIP

Chlorophyll a

Water Clarity

DO

Metals

Organics

TOC

Toxicity

Biological Quality

Benthic Index

Fish Tissue Contaminants





NCA Water Quality Assessment Criteria

Indicator	Region	Good	Fair	Poor
DIN (mg/L)	Gulf	< 0.1	0.1 – 0.5	> 0.5
Dily (mg/L)	FL Bay	< 0.05	0.05 - 0.1	> 0.1
DIR (mg/L)	Gulf	< 0.01	0.01 - 0.05	> 0.05
DIP (mg/L)	FL Bay	< 0.005	0.005 - 0.01	> 0.01
Chl <i>a</i> (μg/L)	Gulf	< 5	5 – 20	> 20
FL Bay < 0.5	< 0.5	0.5 – 1	> 1	
WCI	Gulf	> 2	1 – 2	< 1
DO (mg/L)	Gulf	> 5	2 – 5	< 2




Dissolved Inorganic Nitrogen









Dissolved Inorganic Phosphorus









Dissolved Inorganic Phosphorus








Chlorophyll a








Chlorophyll a








Water Clarity








Water Clarity








Dissolved Oxygen









Water Quality Index

NCA 2000


NEP 2000-2002









Benthic Index








Benthic Index









NCA Sediment Quality Assessment Criteria

Indicator	Good	Fair	Poor
Ampelisca Survival (%)	= 80		< 80
Number of Contaminants	0 > ERM < 5 > ERL	= 5 > ERL	= 1 > ERM
TOC (%)	< 2	2 – 5	> 5




Sediment Toxicity









Sediment Toxicity









Sediment Contaminants









Total Organic Carbon










Sediment Quality Index











NCA Regional Scoring Criteria

Indicator	Good	Fair	Poor
DIN, DIP, Water Clarity	< 10% Poor and > 50% Good	10-25% Poor or > 50% Fair + Poor	> 25% Poor
CHL, WQI, Benthic Index	< 10% Poor and > 50% Good	10-20% Poor or > 50% Fair + Poor	> 20% Poor
Contaminants, DO, SQI	< 5% Poor and > 50% Good	5-15% Poor or > 50% Fair + Poor	> 15% Poor
Toxicity	< 5% Poor		> 5% Poor
TOC	< 20% Poor	20-30% Poor	> 30% Poor



Condition of Gulf of Mexico Estuaries

	NCA	NEP
DIN		
DIP		
CHL		
Clarity		
DO		
Water Quality		

	NCA	NEP
Toxicity		
Contam		
TOC		
Sediment Quality		
Benthic Index		





