
US ERA ARCHIVE DOCUMENT



Cyanobacterial Harmful Blooms (CyanoHABs): Symptomatic of human and climatic alteration of aquatic environments

Urban, agricultural and industrial expansion

Increasing nutrient (Nitrogen & Phosphorus) inputs

Water use and hydrologic modification play key roles

Climate (change) plays a key interactive role Blooms are intensifying and spreading



Why the concern about HABs?

- Toxic to zooplankton, fish, shellfish, domestic animals and humans
 - Cause hypoxia and anoxia, leading to fish kills
 - Odor and taste problems
 - loss of drinking water recreational, fishing use/sustainability















It's a global problem

• Freshwater Ecosystems (lakes, reservoirs, streams, rivers)


















What Controls CyanoHABS? Interacting Physical, Chemical & Biotic Factors



Nutrients: N and P inputs have long been recognized as playing a key role in bloom dynamics, but there's a controversy......

"Eutrophication of lakes cannot be controlled by reducing nitrogen input: Results of a 37-year whole-ecosystem experiment"

Schindler et al. Proceedings of the National Academy of Science USA 105:11254-11258 (2008).


Conclusion by Schindler et al. (2008) (based on one lake: Lake 227)

assumes that CyanoHAB N₂ fixation will supply ecosystem N needs


Therefore, why worry about N? Argument extended to

estuarine and coastal systems

These assumption have been challenged (Lewis and Wurtsbaugh 2008; Conley et al., 2009; Paerl 2009; Scott & McCarthy 2010: Lewis et al. 2011)



Lake Taihu 3rd largest lake in China. Nutrients (Lots!) associated with unprecendented human development in the Taihu Basin (Jiangsu Province). Results: Blooms have increased to "pea soup" conditions within only a few decades




The water crises (2007-?) in the Taihu Basin: **Cessation drinking water use for >20 million (hepato- and neuro-toxins)**


Curtailed recreational use (contact dermatitis)

Fisheries (commercial and recreational)



The nutrient "problem" in Taihu (and other hypertrophic lakes)






Nutrient (N&P) ratios in Taihu

Redfield (balanced growth) ~15:1 (N:P)


HYPOTHESIS Dual (N & P) reductions will be needed to stem eutrophication and CyanoHABs

Xu et al., 2010


Effects of nutrient (N & P) additions on phytoplankton production (Chl a) in Lake Taihu, China: Both N & P inputs matter!!




Xu et al. 2010; Paerl et al. 2011



CyanoHAB Toxicity
Related to nutrient inputs and biomass
Chlorophyll a is a sensitive, relevant and
easy to use indicator



Otten et al., 2011, 2012; Wilhelm et al., 2011



Is Taihu a "looking glass" for hypertrophic shallow ecosystems worldwide?






















Florida lakes: Cylindrospermopsis raciborskii, rapidlyproliferating, toxic N₂ fixing cyanoHAB

- High P uptake and storage capacity
- High NH₄⁺ uptake affinity (competes well for N)
 - N additions (NO₃⁻ + NH₄⁺) often significantly increase growth (chl a and cell counts) and productivity
- N₂ fixer (can supply its own N needs)
- Tolerates low light intensities
 - Eutrophication/decreased transparency favors Cylindro
 - Often in water column with other cyanoHABs






St. Johns R. System, FLorida: Nitrogen <u>and</u>
Phosphorus Effects on CyanoHAB Growth and Bloom
Potential (*Cylindrospermopsis raciborskii*)




Take home message: Cylindrospermopsis raciborskii is opportunistic Dual N & P input constraints will likely be needed to control it

Piehler et al, 2009

Whole-Lake Fertilization Experiments (Canada: ELA, Quebec, NWT, Sweden)




A summary of N & P limitation in lakes worldwide




Lakes: N= 55

Lewis et al., ES&T 45:10300-10305 (2011)

Nutrient limitation along the freshwater to marine continuum Estuaries with CyanoHABs: the Chesapeake Bay






Nutrient load and phytoplankton (dominated by cyanobacteria) growth response in Himmerfjärden (freshwater to brackish), Sweden

Courtesy: Ulf Larsson & Ragnar Elmgren
Stockholm University





The Himmerfjärden case: Baltic coastal area with large Sewage treatment plant,


P removal since 1976

N removal started in 1993 (50%) & 2000 (80%).


No N removal 2004-2008

EFFECTS ON PHYTOPLANKTON (Chl a)?


Plant loads , tonnes/ year




H4 = Eutrophicated station **B1** = Reference station




The results: Reducing DIN inputs reduced Chl a




Larsson and Elmgren, 2012





Developing a N loading-bloom threshold






Lowering nitrogen discharge below 400 tonnes/yr clearly reduced local phytoplankton biomass.


Source: Ulf Larsson, pers.comm.

Lets go back to the 'P only" paradigm from whole-lake experiments, suggesting that P alone controls algal biomass (Schindler et al., 2008)


The Argument:

If nitrogen is in short supply, nitrogen fixation by cyanobacteria will make up the nitrogen deficit:





Lets look at the data for lake 227: What happened following N&P fertilization (1968-1989), then only P fertilization (1989 →)?





Following cessation of N fertilization, Total N, TN:TP, and phytoplankton biomass decreased. N_2 fixation could not keep up with ecosystem N demands (Scott & McCarthy L&O 55:1265-1270 (2010)

WHY?????


Its not because there's a shortage of N2 fixing taxa












Controls on N₂ fixation: Its not just P or N:P...Other controls



Also, N₂ losses from shallow eutrophic systems exceed "new" N inputs via N₂ fixation

Annual estimates of ecosystem N₂ fixation, denitrification, and net ecosystem N₂ flux in lakes.


Location	N ₂ Fixation (g N m ⁻² yr ⁻¹)	Denitrification (g N m ⁻² yr ⁻¹)	Net N ₂ Flux (g N m ⁻² yr ⁻¹) ¹
Lake 227 (ELA) ²	0.5	5-7	-6.5 – -4.5
Lake Mendota ²	1.0	1.2	-0.2
Lake Okeechobee ²	0.8 - 3.5	0.3 - 3.0	-2.2 - 0.5
Lake Erken ²	0.5	1.2	-0.7
Lake Elmdale	10.4^{3}	18^{4}	-7.6
Lake Fayetteville	10.6^{3}	234	-12.4
Lake Wedington	7.0^{3}	124	-5.0

¹Net negative N₂ flux represents reactive N loss, positive represents gain; ²Paerl and Scott (2010); ³J.T. Scott (unpublished data); ⁴Grantz et al. (2012)

Conclusions: 1. N₂ fixation does NOT meet ecosystem N demands

- 2. More N inputs will accelerate eutrophication
- 3. We Gotta get serious about controlling N!!

Confounding Impacts of Climate Change: Its Getting Warmer






Additional Evidence

2003 was the hottest summer in 500 years in Europe!
2005, 2009, 2012 were the hottest years ever in N. America
2010 hottest year in central Asia


Below is the mean summer surface water temperature in a range of Dutch lakes



Huisman et al. 2006; KNMI, De Bilt, the Netherlands


Temperature affects stratification

density profile of water





At higher temperature, stratification is stronger!

density change per 1



Thermal stratification






Buoyant Cyanos favored by Stronger Stratification







Paerl and Huisman 2009


Mid August 2003:

Nieuwe Meer, Holland

Heatwave & little mixing

Microcystis benefits!







Jöhnk et al., 2008

Testing the Model





Lake data




Huisman et al., 2004

The link to CyanoHABs...... Temperature affects growth rates




References: Kraweik 1982, Grzebyk & Berland 1996; Kudo et al., 2000, Litaker et al., 2002, Briand et al., 2004, Butterwick et al., 2005,

Yamamoto & Nakahara 2005, Reynolds 2006



Cyanobacterial dominance along temperature & nutrient gradients in 143 lakes




Percentage of cyanobacterial biovolume in phytoplankton communities as a function of water temperature and nutrients in 143 lakes along a climatic gradient in Europe and South America.

- (a) Combined effects of temperature and nutrients as captured by a logistic regression model
- (b) Response surface obtained from interpolation of the raw data using inverse distance weighting.

From Kosten et al. (2012). Global Change Biology

Temperature increases and longer-lasting, more intense cyanobacterial blooms in Taihu. Is warming changing CyanoHAB thresholds?



Hydrologically: Things are getting more extreme

· Storms, droughts more intense, extensive & frequent

Melillo et al., 2014. 3rd National Climate Assessment, US Global Change Research Program





















Hydrology (flushing) interacts with temperature to determine diatom (fucoxanthin) and cyanobacterial (zeaxanthin) dominance in Neuse R. Estuary






Diatoms like it cool & fast



Cyanos like it hot & slow



Paerl et al. 2009

What's being done to "fix" Taihu?

- Need for dual nutrient reduction strategy recognized
- Wastewater collection and treatment
- Creating buffer zones
- Dredging sediments
- Diversion of Yangtze R. water (to enhance flushing)
- Algal Collection techniques
- Precipitation/flocculation

Since 2007, Wuxi has built 68 new wastewater plants, treating 1,513,000 ton/day, upgraded all wastewater treatment levels for BOD, starting P but not (yet) N removal.






Using vegetative buffers to retain storm water nutrients



(Qin et al., 2012)

Construction of buffer zones and wetlands around the lake





面积(km²)

66.61

44.39

142.55

44.68

60.81

44.50

41.38

7.39

452,31

编号

(1)

(5)

6

区段

无锡滨湖区段

苏州高新区段

苏州吴中区段

苏州吴江市段

湖州吴兴区段

湖州长兴县段

无锡宜兴市段

常州武进区段

合计






图例	土地利用	丽积(km²)
	城镇农村及工交建设用地	72.98
	林地、果园	42.26
	农田	192.61
	河流沟渠	29.68
	水库、坑塘	114.78

(叶春提供)



Yangtze River water diversion through Taihu: Cure or Curse?




- The heavily polluted lake regions (i.e. Meiliang Bay, Zhushan Bay and west littoral) are not remedied by the water transfer route and volume.
- Water residence time is reduced from \sim 320 days to \sim 220 days at best. This has no significant impact on reducing bloom potentials.
- ➤ The diversion may makes things worse...input from the heavily polluted Yangtze River adds ~15% to the N and P loads to the lake, potentially exacerbating eutrophication and blooms.



Polluted sediment dredging

40 km2 in north-western area and 1200 cubic meters (Lake area 2,300 km2). This has actually made the situation worse by releasing a legacy of nutrients and other pollutants, AND dredged sediments deposited in catchments are surrounding the lake Basin



(省水利厅提供)

CyanoHAB collection and removal

*Between 2007 and 2012, 3500,000 tons of algal scum had been removed. However, this is <1% of what was produced!!





Flocculation and precipitation of CyanoHABs

*Trials indicate that sediment resuspension overwhelms


"sediment capping" (i.e., can't hide the problem). *Costs prohibitive. *Doesn't solve the excess nutrient problem

Bottom Line: Need to reduce BOTH N and P inputs, no matter what other "fixes" are proposed

Other physical-chemical approaches

Altering circulation

Solar-Powered, Long-Distance Circulation




- May discourage formation of some buoyant cyanoHABs in confined waters
- May also increase primary production (by obviating light limitation)
- Nutrient transport and circulation??

Anoxic Water

Hudnell 2013

Effects of ultrasound on cyanobacterial biomass



Note: cell content leaks out upon lysis





Figure 2. Changes in algal composition in control and ultrasound-treated ponds.

Zimba et al., 2008; Wu et al., 2012

Hydrogen Peroxide Treatment



II TOXIC II GYANOBACTERIA









Matthijs et al., 2012; Barrington et al., 2013

Management ramifications

- In most cases, both N & P reductions are needed (no matter what other physical/chemical approaches are used)
- Nutrient-bloom threshold are system-specific
- · Nutrient-bloom thresholds may be changing
 - May need to reduce N and P inputs even more in a warmer world
- Nutrient input restrictions year-round
 - Cyanos are favored by higher temperatures, longer warm seasons







Thanks!!

www.unc.edu/ims/paerllab/research/cyanohabs/

Thanks to:

A. Joyner

T. Otten

B. Peierls

B. Qin

M. Piehler

K. Rossignol

5. Wilhelm

H. Xu

G. Zhu


TLLER "crew"











Additional support: Nanjing Instit. of Geography and Limnology, Chinese Academy of Sciences NIGLAS