


US EPA ARCHIVE DOCUMENT

Investigating the effects of atmospheric aging on the radiative properties and climate impacts of black carbon aerosol



Jesse Kroll, Colette Heald

Department of Civil and Environmental Engineering, MIT




Paul Davidovits

Department of Chemistry, Boston College


22 May 2012

Organic aerosol (circa 2005)

Discrepancies in mass loadings between modeled, ambient OA:



*Heald et al.
2005*




*Volkamer et al.
2006*

*also: de Gouw et al. 2005,
Kleinman et al. 2008,
etc...*

Organic aerosol, cont.

Discrepancies in chemical composition (degree of oxidation) between lab, ambient OA:




Aiken et al.
2008

also: Bahreini et al. 2005, Heald et al. 2010, Kroll et al. 2011, etc...


Organic aerosol, cont.

Discrepancies in key properties (volatility) between lab, ambient OA:




Huffman et al.
2009

“Aging” of organic aerosol



Robinson et al. 2007
Jimenez et al. 2009


“Aging” of organic aerosol



Robinson et al. 2007

Jimenez et al. 2009

“Aging” of organic aerosol




Robinson et al. 2007

Jimenez et al. 2009


Aging of OA can describe a large number of processes:

- Oxidation reactions (gas- or condensed-phase)
- Nonoxidative chemical transformations
- Evaporation/condensation

Laboratory experiments of OA aging




1) environmental chamber





2) flow tube reactor

Oxidative aging of OA: Lab studies




squalane
 $C_{30}H_{62}$
 + OH



Oxidative aging: Effects on OA properties

Water uptake




Jimenez et al. 2009

*also: Chang et al. 2010,
Massoli et al. 2010,
Cappa et al. 2011, etc.*

Cappa et al. 2011

Optical properties



This project: Aging of black carbon aerosol

Hypothesis: Similar to the emerging view of organic aerosol, black carbon aerosol is a chemically dynamic system, subject to dramatic changes in physicochemical properties, and therefore climate forcing effects, as a result of atmospheric aging.

This project: Aging of black carbon aerosol


***Hypothesis:** Similar to the emerging view of organic aerosol, black carbon aerosol is a chemically dynamic system, subject to dramatic changes in physicochemical properties, and therefore climate forcing effects, as a result of atmospheric aging.*

Major questions:


- what are the most important atmospheric aging transformations of BC?
- what sort of effects does aging have on climate-relevant properties of BC?
- how do these aging reactions impact BC direct radiative forcing?

Key BC aging reactions

1) Coatings




2) Heterogeneous oxidation




Key BC aging reactions

1) Coatings




2) Heterogeneous oxidation




Effects of aging on BC properties

1) Enhancement of light absorption by coatings

[e.g., Schnaitner 2005, Bond et al. 2006, Schwarz et al. 2008, Lack et al. 2009]



2) Increased water-uptake ability by coated or oxidized BC





This additional hygroscopicity can potentially lead to ...

- more efficient light scattering (due to larger particles from water uptake)
- more facile activation to form cloud droplets
- shorter atmospheric lifetimes due to increased wet deposition

Generic BC aging in global models*

AGING (oxidation, coating):
hydrophobic to hydrophilic,
 $\tau=1.2$ days



*Park et al. [2005]: ~1 day conversion
gave best agreement with measured
BC export efficiency*

*including GEOS-Chem mass-based scheme to be used here

Aging in global models

AeroCom models	Energy Emiss ⁽¹⁾	BB Emiss ⁽¹⁾	Aging ⁽²⁾	BC lifetime days	Ice/snow removal ⁽³⁾	Mass median diameter of emitted particle ⁽⁴⁾	BC density g cm ⁻³ ⁽⁶⁾	Refractive index at 550 nm ⁽⁶⁾	MABS m ² g ⁻¹ ^(5,6)	References for aerosol module
GISS 99	B04	GFED	A	7.2	12%	0.08	1.6	1.56–0.5i	8.4	Koch et al. (2006, 2007), Miller et al. (2006)
ARQM 99	C99	L00, L96	I	6.7	T	0.1	1.5		4.1	Zhang et al. (2001); Gong et al. (2003)
CAM	C99	L96	A		L		X	X	X	Collins et al. (2006)
DLR	CW96	CW96	I		5% accum, strat	0.08, 0.75 FF 0.02, 0.37 BB	X	X	X	Ackermann et al. (1998)
GOCART	C99	GFED, D03	A	6.6	T	0.078	1.0	1.75–0.45i	10.0	Chin et al. (2000, 2002), Ginoux et al. (2001)
SPRINTARS	NK06	NK06	BCOC		L	0.0695 FF, 0.1 others	1.25	1.75–0.44i	2.3	Takemura et al. (2000, 2002, 2005)
LOA B	B04	GFED	A	7.3	LI	0.0118	1.0	1.75–0.45i	8.0 #	Boucher and Anderson (1995); Boucher et al. (2002); Reddy and Boucher (2004); Guibert et al. (2005)
LSCE	G03	G03	A	7.5	L	0.14	1.6	1.75–0.44i	3.5 (4.4 #)	Claquin et al. (1998, 1999); Guelle et al. (1998a, b, 2000); Smith and Harrison (1998); Balkanski et al. (2003); Bauer et al. (2004); Schulz et al. (2006)
MATCH	L96	L96	A		L	0.1	X	X	X	Barth et al. (2000); Rasch et al. (2000, 2001)
MOZGN	C99, O96	M92	A		L	0.1	1.0	1.75–0.44i	8.7	Tie et al. (2001, 2005)
MPIHAM	D06	D06	I #	4.9	S	0.069 (FF, BF) 0.172 (BB)	2.0	1.75–0.44i	7.7 #	Stier et al. (2005)
MIRAGE	C99	CW96, L00	I #	6.1	L	0.19, 0.025	1.7	1.9–0.6i	3 airtk, 6 acc	Ghan et al. (2001); Easter et al. (2004); Ghan and Easter (2006)
TM5	D06	D06	A	5.7	20%	0.034	1.6	1.75–0.44i	4.3	Metzger et al. (2002a, b)
UIOCTM	C99	CW96	A	5.5	L	0.1 (FF), 0.295, 0.852 (BB)	1.0	1.55–0.44i	7.2 #	Grini et al. (2005); Myhre et al. (2003); Berglen et al. (2004); Bernsten et al. (2006)
UIOGCM 99	IPCC	IPCC	I #	5.5	none	0.0236–0.4	2.0	2.0–1.0i	10.5 #	Iversen and Seland (2002); Kirkevåg and Iversen (2002); Kirkevåg et al. (2005)
UMI	L96	P93	N	5.8	L	0.1452 (FF), 0.137 (BB)	1.5	1.80–0.5i	6.8 #	Liu and Perner (2002)
ULAQ 99	IPCC	IPCC	A	11.4	L	0.02–0.32	1.0	2.07–0.6i	7.5 #	Pitari et al. (1993, 2002)

Koch et al. 2009

N = no aging

A = aging at a fixed lifetime

I = aging with coagulation and condensation

= aging affects optical properties

This project: Aging of black carbon aerosol

***Hypothesis:** Similar to the emerging view of organic aerosol, black carbon aerosol is a chemically dynamic system, subject to dramatic changes in physicochemical properties, and therefore climate forcing effects, as a result of atmospheric aging.*


Major questions:

- what are the most important atmospheric aging transformations of BC?
- what sort of effects does aging have on climate-relevant properties of BC?
- how do these aging reactions impact BC direct radiative forcing?


Approach:

- detailed **laboratory studies** of individual aging reactions
- **parameterization of optical properties** as a function of size, RH, mixing state
- **incorporation into a global model** framework for calculation of direct radiative forcing


Laboratory setup



soot particle
generation




reagent/oxidant
preparation



Key measurements



Particle mass, density



Particle chemical composition

Particle hygroscopicity (κ)
CCN activity (κ)

Multi-wavelength absorption, extinction, scattering (MAC, SSA, etc.)



Experimental matrix

Based upon “BC²” intercomparison study, 318 runs [Cross *et al.* 2010]

BC source

- fractal soot from McKenna burner (denuded at 300°C)
- also atomized black carbon spheres

Particle size


- monodisperse, 30-300 nm

Relative humidity

- controlled after reactor, but before instruments (multiplex of 0%, 30%, 60%, 90%)


Aging type

- Heterogeneous oxidation (OH, O₃)
- coated with sulfuric acid (SO₂)
- coated with SOA (fresh, aged)
- SOA + H₂SO₄




Parameterization

- Calculation of radiative forcing in models requires knowledge of key optical parameters as a function of particle properties
- This will be done by construction of a “lookup table” (or interpolated function) based on experimental results



BC in GEOS-Chem (current treatment)


AGING (oxidation, coating):
hydrophobic to hydrophilic,
 $\tau=1.2$ days



WET
DEPOSITION


Currently aerosol is treated as (optically) externally mixed, with hydrophilic and hydrophobic BC having the same optical properties (with the exception of water uptake).

BC AOD



0.00 0.01 0.03 0.04 0.05

% of AOD from BC




0 2 4 6 8 10 [%]

$$\tau = \frac{3}{4} \frac{Q_{\text{ext}} M}{r_{\text{eff}} \rho} = \alpha M$$


Modeling BC aging in GEOS-Chem

MASS-ONLY SCHEME

AGING (oxidation, coating):
hydrophobic to hydrophilic,
 $\tau=1.2$ days



SECTIONAL SCHEME (TOMAS)




30 internally mixed bins from 0.01-10 μm (dry)

[Trivitayanurak et al., 2008]

Here: allow for physical aging through either (1) 1-day timescale (2) assume internal mixing (for all or within bins) or (3) implement a kappa-based transition to soluble aerosol

Calculation of direct radiative forcing by BC

- (1) AER's *Rapid Radiative Transfer Model (RRTMG)* will be integrated into GEOS-Chem (online) this Fall for application here. (RRTMG utilizes the correlated-k approach to calculate fluxes and heating rates efficiently and accurately. Has been applied in many GCMs, well-documented comparison with line-by-line code.)
- (2) Optical properties (mass extinction efficiency, SSA, asymmetry parameters and phase function) will be applied from lookup table in shortwave to GEOS-Chem simulated BC to estimate DRF.




Investigating the sensitivity of DRF to BC aging

Laboratory measured optics



Simulated direct radiative forcing



The modeling, based upon detailed laboratory studies of BC aging, will allow for the estimation of how chemical transformations of BC particles impact direct radiative forcing (and export efficiency, deposition to the Arctic, other ancillary effects of aging timescales).