

US EPA ARCHIVE DOCUMENT



# Tracing Calcium in Forest Ecosystems: Ca/Sr and Ca isotopes

## BACKGROUND

Acid deposition significantly affects forest ecosystems (e.g. the Northeast U.S.)

- o Effects:
  - o Soil acidification
  - o Leaching of base cations from the soil, especially calcium
  - o Mobilization of aluminum (Al is toxic to both plants and animals)
  - o Increased nitrate availability
  - o Declines in symbioses between plants and mycorrhizal fungi


Resulting Ca limitation is correlated with declines in forest health

Ca pools (boxes) and fluxes (arrows) are poorly identified and unquantified.

o Ca/Sr values are commonly used to identify Ca sources and estimate fluxes (e.g. Blum et al. 2002 and Bailey et al. 1996)

o Current methods assume that there is little fractionation between Ca and Sr along the source - foliage path.

o There is evidence that trees are directly accessing mineral Ca through symbiotic relationships with mycorrhizal fungi - bypassing the soil Ca pool (Landeweert et al. 2001)



## RESEARCH DIRECTION

- 1.) Are Ca and Sr fractionated in terrestrial biogeochemical cycles?
  - a.) Is there significant fractionation between Ca and Sr along the source-foliage path?
  - b.) If so, what is the magnitude?
- 2.) Do environmental conditions, N-deposition or fungal associations, enhance this fractionation?
- 3.) Do Ca isotope ratios provide a more reliable way of tracing the movement of Ca in forest ecosystems? Questions 1 and 2 must be addressed for Ca isotopes.

➤ I am investigating these questions in cultured Scotch pines, controlling the following variables: N species, N supply rate, and presence of mycorrhizal fungi.



## PRELIMINARY RESULTS






Figure 1: Results from a culture study showing biological fractionation within individual pine seedlings at the one-sigma level (ICP-AES).

Treatments:  
 H = high nutrient supply rate  
 L = low nutrient supply rate  
 Non = non-mycorrhizal  
 S = *Suillus luteus*  
 T = *Thelephora terrestris*

Figures 2 and 3: Effects of nitrogen (high vs. low and nitrate vs. ammonium) and mycorrhizal (non = non-mycorrhizal, S = *Suillus bovinus*, L = *Laccaria laccata*) on enrichment factors, F.

$$F_{\text{ratio}} = (\text{Needle ratio}) / (\text{Root ratio})$$

- *Suillus* may block plant uptake of aluminum (Fig. 3 below)



ACKNOWLEDGEMENTS: J.G. Bryce, E.A. Hobbie, R. Hallett, J. Colpaert, & C.-T. Lee

## CONCLUSIONS AND IMPACTS

- Ca/Sr is not a conservative tracer; ratios of foliage do not represent ratios at soil-root interface.
- Other tracers, e.g. Ca isotopes, are required to constrain Ca cycling in terrestrial systems.
- Development of new tools (tracers) will further our understanding of the Ca cycle in forests.
- Effective forest management practices (e.g. liming), environmental policies, pollution controls, commercial industries (e.g. maple syrup), etc. require a more comprehensive understanding of terrestrial calcium.

REFERENCES: Bailey, S.W., et al. 1996. Water Resour. Res. 32: 707-719. Blum, J.D., et al. 2002. Nature. 417: 729 - 731. AND Landeweert, R., E. et al. 2001. TREE. 16: 248-254.

US EPA ARCHIVE DOCUMENT