



US EPA ARCHIVE DOCUMENT





## Death on the Danube


**T**HE leakage of 100,000 tonnes of cyanide-laden sludge into the Danube river has already taken a heavy ecological toll. The cyanide, which was used at a mine in Baia Mare, in Romania, to separate gold from ore, was washed from an open storage basin into the Tisza, a tributary of the Danube, on January 30th. Three weeks and almost 1,000km (625 miles) later, almost everything in the Tisza's murky, brown water is lifeless. Well over 100 tonnes of dead fish have been scooped out downstream in Hungary and Yugoslavia; plankton, invertebrates and birds have been killed too. Fishermen have hung black banners from houses and bridges, mourning their livelihoods. Wells near the river have been closed.

With concentrations of 2.7 milligrams

# Why Nanotechnology?



**Sensors:**  
Small  
Cheap  
Low Power  
Highly distributed




*Nature Materials* **2002**, *1*, 39-41.

## Nanosensors (“Smart Dust”)






*What functionalities can be incorporated into a grain of sand?*




- Filtration
- Sample concentration
- Chemical sensing
- Biological sensing
- Environmental stability
- Internal referencing/drift correction
- Remote identification
- Remote interrogation (>100 m)
- Remotely triggered chemical processing
- Targeted motion
- Collective behavior (swarming)

# Photonic Crystals





# Chemical Sensing with a Si Photonic Crystal

QuickTime™ and a MPEG-4 Video decompressor are needed to see this picture.




$$I = 4I_0 \cos^2\left(2\pi \cdot \frac{nd}{\lambda}\right)$$



# Amplified Vapor Sensing in Microporous Media

**Capillary condensation:** Liquids spontaneously condense from vapor into cracks and pores as bulk liquid.



*Schematic of a PS sample*

**Kelvin Equation:**


$$k_B T \ln\left(\frac{P}{P_s}\right) = \gamma V_m \left(\frac{1}{R_K}\right)$$

Surface energy

Molar volume of the liquid

Effective pore radius

# Robotic platform integrating “smart dust” photonic crystal VOC sensors




QuickTime™ and a H.263 decompressor are needed to see this picture.

**Photo  
transistors**

**Porous Si  
“Smart Dust”  
Particle**

**LED**



The robot follows a chemical track by sensing light transmitted through a chemically responsive photonic crystal. The robot uses two phototransistors, each covered with a porous silicon film as the chemical detector. Each sensor closes a circuit to one of the two drive motors when it detects a chemical vapor. The circuit memorizes which side the chemical it is on and activates the motor on the opposite side, steering the robot towards the chemical.





# Smart Dust

## Kris Pister, UCB

“The Smart Dust project is probing microfabrication technology's limitations to determine whether an autonomous sensing, computing, and communication system can be packed into a cubic millimeter mote (a small particle or speck) to form the basis of integrated, massively distributed sensor networks.”

-*Computer*, **34**, 44-51 (2001).



## Death on the Danube

**T**HE leakage of 100,000 tonnes of cyanide-laden sludge into the Danube river has already taken a heavy ecological toll. The cyanide, which was used at a mine in Baia Mare, in Romania, to separate gold from ore, was washed from an open storage basin into the Tisza, a tributary of the Danube, on January 30th. Three weeks and almost 1,000km (625 miles) later, almost everything in the Tisza's murky, brown water is lifeless. Well over 100 tonnes of dead fish have been scooped out downstream in Hungary and Yugoslavia; plankton, invertebrates and birds have been killed too. Fishermen have hung black banners from houses and bridges, mourning their livelihoods. Wells near the river have been closed. With concentrations of 2.7 milligrams

# Field trials with Smart Dust at SMER



*Smart Dust*



Santa Margarita Ecological Reserve (SMER), San Diego county CA

In Aug 2002, we performed the first field trials of porous Si “smart dust” and small wireless chip-based systems for detection of environmental pollutants. The field trials were performed in the Santa Margarita Ecological Reserve, over 24 hours, and involved two undergraduates, two high school students, a graduate student, and a post-doc. A simulant (ethanol) was successfully detected from a distance of 25 meters.



From left: Jamie Link (UCSD Grad), Pat Wang (Torrey Pines HS), Jason Dorvee (UCSD Undergrad), Frederique Cunin (UCSD Post-doc), Prof. Mike Sailor, Stephanie Tsai (Torrey Pines HS) at the entrance to SMER.




Jason Dorvee (UCSD Undergrad) doses smart dust sample with simulant vapor, assisted by Dr. Frederique Cunin (UCSD Post-doc)





# Smart Dust: sensors in a grain of sand



1 mm


# Synthesis of bilayered 1-D photonic crystals





Bilayered rugate at  
 $\text{CH}_2\text{Cl}_2$ /water  
interface

# Specificity: Fluorescent molecular wires for detection of Cr(VI) in water (with W. C. Trogler)



# Chromate Sensing with Silole-amine Nanoparticulate

Monomer

Nanoaggregate

0.5ppm Cr<sup>VI</sup>


1ppm Cr<sup>VI</sup>



William Trogler, Sarah Toal




# Nanoextraction



Hydrosilylated Porous Si film immersed in water traps hydrophobic liquids.

- Hydrophobic liquids are trapped in hydrophobic porous Si matrix.
- Pollutants can be extracted and concentrated in this film for subsequent detection.

# Application areas



Crosslinked Porous Si/polynorbornene composite film held between two pairs of tweezers, showing the mechanical stability of the materials.

- Chem/Bio detection in distributed sensor networks: buildings (HVAC), environment (air and water pollution monitoring), and public spaces (homeland security). Provide higher selectivity and sensitivity at lower cost, reduced power requirements.
- Medical implants, monitoring.
- High throughput screening for drug discovery, genomic sequencing.