


US EPA ARCHIVE DOCUMENT

# Fundamental Studies of Secondary Organic Aerosol (SOA) Formation




**John H. Seinfeld**

California Institute of Technology

EPA Atmospheric Science Progress Review Meeting

June 22, 2007

# Formation of secondary organic aerosol (SOA)



# Secondary organic aerosol (SOA)

---

- Globally: ~10-40% of organic aerosol is secondary [*Kanakidou et al.*, 2005]
- Urban areas: ~50-80% [*Kanakidou et al.*, 2005]
- SOA precursors
  - Biogenic emissions (isoprene, monoterpenes, sesquiterpenes)
  - Anthropogenic emissions (aromatic hydrocarbons)

Hydrocarbon	Emission [Tg / y]	SOA Production [Tg / y]
aromatics	19	3.7
terpenes	153	12
alcohols	41	1.9
sesquiterpenes	15	2.3
isoprene	461	?
Total	689	19.9

*Henze et al.*, 2007

# Caltech environmental chambers



- 2 Teflon chambers, 28 m<sup>3</sup> each
- Differential Mobility Analyzer (DMA): particle size distribution, volume
- Time-of-flight Aerodyne Aerosol Mass Spectrometer (AMS): particle mass, composition
- GC-FID: hydrocarbon
- Proton Transfer Reaction Mass Spectrometer (PTR-MS): hydrocarbon, reaction products
- Filter samples: off-line chemical analysis
- O<sub>3</sub>, NO<sub>x</sub>, RH, T

# Oxidation of biogenic hydrocarbons

## Ozonolysis:

- T=20°C, RH<10%
- OH Scavenger: cyclohexane
- (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> seed
- Reaction initiated upon addition of O<sub>3</sub>

## Photooxidation:

- T=20-22°C, RH~50%
- HONO as OH precursor: dropwise addition of 1% NaNO<sub>2</sub> into 10% H<sub>2</sub>SO<sub>4</sub>
- (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> seed
- Reaction initiated by irradiation with UV lights

Isoprene (C<sub>5</sub>H<sub>8</sub>)



Monoterpenes (C<sub>10</sub>H<sub>16</sub>)




α-pinene




β-pinene




Δ<sup>3</sup>-carene




myrcene




terpinolene



limonene




α-terpinene




γ-terpinene


Sesquiterpenes (C<sub>15</sub>H<sub>24</sub>)




α-humulene



β-caryophyllene




aromadendrene




longifolene


Oxygenated Terpenes



methyl chavicol






linalool



verbenone

# $\alpha$ -pinene ozonolysis










- SOA is formed from the condensation of first-generation products and the first oxidation step is the rate-limiting step
- Organic acids have been identified as major particle-phase products: monocarboxylic acids (pinonic acid and norpinonic acid), dicarboxylic acids (pinic acid and norpinic acid), and hydroxy pinonic acid [e.g. *Yu et al.*, 1999]
- *Jenkin et al.* [2000] proposed pinic and hydroxy pinonic acid are first-generation products, which is consistent with our study

# Compounds with one double bond





Isoprene ( $C_5H_8$ )






Monoterpenes ( $C_{10}H_{16}$ )



Sesquiterpenes ( $C_{15}H_{24}$ )






Oxygenated Terpenes








# Terpinolene ozonolysis



- Terpinolene has multiple double bonds, so its first-generation products formed are still unsaturated, and they will further react with the ozone in the chamber to produce additional condensable products
- Further oxidation of first-generation products contributes significantly to SOA and this oxidation step may also be rate-limiting





# Time dependent growth vs. Final SOA growth $\alpha$ -pinene ozonolysis




- Time-dependent data overlap remarkably well with the final SOA growth curve
  - Odum equation (as well as the growth curve equation) is valid for the final growth as well as the time-dependent data

# Time dependent growth vs. Final SOA growth terpinolene ozonolysis




- Time-dependent growth curves and final growth curve do not overlap; time-dependent growth curves show clearly the contribution of the secondary reactions
- Cannot fit the time-dependent growth curves for terpinolene ozonolysis with Odum equation, confirming that this model is only valid when the data represent final SOA growth

# Conclusions



- Growth stops when all hydrocarbon is consumed
  - First step is rate-limiting
  - SOA formed from nonvolatile first-generation products




- Growth continues after all hydrocarbon is consumed
- Aerosol formed from further oxidation of first-generation products and this second oxidation step may also be rate-limiting



# Effect of $\text{NO}_x$ level on SOA Formation from Photooxidation of Biogenic Hydrocarbons



# Isoprene





- Global emissions of ~500 Tg/year [*Henze et al.*, 2007]
- First-generation oxidation products: all are volatile, not expected to partition into the aerosol phase
- *Pandis et al.* [1991], *Edney et al.* [2005] observed no SOA formation from irradiation of isoprene/NO<sub>x</sub> mixtures
- Possible contributions of isoprene to organic aerosol by heterogeneous chemistry [*Limbeck et al.*, 2003], [*Claeys et al.*, 2004], [*Edney et al.*, 2005]

# Experimental conditions

---

- Ammonium sulfate seed;  $T \sim 25^\circ\text{C}$ ,  $\text{RH} < 10\%$
- Low- $\text{NO}_x$  experiments
  - Radical source:  $\text{H}_2\text{O}_2 + h\nu \rightarrow \text{OH} + \text{OH}$
  - Peroxy radicals react with  $\text{HO}_2$
- High- $\text{NO}_x$  experiments
  - Radical source:  $\text{HONO} + h\nu \rightarrow \text{OH} + \text{NO}$
  - $\text{NO}_x$  is produced as side product
  - Peroxy radicals react with  $\text{NO}$


# Isoprene and SOA formation






# Isoprene: NO<sub>x</sub> dependence

40-45 ppb isoprene



- Decrease in SOA yield at high NO<sub>x</sub> [*Pandis et al.* 1991; *Zhang et al.*, 1992; *Hurley et al.*, 2001; *Johnson et al.*, 2004; *Song et al.*, 2005; *Presto et al.*, 2005]


# Growth curve: $\alpha$ -pinene photooxidation



- Same  $NO_x$  dependence as isoprene: higher  $NO_x$ , lower SOA growth
- SOA formed from the condensation of first-generation products and the first oxidation step is rate-limiting
- $H_2O_2+NO$ : Multiple SOA formation steps;  $\alpha$ -pinene only has one double bond, further SOA growth by
  - Particle-phase reaction
  - Further gas-phase reaction of reactive oxidation products (aldehydes, furans etc)


# Peroxy radical chemistry

---




- Small alkoxy radical easily fragmented
- Organic nitrates relatively volatile [*Presto et al.*, 2005]
- Peroxides: important SOA components [*Bonn et al.*, 2004; *Docherty et al.*, 2005]

# Growth curve: longifolene photooxidation



- **Reversed  $\text{NO}_x$  dependence: higher  $\text{NO}_x$ , higher SOA growth**
- High- $\text{NO}_x$ : maximum yield = 100-120%
- Low- $\text{NO}_x$ : constant yield = 75%


# Longifolene: $\text{NO}_x$ dependence



- Low  $\text{NO}_x$ : SOA Yield = 75%
- With  $\sim 300$  ppb  $\text{NO}$ : SOA Yield = 127%;


# Longifolene: NO<sub>x</sub> dependence

~5 ppb longifolene: increase in yield at high NO<sub>x</sub>





# Aromadendrene: NO<sub>x</sub> dependence

~5 ppb aromadendrene: increase in yield at high NO<sub>x</sub>




# Isomerization of alkoxy radicals





# Isomerization of alkoxy radicals


- SOA growth from large alkanes at ppm levels of NO [Lim et al., 2006]



Lim et al., 2006

# Organic nitrates: UPLC/ESI-TOFMS

- Extracted ion chromatograms shows the presence of *acidic* nitrates in longifolene SOA



# Conclusions

---

- Isoprene is an important SOA precursor (SOA: 14 Tg / year, *Henze et al.*, 2007)
  - Condensable products are second-generation
- Change in NO<sub>x</sub> dependence going from isoprene to sesquiterpene
  - Isoprene (C<sub>5</sub>H<sub>8</sub>) and α-pinene (C<sub>10</sub>H<sub>16</sub>): SOA yield decreases at high NO<sub>x</sub>
  - Longifolene and aromadendrene (C<sub>15</sub>H<sub>24</sub>): SOA yield increases at high NO<sub>x</sub>
- Isomerization of alkoxy radicals and formation of nonvolatile organic nitrates could be an efficient channel of SOA formation (for large hydrocarbon precursors)

# SOA Formation from Photooxidation of Aromatic Hydrocarbons




# Background

---

- Field studies suggest higher SOA formation than models predict [*De Gouw et al.*, 2005; *Volkamer et al.*, 2006]
  - SOA formed from anthropogenic sources is higher than currently thought
- SOA formation from aromatic hydrocarbons
  - Mechanisms poorly understood
  - Poor carbon balance, typically < 50% [*Calvert et al.*, 2002]
  - SOA yields vary with different NO<sub>x</sub> levels [*Hurley et al.*, 2001; *Johnson et al.*, 2005; *Martin-Reviejo et al.*, 2005, *Song et al.*, 2005]

# Previous studies

- Irradiation of aromatics/ $\text{NO}_x$  mixture [*Hurley et al.*, 2001; *Johnson et al.*, 2005; *Martin-Reviejo et al.*, 2005, *Song et al.*, 2005]
  - Changing oxidation conditions over the course of the experiments
  - Aerosol growth does not begin until NO approaches zero




- Problems
  - Urban areas are high- $\text{NO}_x$ , so this would suggest no aerosol formed from anthropogenic hydrocarbons
  - Experiments with higher levels of NO, and  $\text{CH}_3\text{ONO}$ , aerosols observed before NO approaches zero [*Stroud et al.*, 2004]


# Goals of experiments

---


- Study systematically the effect of  $\text{NO}_x$  on SOA formation from selected aromatic hydrocarbons



Benzene



Toluene



*m*-xylene

- Obtain SOA yields at high- and low- $\text{NO}_x$  conditions (the limiting cases), parameterize the  $\text{NO}_x$  dependence for modeling purposes
- Investigate the effect of particle phase acidity on aerosol growth


# Experimental conditions

---


- Ammonium sulfate seed;  $T \sim 25^\circ\text{C}$ ,  $\text{RH} < 10\%$
- Low- $\text{NO}_x$  experiments
  - Radical source:  $\text{H}_2\text{O}_2 + h\nu \rightarrow \text{OH} + \text{OH}$
  - Peroxy radicals react with  $\text{HO}_2$
- High- $\text{NO}_x$  experiments
  - Radical source:  $\text{HONO} + h\nu \rightarrow \text{OH} + \text{NO}$
  - $\text{NO}_x$  is produced as side product
  - Peroxy radicals react with  $\text{NO}$



# Growth curves: *m*-xylene




## High-NO<sub>x</sub>




- High-NO<sub>x</sub>: Growth curves do not overlap, multiple rate-limiting steps in SOA formation (first step is the slowest)
- Further-generation oxidation products

## Low-NO<sub>x</sub>




- SOA yields much higher than high-NO<sub>x</sub> experiments
- Constant SOA yield implies essentially nonvolatile oxidation products (36% yield)


# Growth curves: toluene





## High-NO<sub>x</sub>



## Low-NO<sub>x</sub>




# Growth curves: benzene




- ~400 ppb benzene (slow reaction rate, <20% reacted)
- Same NO<sub>x</sub> dependence as *m*-xylene and toluene: high NO<sub>x</sub>, lower yields
- Low NO<sub>x</sub>: constant yield of 37%

# Peroxy radical chemistry




# Effect of oxidation rate

HONO experiment




Classical experiment



- Loss of semivolatiles → rate effect


# Loss of semivolatiles

---



- Loss of semivolatiles (by photolysis, reactions in the gas phase to form volatile products, or deposition to chamber walls)
  - Lowers the concentration of the gas-phase semivolatile, thereby reducing the amount that partitions into the aerosol phase

# Seed acidity: acid seed vs. non-acid seed



- No acid effect observed
- Same observations in *m*-xylene oxidation

# Conclusions: Aromatic SOA

---

- SOA yields are highly dependent on  $\text{NO}_x$  levels (peroxy radical chemistry)
  - High  $\text{NO}_x$ : Usual Odum yield curve behavior (Yield ~ 5 -10%)
  - Low  $\text{NO}_x$ : Constant yield (Yield ~30%)
- Condensable compounds are second-generation products (further gas-phase and/or particle-phase reactions)
- No effect of particle phase acidity observed



# What do all these studies tell us?

---

- Growth curve ( $\Delta M_o$  vs.  $\Delta HC$ ) as a powerful approach to infer the general mechanism of SOA growth
- Profound effect of  $NO_x$  level on SOA formation (isoprene, monoterpenes, aromatics, sesquiterpenes)
- Discrepancy between modeled vs. measured SOA:
  - Of compounds studied in the laboratory, biogenics are the largest contributor to ambient SOA, and isoprene is the most important single precursor
  - SOA formation from aromatics significantly higher than previously measured but not sufficiently large to rival that of biogenics on a continental scale
  - According to recent CMU study (Robinson et al., 2007), SVOCs from primary organic aerosol emissions may themselves constitute a major class of SOA precursors