

US EPA ARCHIVE DOCUMENT

DATA EVALUATION RECORD

9-26-80

1. Chemical: Cited in report as 1-bromo-3-chloro-5, 5-dimethylhydantoin (DiHalo) (Shaughnessy #006315).  
Reported by Great Lakes Chemical Co. (Dr. John Sands) as "Bromochloro-5,5-dimethylhydantoin", an equal-molar combination of:
  - 1) Dibromo-5, 5-dimethylhydantoin (Shaughnessy #006317)
  - 2) Dichloro-5, 5-dimethylhydantoin (Shaughnessy #028501)
  - 3) 1-bromo-3-chloro-5, 5-dimethylhydantoin (Shaughnessy #006315)
  - 4) 3-bromo-1-chloro-5, 5-dimethylhydantoin (Shaughnessy Number not avail.)
2. Formulation: Report: Test material not labelled as to its percent active ingredient.  
GLCC (DR. Sands): 92.2% active ingredient.
3. Citation: Horne, J.D., et al. 1980. 96-hour static bioassays using two Great Lakes Chemical Corporation compounds with three marine and three freshwater species. NUS Corporation--Northern Environmental Services Div. (Pittsburgh, PA) and Southern Environmental Services Div. (Clear Lake, Texas). (Within Accession #243015.)
4. Reviewed by: James. D. Felkel  
Wildlife Biologist  
Ecological Effects Branch/HED
5. Date Reviewd: 9/26/80
6. Test Type: Aquatic Invertebrate 48-hour LC<sub>50</sub>
  - A. Test Species: Daphnia Magna
7. Reported Results: The 48-hour LC<sub>50</sub> is reported to be 0.48 mg/l with 95% confidence limits of 0.53 mg/l and 0.42 mg/l.
8. Reviewer's Conclusions: The study is scientifically sound and fulfills the requirement for an aquatic invertebrate 48-hour LC<sub>50</sub> for a manufacturing -use. The reported LC<sub>50</sub> value indicates that the test material is highly toxic to daphnids. Depending on future use patterns, a flow-through test may be advisable.

## Materials/Methods Reported

### A. Test procedure

Dechlorinated Pittsburgh tap water (pretreatment included activated carbon filters and U.V.-light) was used to prepare the test concentrations. The Daphnia cultures were purchased from Wards Natural Science Estab., Inc. on 2/14/80 and acclimated at 20° (+1°)C. First instars were used in all tests. Cultures were fed yeast solution every other day.

Rangefinder tests were conducted to estimate the upper and lower bounds of the concentration range expected to include the median lethal concentration and five concentrations spanning this range were selected for the definitive test.

All definitive treatment levels and controls were established in duplicate with 10 Daphnia magna individuals pipetted into each of the 250 ml test beakers containing a known concentration of test material. Water quality measurements taken at the start and finish of the experiment included:

- 1). Temperature (at all test levels); and
2. D.O. and pH. (at high, middle, and low concentrations and controls).

Parameters measured at the start of the experiment only, for the controls and highest test concentration, were alkalinity, hardness, and conductivity.

### B. Statistical Analysis

The Litchfield- Wilcoxon method was used to fit a line to mortality data plotted on log-probit paper.

## Discussion/Results Reported

After rangefinder testing, definitive test concentrations selected were 0.3, 0.4, 0.5, 0.7, and 1.0 mg/l. Cumulative mortality was as follows:

<u>Treatment Level (mg/l)</u>	<u>Time (hrs.)</u>			
	12	24	36	48
1.0	20	20	20	20
0.7	10	13	13	13
0.5	7	11	11	11
0.4	2	3	3	3
0.3	0	0	0	0
Control	0	0	0	0

The table shows that all mortality occurred within the first 24 hours. The 48-hour LC<sub>50</sub> was reported to be 0.48 mg/l with confidence limits of 0.53 mg/l and 0.42 mg/l.

## Reviewer's Evaluation

### A. Test Procedures

Procedures generally followed the 1978 Proposed Guidelines (subpart E) and approved protocols (Committee on Methods for Toxicity Tests with Aquatic Organisms, 1975). However, missing information included the method of storing the test material and the photoperiod/lighting conditions.

### B. Statistical Analysis

Statistical analysis by this reviewer of the submitted mortality data indicates LC<sub>50</sub> values of 0.54 mg/l using the probit and moving average methods, and 0.48 mg/l using the binomial test. These values closely approximate the submitted value.

### C. Results/Discussion

The submitted results indicate that the compound is highly toxic to daphnids.

### D. Conclusions

1. Category: Core

2. Rationale: The test appeared to satisfy the intent of the 1978 proposed guidelines (subpart E) for a manufacturing-use. Depending on future use patterns, however, a flow-through test may be advisable.

3. Repairability: N/A

data 1.0,.7,.5,.4,.3  
 data 20,20,20,20,20  
 data 20,13,11,3,0  
 run

80/09/24. 14.37.29.  
 BASIC PROGRAM S79LC50

DiHalo/Daphnia magna Acc. #243015

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.(PERCENT)
1	20	20	100	9.53674E-5
.7	20	13	65.	13.1588
.5	20	11	55.	41.1901
.4	20	3	15.	.128841
.3	20	0	0	9.53674E-5

THE BINOMIAL TEST SHOWS THAT .4 AND 1 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS .487282

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD  
 SPAN 4 G LC50 5.44358E-2 .544681 95 PERCENT CONFIDENCE LIMITS .497148 .602398

-----RESULTS CALCULATED USING THE PROBIT METHOD  
 ITERATIONS 6 G H GOODNESS OF FIT PROBABILITY .102371 1 .13592

SLOPE = 7.38204  
 95 PERCENT CONFIDENCE LIMITS = 5.02013 AND 9.74396

LC50 = .54575  
 95 PERCENT CONFIDENCE LIMITS = .492563 AND .609742

SRU 1.314 UNTS.

RUN COMPLETE.

4

DATA EVALUATION RECORD

1. Chemical: Cited in report as 1-bromo-3-chloro-5, 5-dimethylhydantoin (DiHalo) (Shaughnessy #006315).  
Reported by Great Lakes Chemical Co. (Dr. John Sands) as "Bromochloro-5, 5-dimethylhydantoin", an equal-molar combination of:
  - 1) Dibromo-5, 5-dimethylhydantoin (Shaughnessy #006317)
  - 2) Dichloro-5, 5-dimethylhydantoin (Shaughnessy #028501)
  - 3) 1-bromo-3-chloro-5, 5-dimethylhydantoin (Shaughnessy #006315)
  - 4) 3-bromo-1-chloro-5, 5-dimethylhydantoin (Shaughnessy Number not avail.)
2. Formulation: Report: Test material not labelled as to its percent active ingredient.  
GLCC (Dr. Sands): 92.2% active ingredient.
3. Citation: Horne, J.D., et al. 1980. 96-hour static bioassays using two Great Lakes Chemical Corporation compounds with three marine and three freshwater species. NUS Corporation--Northern Environmental Services Div. (Pittsburgh, PA) and Southern Environmental Services Div. (Clear Lake, Texas). (Within Accession #243015).
4. Reviewed by: James D. Felkel  
Wildlife Biologist  
Ecological Effects Branch/HED
5. Date Reviewed: 9/4/80
6. Test Type: Fish acute 96-hour LC<sub>50</sub>
  - A. Test Species: Rainbow Trout (Salmo gairdneri)  
Fathead Minnows (Pimephales promelas)
7. Reported Results: The 96-hr. LC<sub>50</sub> for the rainbow trout is reported to be 0.87 mg/l (95% C.L. of 0.76-1.00 mg/l) and 2.25 mg/l (2.05-2.48 mg/l) for the fathead minnow.
8. Reviewer's Conclusions: The studies appear to be scientifically sound and meet the intent of Proposed Guidelines for a manufacturing-use. The reported LC<sub>50</sub> values indicate that the material tested is highly toxic to rainbow trout and moderately toxic to fathead minnows. Depending on future use patterns, a flow-through test may be advisable.

## Materials/Methods Reported

### A. Test Procedure

Dechlorinated Pittsburgh tap water (pretreatment included activated carbon filters and U.V. light) was used to prepare the test concentrations. The fish were purchased from commercial hatcheries. The trout ranged from 31-46 mm in length with an average weight of 0.84 g/fish and were acclimated to the test temperature of 10° (+1°) C. The juvenile fathead minnows ranged from 27-33 mm in length with an average weight of 0.29 g/fish and were acclimated to 17.5° (+1°) C and tested at 18°C. Food was withheld one day prior to, and during, all tests.

Rangefinder tests were conducted to estimate the upper and lower bounds of the concentration range expected to include the median lethal concentration and five concentrations spanning this range were selected for the definitive test.

All definitive treatment levels and controls were established in duplicate with 10 fish in each chamber. Test chambers were arranged randomly in temperature-controlled water baths and fish were randomly selected from the holding tank and placed two at a time in all chambers. Fish behavior and mortality were monitored daily with dead fish removed and preserved for length and weight measurements. Survivors were sacrificed and similarly measured after each bioassay. Water quality measurements included water temperature for all test concentrations every 24 hours; D.O. and pH for the high, middle, and low concentrations and controls at the start and finish of testing; plus the alkalinity, hardness, and conductivity for the highest concentrations and controls at the start of testing.

### B. Statistical Analysis

The Litchfield-Wilcoxon method was used to fit a line to mortality data plotted on log-probit paper.

## Discussion/Results Reported

### 1. Rainbow Trout

Test concentrations selected after rangefinder testing were 1, 1.3, 1.5, 1.7, and 2 mg/l. However, heavy mortality, including all fish dead in the upper three concentrations after 12 hours, necessitated the addition of three additional test levels: 0.3, 0.5, and 0.7 mg/l. All fish at the lowest two levels remained alive for the 96-hour period. Cumulative mortality was as follows:

6

Treatment Level (mg/l)

	<u>Time (hrs.)</u>			
	<u>24</u>	<u>48</u>	<u>72</u>	<u>96</u>
1.5	20	20	20	20
1.3	18	19	19	19
1.0	16	17	17	17
0.7	2	3	4	4
0.5	0	0	0	0
Control	0	0	0	0

This table shows that the majority of deaths at all treatment levels occurred within the first 24 hours. The 96-hour LC<sub>50</sub> was reported to be 0.87 mg/l (95% confidence limits of 0.76 and 1.00 mg/l).

2. Fathead Minnow

Definitive test concentrations selected after a rangefinder test were 3, 2.5, 2, 1.7, and 1 mg/l. Cumulative mortality was as follows:

Treatment Level (mg/l)

	<u>Time (hrs.)</u>			
	<u>24</u>	<u>48</u>	<u>72</u>	<u>96</u>
3	20	20	20	20
2.5	13	13	13	13
2	7	7	7	7
1.7	0	0	0	0
1	0	0	0	0
Control	0	0	0	0

All mortality occurred within the first 24 hours. The 96-hour LC<sub>50</sub> was reported to be 2.25 mg/l with 95% confidence limits of 2.48 and 2.05 mg/l.

3. Water Quality

At the start of testing with Rainbow Trout, D.O. was 10.5 mg/l at all test levels checked (ca. 93% saturation) and declined after 96 hours to levels ranging from 7.6 mg/l in the controls (ca. 62% saturation) to 9.8 mg/l at 1.0 mg/l test concentration (ca. 87% saturation). At the start of testing with fathead minnow, D.O. was 9.7 mg/l at all test levels checked (ca. 85% saturation) and declined after 96 hours to levels ranging from 5.2 mg/l (ca. 46% saturation) in the controls to 6.2 mg/l (ca. 54% saturation) at 1 mg/l test concentration.

pH was 6.7 - 6.8 at the start of all testing and declined slightly to 6.5 - 6.7 or remained constant by the end of the 96 hours in the test vessels. Conductivity level was 375  $\mu$ mhos in the trout experiment and 225 with the minnow experiment. Alkalinity was 44 and 34 mg/l (as CaCO<sub>3</sub>) in these two experiments, respectively; and hardness was 134 and 100 mg/l (as CaCO<sub>3</sub>), respectively.

7



## Reviewer's Evaluation

### A. Test Procedure

Procedures generally followed the 1978 Proposed Guidelines (subpart E) and approved protocols (Committee on Methods for Toxicity Tests with Aquatic Organisms, 1975). However, discrepancies include:

- 1) Fathead Minnows, with an average weight of 0.29 g/fish, were below the recommended range of 0.5-5g;
- 2) Food was withheld 1 day rather than 96 hours prior to testing;
- 3) The test temperature of 18°C for the fathead minnow was below the recommended 22 °C of the 1975 protocols; however, recent ASTM standard practices (E 729-80) permit a test temperature of 17 or 22°C.;
- 4) Information on storage of the test material was missing; and
- 5) Missing environmental conditions included the photoperiod/lighting and depth of solution in test chambers; and
- 6) D.O. was not measured every 48 hours in the control and the high, medium, and low toxicant concentrations. (These measurements were taken at the beginning and end of testing.)

Minor internal discrepancies within the report include:

- 1) Page 16 of the text states that 1 trout died at 1.3 mg/l and 3 died at 1.0 mg/l of the test chemical, but Table 16 indicates 2 and 4 deaths, respectively; The submitted LC<sub>50</sub> was confirmed using the data of Table 16; and
- 2) The description of water quality measurements taken on p.17 is inconsistent with the results in Table 22 in terms of the tests conducted. It is assumed that the tests conducted were those for which results were reported (Table 22).

### B. Statistical Analysis

A probit analysis by this reviewer of the cumulative mortality data submitted in Tables 16 and 17 gave LC<sub>50</sub> values and 95% confidence limits of 0.84 (0.76-0.91) mg/l for rainbow trout and 2.24 (2.11-2.38) mg/l for fathead minnow, closely approximating the submitted values.

8

C. Results/Discussion

The submitted results indicate that this compound is highly toxic to rainbow trout and moderately toxic to fathead minnows.

D. Conclusions

1. Category: Core
2. Rationale: While there were certain discrepancies from approved protocols as described above, the studies appear to be scientifically sound and meet the intent of the 1978 proposed subpart E guidelines for a manufacturing-use. Depending on future use patterns, a flow-through test may be advisable.
3. Repairability: N/A

```

*****
CONC.          NUMBER      NUMBER      PERCENT      BINOMIAL
              EXPOSED     DEAD        DEAD          PROB. (1)
1.5           20             20          100           9.53673
1.3           20             19          95.           2.0027
1             20             17          85.           .128841
.7            20             4           20.           .590897
.5            20             0           0             9.53674

```

THE BINOMIAL TEST SHOWS THAT .7 AND 1 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS .823179

```

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD
SPAN          G          LC50          95 PERCENT CONFIDENCE LIMITS
3             6.12738E-2    .829708      .756731      .909125

```

```

-----RESULTS CALCULATED USING THE PROBIT METHOD
ITERATIONS    G          H          GOODNESS OF FIT PROBABILITY
5             .111283     1          .674846

```

SLOPE = 10.7927  
95 PERCENT CONFIDENCE LIMITS = 7.19235 AND 14.393

LC50 = .836785  
95 PERCENT CONFIDENCE LIMITS = .760438 AND .913542

\*\*\*\*\*

SRU 1.300 UNTS.

RUN COMPLETE.

9001 data 3,2.5,2,1.7,1  
 9002 data 20,20,20,20,20  
 9003 data 20,13,7,0,0  
 run

80/09/24. 14.33.24.  
 BASIC PROGRAM S79LC50

DiHalo/ Fathead minnow Acc. #243015

\*\*\*\*\*

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.( PERCENT)
3	20	20	100	9.53674E-5
2.5	20	13	65.	13.1588
2	20	7	35.	13.1588
1.7	20	0	0	9.53674E-5
1	20	0	0	9.53674E-5

THE BINOMIAL TEST SHOWS THAT 1.7 AND 3 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 2.23607

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD

SPAN	G	LC50	95 PERCENT CONFIDENCE LIMITS	
4	4.90922E-2	2.22507	2.06362	2.42224

-----RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS	G	H	GOODNESS OF FIT PROBABILITY
8	.113443	1	.230874

SLOPE = 14.8701  
 95 PERCENT CONFIDENCE LIMITS = 9.86162 AND 19.8785

LC50 = 2.24066  
 95 PERCENT CONFIDENCE LIMITS = 2.10911 AND 2.38308

SRU 1.332 UNTS.

RUN COMPLETE.

11

DATA EVALUATION RECORD

1. Chemical: 5,5-dimethylhydantoin (No Shaughnessy number available)
2. Formulation: Test material not labelled as to purity or percent active ingredient. GLCC (Dr. John Sands) reports a.i. to be a minimum of 97%.
3. Citation: Horne, J.D., et al. 1980. 96-hour static bioassays using two Great Lakes Chemical Corporation compounds with three marine and three freshwater species. NUS Corporation--Northern Environmental Services Div. (Pittsburgh, PA) and Southern Environmental Services Div. (Clear Lake, Texas.) (Within Accession #243015.)
4. Reviewed by: James D. Felkel  
Wildlife Biologist  
Ecological Effects Branch/HED
5. Date Reviewed: 10/21/80
6. Test Type: Acute toxicity to estuarine and marine organisms
  - A. Test Species: Juvenile sheepshead minnow (Cyprinodon variegatus)  
Grass shrimp (Palaemonetes pugio)  
American oyster (Crassostrea virginica)
7. Reported Results: The LC<sub>50</sub> for sheepshead minnow is 8,100 mg/l, 1,300 mg/l for grass shrimp, and 13,300 mg/l for American oyster.
8. Reviewer's Conclusions: The reported results indicate that this degradation product is practically non-toxic to the species tested. While the studies may be scientifically sound, further information cited in the Repairability section below is needed before the studies on sheepshead minnow and grass shrimp could satisfy a potential requirement for this type of test. Flow-through tests may be advisable. The test on the American oyster could not satisfy a potential requirement for an acute toxicity test with this material.

## I. Materials/Methods Reported

### A. Test Procedures

A synthetic seasalt ("Biocrystals Marinemix") was dissolved in deionized water with a salinity of 20(+2) ppt (o/oo). This water was used in the culture/acclimation tanks, the test tanks, and to prepare the stock solutions.

The sheepshead minnows were spawned and reared at the NUS laboratory. They were 45(+2) days old, averaged 0.13 g/individual, and ranged from 10-26 mm in length at the start of testing. The grass shrimp were collected locally in Texas and acclimated to a final temperature of 22°C and 20 ppt (o/oo) salinity. They averaged 0.18 g/individual and ranged from 17-38 mm in length. The oysters were purchased from a commercial lease-bed operator. The average wet-tissue weight of the oysters was 3.8 g/individual.

Rangefinder tests were conducted to determine the appropriate definitive test concentrations for all species. Five test levels were selected for the tests with sheepshead minnows (10, 6.5, 4.2, 2.7, and 1.8 g/l), seven for the grass shrimp (10, 6.5, 4.2, 2.7, 1.8, 1.17, and 0.76 g/l), and five for the American oysters (20, 17, 14.5, 12.25, and 10.4 g/l). Each definitive bioassay included a control and all treatment levels were established in duplicate. The minnows and shrimp were tested in 3 l of the test solution in 1-gal. jars with 10 organisms/container. Test containers were arranged randomly in the test area and organisms were assigned randomly to the containers.

Behavior and mortality were checked daily. Temperature, D.O., and salinity were measured daily for all test solutions.

### B. Statistical Analysis

The Litchfield-Wilcoxon method was used to fit a line to mortality data plotted on log-probit paper for the shrimp and oyster data and the straight-line interpolation method described in "Standard Methods, 14th ed." (APHA, 1975) was used for the minnow data.

## II. Discussion/Results Reported

### A. Sheepshead Minnow

100% mortality occurred at the 10 g/l level with no mortality at or below the 6.5 g/l test level. The submitted LC<sub>50</sub> for this species is 8.1 g/l. 95% confidence limits could not be determined using the analytical procedure employed.

### B. Grass shrimp

100% mortality occurred at the highest four test levels after 96 hours and partial mortalities occurred at the lower three levels

during the same period. The submitted  $LC_{50}$  is 1.3 g/l (95% C.I. = 1.1-1.6 g/l).

C. American oyster

Partial mortalities occurred at all concentration levels tested. The submitted  $LC_{50}$  is 13.3 g/l (95% C.I. = 10.3-17.5 g/l).

D. Water Quality

Tables of water quality data submitted indicate that temperature rose from 20°C at the start of testing to a high of 26°C after 24 hours in all the sheepshead minnow test vessels while salinity levels ranged from 20 ppt (o/oo) at the start to 22-31 ppt (o/oo) by the end of testing. D.O. levels declined from 7.2-7.6 mg/l at the start to 2.2-3.9 mg/l after 24 hours in the minnow test vessels before generally increasing as the test progressed. Similar variation in these parameters occurred in the grass shrimp test vessels while water quality was considerably more constant in the oyster bioassays.

III. Reviewer's Evaluation

A. Test procedure

Discrepancies from accepted protocols (Stephen, 1975) include:

- 1) sheepshead minnows averaged 0.13 g/individual rather than 0.5-5g and the standard length of the largest fish was more than twice that of the shortest;
- 2) oysters are generally used in embryolarvae or shell deposition studies rather than in adult 96-hr.  $LC_{50}$  studies; and
- 3) information was not provided on the photoperiod.

B. Statistical Analysis

An analysis by this reviewer of the cumulative mortality data submitted provided  $LC_{50}$  values that generally approximated the submitted values except for the 95% C.I. for the oyster  $LC_{50}$ . For the sheepshead minnow data, it was necessary to rely on the binomial test since where there are less than two concentrations at which the percent dead is between 0 and 100, neither the moving average nor the probit method can give any statistically sound results (analyses attached).

C. Results/Discussion

Discrepancies from accepted protocols (Stephen, 1975) include:

- 1) water temperature varied substantially more than 1°C from the initial temperature in all test vessels;

- 2) dissolved oxygen was under 60% saturation at some point during the 1st 48 hours in all the sheepshead minnow and grass shrimp test vessels; and
- 3) in many of the minnow and shrimp test vessels, D.O. increased sharply from the 24-hour check to the later checks in the bioassays; while this is not a discrepancy per se, aeration during the tests, if it occurred, would be.

#### D. Conclusions

1. Category: Supplemental
2. Rationale: At present, neither marine/estuarine organism toxicity testing nor tests on degradation products, such as the material tested here, are required for the present registration proposals and thus "Core" status (satisfying an existing requirement) could not apply.

Also, there were substantial discrepancies from accepted protocols regarding water temperature and D.O. and it also is not clear that aeration was not conducted. The American oyster is not an acceptable species for the type of acute toxicity test conducted.

3. Repairability: Although this degradate product appears to be relatively non-toxic, the water temperature and D.O. patterns must be explained and adequately defended and it must be verified whether aeration took place during the tests. If this is done and if tests on degradation products are required for marine/estuarine species in the future, the tests on sheepshead minnow and grass shrimp may satisfy such a requirement. The test on the American oyster would not be repairable for the reasons cited earlier. Flow-through tests are generally recommended for studies using salt water since salt water is particularly vulnerable to fouling.



data 5  
 data 10,6.5,4.2,2.7,1.8  
 data 20,20,20,20,20  
 data 20,0,0,0,0

80/09/24. 14.40.34.  
 BASIC PROGRAM S79LC50

5.5-Dimethylhydantoin/Sheepshead Minnow \*\*\*\*\* ACC #243015 \*\*\*\*\*

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.(PERCENT)
10	20	20	100	9.53674E-5
6.5	20	0	0	9.53674E-5
4.2	20	0	0	9.53674E-5
2.7	20	0	0	9.53674E-5
1.8	20	0	0	9.53674E-5

A. THE BINOMIAL TEST SHOWS THAT 6.5 AND 10 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 8.06226

WHEN THERE ARE LESS THAN TWO CONCENTRATIONS AT WHICH THE PERCENT DEAD IS BETWEEN 0 AND 100, NEITHER THE MOVING AVERAGE NOR THE PROBIT METHOD CAN GIVE ANY STATISTICALLY SOUND RESULTS.  
 \*\*\*\*\*

SRU 1.308 UNTS.

RUN COMPLETE.

run

80/09/25. 12.43.53.  
BASIC PROGRAM S79LC50

5,5-Dimethylhydantoin/Crassostrea virginica) Acc. #243015;

\*\*\*\*\*

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.(PERCENT)
20	20	11	55.	41.1901
17	20	15	75	2.06947
14.5	20	15	75	2.06947
12.25	20	8	40.	25.1722
10.4	20	4	20.	.590897

THE BINOMIAL TEST SHOWS THAT 10.4 AND +INFINITY CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 12.839

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD

SPAN	G	LC50	95 PERCENT CONFIDENCE LIMITS	
3	.299595	12.8431	11.1723	14.304

-----RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS	G	H	GOODNESS OF FIT PROBABILITY
4	3.73209	3.20942	2.20054E-2

SINCE THE PROBABILITY IS LESS THAN 0.05, RESULTS CALCULATED USING THE PROBIT METHOD PROBABLY SHOULD NOT BE USED.

SLOPE = 3.86461  
95 PERCENT CONFIDENCE LIMITS = -3.60128 AND 11.3305

LC50 = 13.8122  
95 PERCENT CONFIDENCE LIMITS = 0 AND +INFINITY

\*\*\*\*\*

SRU 1.336 UNTS.

RUN COMPLETE.

17

TERMINAL: 37, P 33/TTY  
 USERS, TYPE\*WRITEUP(NOTE)\* 80/09/25.  
 RECOVER /SYSTEM: basic,old,s79lc50  
 READY.

9000 data 7  
 9001 data 10,6.5,4.2,2.7,1.8,1.17,0.76  
 9001 data 10,6.5,4.2,2.7,1.8,1.17,0.76  
 9002 data 20,20,20,20,20,20,20  
 9003 data 20,20,20,20,16,7,3  
 run

80/09/25. 12.33.02.  
 BASIC PROGRAM S79LC50

\*\*\*\*\*

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB. (PERCENT)
10	20	20	100	9.53674E-5
6.5	20	20	100	9.53674E-5
4.2	20	20	100	9.53674E-5
2.7	20	20	100	9.53674E-5
1.8	20	16	80.	.590897
1.17	20	7	35.	13.1588
.76	20	3	15.	.128841

THE BINOMIAL TEST SHOWS THAT .76 AND 1.8 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 1.34439

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD

SPAN	G	LC50	95 PERCENT CONFIDENCE LIMITS	
3	8.50121E-2	1.27142	1.0807	1.45953

-----RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS	G	H	GOODNESS OF FIT PROBABILITY
7	.119289	1	.896154

SLOPE = 5.8511  
 95 PERCENT CONFIDENCE LIMITS = 3.83023 AND 7.87197

LC50 = 1.25593  
 95 PERCENT CONFIDENCE LIMITS = 1.07766 AND 1.44501

SRU 1.317 UNTS.

RUN COMPLETE.

DATA EVALUATION RECORD

1. Chemical: 5,5-dimethylhydantoin (No Shaughnessy number available)
2. Formulation: Test material not labelled as to purity or percent active ingredient. GLCC (Dr. John Sanos) reports a.i. to be a minimum of 97%.
3. Citation: Horne, J.D., et al. 1980. 96-hour static bioassays using two Great Lakes Chemical Corporation compounds with three marine and three freshwater species. NUS Corporation--Northern Environmental Services Div. (Pittsburgh, PA) and Southern Environmental Services Div. (Clear Lake, Texas.) (Within Accession #243015.)
4. Reviewed by: James D. Felkel  
Wildlife Biologist  
Ecological Effects Branch/HED
5. Date Reviewed: 10/23/80
6. Test Type: 96-hour fish and 48-hour aquatic invertebrate LC50
  - A. Test Species: Rainbow trout (Salmo gairdneri)  
Fathead minnows (Pimephales promelas)  
Daphnia magna
7. Reported Results: LC50 values are 12.7 g/l for rainbow trout, 14.2 g/l for fathead minnow, and 6.2 g/l for Daphnia magna.
8. Reviewer's Conclusions: The reported results indicate that the test material is practically non-toxic to the species tested. The tests appear to be scientifically sound and should acute toxicity tests be required in the future for these species with this degradation product as the test material, the tests with the fathead minnow and Daphnia magna would satisfy such a requirement. Further information regarding the aeration, as noted under Repairability below, would be needed before the trout study would satisfy such a requirement. Depending on future use proposals, flow-through testing may also be advisable.

## I. Materials/Methods Reported

### A. Test Procedures

Dechlorinated Pittsburgh tap water (pretreatment included activated carbon filters and U.V. light) was used to prepare the test concentrations. The fish were purchased from commercial hatcheries and the Daphnia magna from Wards Natural Science Estab., Inc. The trout ranged from 32-58 mm with an average weight of 2.18 g/fish. The minnows were 21-31 mm in length with an average weight of 0.51 g/individual. The fish were not fed one day prior to, or during, testing.

Rangefinder tests were conducted to estimate the upper and lower bounds of the concentration range expected to include the median lethal concentration and five concentrations spanning this range were selected for the definitive test.

All definitive treatment levels and controls were established in duplicate with 10 organisms in each chamber. Test chambers were arranged randomly in temperature-controlled water baths and fish were randomly selected from the holding tank and placed two at a time in all chambers. Fish behavior and mortality were monitored daily with dead fish removed and preserved for length and weight measurements. Survivors were sacrificed and measured after each bioassay.

Water quality measurements in the tests included (1) temperature for all test concentrations every 24 hours for the fish test vessels and at the start and end of testing for the Daphnia containers; (2) D.O., pH, alkalinity, hardness, and conductivity for the control and high, middle, and low fish test concentrations (D.O. and pH every 24 hours and the last three parameters at the start and end of testing) and; (3) D.O. and pH for the control and high, middle, and low Daphnia test concentrations at the start and end of testing and alkalinity, hardness, and conductivity for the control and high Daphnia test concentration at the start of testing.

It is reported that "gentle aeration" was required in the test with rainbow trout from the 48-hr. period to the end of the test and that this was the only freshwater test in which this was done.

Initial test temperatures were 10.0-10.9°C for the trout, 17.3-17.5°C for the minnows, and 20°C for the Daphnia. Treatment levels were 25, 15, 12.5, 10, and 5 g/l for the trout; 15, 10, 7, 5, and 3 g/l for the Daphnia; and 20, 15, 12.5, 10 and 5 g/l for the minnows.

### B. Statistical Analysis

The Litchfield-Wilcoxon method was used to fit a line to mortality data plotted on log-probit paper.

## II. Discussion/Results Reported

### A. Rainbow trout

Cumulative mortality was as follows:

<u>treatment level (g/l)</u>	<u>time (hrs.)</u>			
	24	48	72	96
25	9	17	20	20
15	0	3	11	17
12.5	0	0	1	8
10	0	0	0	2
5	0	0	0	0
Control	0	0	0	0

An  $LC_{50}$  of 12.7 g/l (95% C.I. = 11.3-14.2 g/l) is reported.

D.O. levels dropped from a starting level of 10.4-10.6 mg/l to 2.2-5.5 after 48 hours and "gentle aeration" was conducted. Other parameters remained relatively constant.

### B. Fathead minnow

Cumulative mortality was as follows:

<u>treatment level (g/l)</u>	<u>time (hrs.)</u>			
	24	48	72	96
20	20	20	20	20
15	11	13	14	14
12.5	3	3	3	3
10	0	0	0	0
5	0	0	0	0
Control	0	0	0	0

An  $LC_{50}$  of 14.2 g/l (C.I. = 13.2-15.3 g/l) is reported. Water quality parameters were within acceptable limits.

### C. Daphnia magna

Cumulative mortality was as follows:

<u>treatment level (g/l)</u>	<u>time (hrs.)</u>			
	12	24	36	48
15	20	20	20	20
10	9	12	17	20
7	0	5	9	11
5	0	1	6	7
3	0	0	0	0
Control	0	0	0	0

An  $LC_{50}$  of 6.2 g/l (95% C.I. = 5.3-7.3 g/l) is reported. Water quality parameters were within acceptable limits.

### III. Reviewer's Evaluation

#### A. Test Procedures

Procedures generally followed the 1978 proposed subpart E guidelines and approved protocols (Stephen, 1975). The principal discrepancy was an overloading of the trout test vessels (1.36 g/l). This resulted in the low D.O. levels and necessitated aeration. Aeration is not permitted by accepted protocols unless it can be shown that it did not reduce toxicant levels (through volatilization) by more than 20% from the levels found in non-aerated vessels. Other discrepancies included:

- 1) food was withheld 1 day rather than 96 hours prior to testing;
- 2) test temperature for the fathead minnow was below that of the 1975 protocols but consistent with recently-published ASTM standard practices (E 729-80);
- 3) the test temperatures for the trout varied by slightly more than 1°C in some vessels;
- 4) information on the storage of the test material was missing; and
- 5) missing environmental conditions included the photoperiod and depth of solution in the test chambers.

#### B. Statistical Analysis

Probit analyses of the submitted data by this reviewer gave LC<sub>50</sub> values closely approximating the submitted values (analyses attached).

#### C. Results/Discussion

The submitted results indicate that this degradation is practically non-toxic to the species tested.

#### D. Conclusions

1. Category: Supplemental
2. Rationale: Guidelines do not require fish and wildlife testing on degradation products for the present registrations (1729-122 and 1729-123). Aeration in the test with rainbow trout, unless it can be shown that this would not have affected toxicant levels available to the fish by more than 20%, would prevent this study from meeting "Core" status.
3. Repairability: If 96-hour acute toxicity studies with an aquatic invertebrate and a warmwater fish species are required

22

for this degradation product in the future, these tests would meet "Core" status. Only if it can be shown that the aeration of the trout test vessels did not substantially effect toxicity, as noted above, would the trout meet "Core" status if this test is required in the future. Depending on future use patterns, a flow-through test may also be advisable.



11.16.02.  
PROGRAM S79LC50

Dimethylhydantoin/Rainbow trout Acc. #243015;

NO.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.(PERCENT)
25	20	20	100	9.53674E-5
15	20	17	85.	.128841
12.5	20	8	40.	25.1722
10	20	2	10.	2.01225E-2
5	20	0	0	9.53674E-5

THE BINOMIAL TEST SHOWS THAT 10 AND 15 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 12.9826

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD  
 SPAN G LC50 95 PERCENT CONFIDENCE LIMITS  
 3 .112353 12.7795 11.4769 14.0927

-----RESULTS CALCULATED USING THE PROBIT METHOD  
 ITERATIONS G H GOODNESS OF FIT PROBABILITY  
 7 .189957 1 .935389

SLOPE = 13.2322  
 95 PERCENT CONFIDENCE LIMITS = 7.46504 AND 18.9993

LC50 = 12.7575  
 95 PERCENT CONFIDENCE LIMITS = 11.8959 AND 13.7326

SRU 1.324 UNTS.

RUN COMPLETE.

data 15,10,7,5,3  
data 20,20,11,7,0

80/09/25. 13.24.10.  
BASIC PROGRAM S79LC50

5,5-Dimethylhydantoin/Daphnia magna 48-hr. Bioassay; Acc.#243015;  
\*\*\*\*\*

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.(PERCENT)
15	20	20	100	9.53674E-5
10	20	20	100	9.53674E-5
7	20	11	55.	41.1901
5	20	7	35.	13.1588
3	20	0	0	9.53674E-5

THE BINOMIAL TEST SHOWS THAT 3 AND 10 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 6.44038

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD  
SPAN 3 G 5.39519E-2 LC50 5.97328 95 PERCENT CONFIDENCE LIMITS 5.38581 6.62557

-----RESULTS CALCULATED USING THE PROBIT METHOD  
ITERATIONS 7 G .136787 H 1 GOODNESS OF FIT PROBABILITY .274481

SLOPE = 7.934  
95 PERCENT CONFIDENCE LIMITS = 4.99963 AND 10.8684

LC50 = 6.05161  
95 PERCENT CONFIDENCE LIMITS = 5.354 AND 6.76519  
\*\*\*\*\*

SRU 1.334 UNTS.

RUN COMPLETE.

bye

25

9000 data 5  
9001 data 20,15,12.5,10,5  
9003 data 20,14,3,0,0  
run

80/09/25. 13.19.35.  
BASIC PROGRAM S79LC50

5,5-Dimethylhydantoin/Fathead minnow 96-hr. LC50 Acc.#243015:  
\*\*\*\*\*

CONC. (g/l)	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.(PERCENT)
20	20	20	100	9.53674E-5
15	20	14	70.	5.76591
12.5	20	3	15.	.128841
10	20	0	0	9.53674E-5
5	20	0	0	9.53674E-5

THE BINOMIAL TEST SHOWS THAT 12.5 AND 20 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 14.0765

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD

SPAN	G	LC50	95 PERCENT CONFIDENCE LIMITS	
3	6.23526E-2	14.2861	13.4176	15.307

-----RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS	G	H	GOODNESS OF FIT PROBABILITY
8	.2253	1	.996363

SLOPE = 20.4134  
95 PERCENT CONFIDENCE LIMITS = 10.724 AND 30.1028

LC50 = 14.1019  
95 PERCENT CONFIDENCE LIMITS = 13.3841 AND 14.9903

\*\*\*\*\*

SRU 1.324 UNTS.

RUN COMPLETE.

26

DATA EVALUATION RECORD

1. Chemical: Cited in report as 1-bromo-3-chloro-5, 5-dimethylhydantoin (DiHalo) (Shaughnessy #006315.)  
Reported by Great Lakes Chemical Co. (Dr. John Sands)  
as "Bromochloro-5, 5-diemthylhydantoin", an equal-molar  
combination of:

- 1) Dibromo-5, 5-dimethylhydantoin (Shaughnessy #006317)
- 2) Dichloro-5, 5-dimethylhydantoin (Shaughnessy #028501)
- 3) 1-bromo-3-chloro-5, 5-dimethylhydantoin (Shaughnessey #006315)
- 4) 3-bromo-1-chloro-5, 5-dimethylhydantoin (Shaughnessy Number not avail.)

2. Formulation: Report: Test material not labelled as to its percent  
active ingredient.  
GLCC (Dr. Sands): 92.2% active ingredient.

3. Citation: Horne, J.D., et al. 1980. 96-hour static bioassys using  
two Great Lakes Chemical Corportation compounds with three marine and three  
freshwater species. NUS Corporation--Northern Environmental Services Div.  
(Pittsburgh, PA) and Southern Environmental Services Div. (Clear Lake,  
Texas). (Within Accession #243015).

4. Reviewed by: James D. Felkel  
Wildlife Biologist  
Ecological Effects Branch/HED

5. Date Reviewed: 10/16/80

6. Test Type: Acute toxicity (96-hour) to estuarine and marine organisms

A. Test Species: Juvnile sheepshead minnow (Cyprinodon variegatus) Grass shrimp (Palaemonetes pugio) American Oyster (Crassostrea virginica)

7. Reported Results:

The 96-hour LC<sub>50</sub> for sheepshead minnow is 20 mg/l (95% C.I. = 17-23 mg/l) and 13 mg/l (95% C.I. = 10.1-16.6 mg/l) for the grass shrimp. No LC<sub>50</sub> was calculable for the American oyster.

8. Reviewer's Conclusions:

The studies appear to be scientifically sound. However, further information cited in the Repairability section below is needed before the tests on the sheepshead minnow and grass shrimp could meet proposed subpart E guideline requirements. The test on the American oyster cannot meet the intent of the guidelines. The submitted results indicate that the material tested is "slightly toxic" to the minnows and shrimps, following EPA-accepted toxicity category criteria. Depending on future use patterns, flow-through tests may be advisable.

11.09.37.  
PROGRAM S79LC50

11.09.37.  
PROGRAM S79LC50

Crassostrea virginica Acc. 243015; Shaugh. #006315/ 006317/028501

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.(PERCENT)
640	20	2	10.	2.01225E-2
200	20	2	10.	2.01225E-2
112	20	1	5.	2.00272E-3
64	20	1	5.	2.00272E-3
36	20	2	10.	2.01225E-2
20	20	2	10.	2.01225E-2
11.2	20	0	0	9.53674E-5

THIS DATA SET DOES NOT MEET THE CRITERIA ESTABLISHED BY THE COMMITTEE ON METHODS FOR TOXICITY TESTS WITH AQUATIC ORGANISMS BECAUSE NO PERCENT DEAD IS GREATER THAN 65 PERCENT.

NEITHER THE BINOMIAL TEST NOR THE MOVING AVERAGE METHOD CAN GIVE ANY RESULTS FOR THIS DATA SET. EITHER THE HIGHEST CONCENTRATION KILLED LESS THAN 50 PERCENT OR THE LOWEST KILLED MORE THAN 50. IF THE PROBIT SLOPE IS NEGATIVE, ENTER DATA AGAIN USING NUMBER ALIVE INSTEAD OF NUMBER DEAD.

-----RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS 5 G 5.58841 H 1 GOODNESS OF FIT PROBABILITY .743748

SLOPE = .235321  
95 PERCENT CONFIDENCE LIMITS = -.320974 AND .791617

LC50 = 1.31570E+8  
95 PERCENT CONFIDENCE LIMITS = 5065.44 AND +INFINITY

SRU 1.261 UNTS.

RUN COMPLETE.

ILLEGAL PARAMETER.

28

data 52,36,20,11.2,6.4,3.6,2.0  
 data 20,20,20,20,20,20,20  
 data 20,20,14,9,0,0,0

run

80/09/25. 13.00.37.  
 BASIC PROGRAM S79LC50

DiHalo/Palaemonetes pugio Acc. #243015; Shaugh. #006315 /006317/028501

\*\*\*\*\*

CONC. (mg/l)	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB.(PERCENT)
52	20	20	100	9.53674E-5
36	20	20	100	9.53674E-5
20	20	14	70.	5.76591
11.2	20	9	45.	41.1901
6.4	20	0	0	9.53674E-5
3.6	20	0	0	9.53674E-5
2	20	0	0	9.53674E-5

THE BINOMIAL TEST SHOWS THAT 6.4 AND 36 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 12.549

-----RESULTS CALCULATED USING THE MOVING AVERAGE METHOD

SPAN	G	LC50	95 PERCENT CONFIDENCE LIMITS	
6	2.67807E-2	12.8789	10.8127	15.407

-----RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS	G	H	GOODNESS OF FIT PROBABILITY
8	.109279	1	.531515

SLOPE = 5.27602  
 95 PERCENT CONFIDENCE LIMITS = 3.5319 AND 7.02014

LC50 = 13.8465  
 95 PERCENT CONFIDENCE LIMITS = 11.6506 AND 16.45

\*\*\*\*\*

SRU 1.297 UNITS.

RUN COMPLETE.

29

11.2, 64, 36, 20, 11.2  
 20, 20, 20, 20  
 20, 20, 20, 10, 0  
 20, 20, 20, 20, 20, 20

12.52.05.  
 PROGRAM S79LC50

Halalo/ Juv. Sheepshead minnow Acc. #243015; Shaugh. #006315/006317/028501  
 \*\*\*\*\*

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB. (PERCENT)
200	20	20	100	9.53674E-5
112	20	20	100	9.53674E-5
64	20	20	100	9.53674E-5
36	20	20	100	9.53674E-5
20	20	10	50	58.8099
11.2	20	0	0	9.53674E-5

THE BINOMIAL TEST SHOWS THAT 11.2 AND 36 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS SINCE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 20.

WHEN THERE ARE LESS THAN TWO CONCENTRATIONS AT WHICH THE PERCENT DEAD IS BETWEEN 0 AND 100, NEITHER THE MOVING AVERAGE NOR THE PROBIT METHOD CAN GIVE ANY STATISTICALLY SOUND RESULTS.  
 \*\*\*\*\*

SRU 1.285 UNTS.

RUN COMPLETE.

30

- 4) on several occasions, such as with the sheepshead minnow 20 mg/l and control vessels, the grass shrimp 20 mg/l and 11.2 mg/l vessels, and the oyster 11.2 mg/l and control vessels, D.O. increased from the 48-hour sample to the 96-hour sample. The cause of this increase should be clarified since aeration during the tests is strictly prohibited; and
- 5) a reliable LC<sub>50</sub> was not calculable for the American oyster.

The results indicate that the material tested is "slightly toxic" to sheepshead minnows and grass shrimps, following EPA-accepted toxicity category criteria..

#### D. Conclusions

1. Category: Supplemental
2. Rationale: Estuarine and marine organism toxicity tests are required to support the registration of a formulated product where such product is intended for direct application to the marine or estuarine environment or if it may be expected to enter this environment in significant concentrations because of its use or mobility pattern. Since such studies are not a requirement for the proposed registrations (1729-22 and 1729-23), they could not technically be considered "Core" (i.e., satisfying an existing guideline requirement).

Also, there were some discrepancies from accepted protocols (see Reviewer's Evaluation of procedures and results above). In particular, it should be clarified whether there was any aeration during the tests, and if not, what the cause of the D.O. pattern referred to in III. C. 4. above was. The acute toxicity results submitted for the American oyster were not suitable for calculation of a reliable LC<sub>50</sub> value; this species is not generally acceptable for such tests and is more suited for embryolarvae and shell deposition studies.

3. Repairability: a.) Sheepshead minnow and grass shrimp tests - If the temperature and D.O. pattern discrepancies can be adequately defended and if it can be verified that no aeration took place during the bioassays, then these tests may be suitable for Core status. [If acute toxicity tests on estuarine and marine organisms are required in the future, subpart E guidelines specify that a 96-hour LC<sub>50</sub> on a species of crab and either a 48-hour EC<sub>50</sub> for mollusc embryolarvae or a 96-hour EC<sub>50</sub> mollusc shell deposition also be provided (using the technical grade of the active ingredient). Flow-through tests may also be required since salt water is particularly vulnerable to fouling in static tests and since the test material (in minnow appears to be active for a very short period.] test)

- 6.) American oyster acute toxicity test is not repairable for the reasons noted above.



### III. Reviewer's Evaluation

#### A. Test Procedure

Discrepancies from accepted protocols (Stephan, 1975) include:

- 1) sheepshead minnows averaged 0.05 g/individual rather than 0.5 - 5 g and the standard length of the largest fish was more than twice that of the shortest;
- 2) oysters are generally used in embryolarvae or shell deposition studies rather than adult 96-hour LC<sub>50</sub> studies; and
- 3) information was not provided on the photoperiod.

#### B. Statistical Analysis

An analysis by this reviewer of the cumulative mortality data submitted confirmed the two LC<sub>50</sub> values reported and the fact that a reliable LC<sub>50</sub> value could not be calculated for the American oyster. However, it should be noted that the LC<sub>50</sub> value for the sheepshead minnow is more approximate than for the grass shrimp since there were less than two concentrations at which the percent dead was between 0 and 100.

#### C. Results/Discussion

Discrepancies from accepted protocols (Stephan, 1975) include:

- 1) with the sheepshead minnow and American oyster, water temperature occasionally varied by more than 1°C from the selected temperature (maximum of 3°C above starting temperature with the minnows and 2°C with the oysters);
- 2) D.O. was less than 60% saturation during at least part of the 1st 48 hours of testing with
  - a.) the sheepshead minnow test solution in the 20 mg/l test vessel;
  - b.) the grass shrimp test solutions in the 11.2, 6.4, 3.6, 2.0 mg/kg, and control vessels; and
  - c.) the American oyster test solution in the 11.2 mg/l and control test vessels.
- 3) D.O. was less than 40% saturation during at least part of the 2nd 48 hours of testing with the grass shrimps test solutions in the 3.6, 2.0 mg/l and the control vessels.

32

3. American Oyster

<u>Treatment Level</u> (mg/l)	<u>Time (hrs.)</u>			
	24	48	72	96
640	0	1	2	2
200	0	0	0	2
112	0	0	1	1
64	0	0	0	1
36	0	1	1	2
20	0	0	1	2
11.2	0	0	0	0
Control	0	0	0	0

Since greater than 50% mortality did not occur at any treatment level, an LC<sub>50</sub> value could not be calculated.

B. Water Quality1. Sheepshead Minnow

Water temperature was 22-23°C in all vessels at the start of testing, rose to 24°C in all vessels after 24 hours, rose to 25°C after 48-72 hours in all vessels for which measurements were taken (i.e., those still containing live fish), and then returned to 24°C after 96 hours in these latter vessels.

Dissolved Oxygen was 6.5-7.4 mg/l at the start of testing and declined to a low of 4.0 mg/l at 48 hours in the 20 mg/l test vessel. Salinity was constant at 21 ppt (0/00) except in the 200 mg/l vessel where it rose to 22 ppt (0/00).

2. Grass Shrimp

Water temperature was 19-20°C throughout the test. D.O. at the start of testing was 7.9-8.0 mg/l in all vessels and declined to a low of 2.5 mg/l after 96 hours in the 2 mg/l test vessel. Salinity was constant at 20 ppt (0/00).

3. American Oyster

Water temperature ranged from 22-24°C during the test. D.O. at the start of testing was 7.0 - 7.2 mg/l and declined to a low of 2.7 mg/l at 48 hours in the control group. Salinity ranged from 20-22 ppt.

1. Sheepshead Minnow

<u>Treatment Level</u> (mg/l)	<u>Time (hrs.)</u>			
	← 24	48	72	96
200	20	20	20	20
112	20	20	20	20
64	20	20	20	20
36	20	20	20	20
20	4	10	10	10
11.2	0	0	0	0
Control	0	0	0	0

The LC<sub>50</sub> is reported to be 20 mg/l (95% confidence interval of 17-23 mg/l).

2. Grass Shrimp

<u>Treatment Level</u> (mg/l)	<u>Time (hrs.)</u>			
	← 24	48	72	96
52	20	20	20	20
36	20	20	20	20
20	7	11	13	14
11.2	2	7	9	9
6.4	0	0	0	0
3.6	0	0	0	0
2.0	0	0	0	0
Control	0	0	0	1

The LC<sub>50</sub> is reported to be 13 mg/l (95% C.I. of 10.1 - 16.6 mg/l)

The one control mortality resulted from the shrimp escaping the test vessel.

## Materials/Methods Reported

### A. Test procedure

A synthetic seasalt ("Biocrystals Marinemix") was dissolved in deionized water to produce water with a salinity of 20 (+2) ppt (0/00). This water was used in the culture/acclimation tanks, the test tanks, and to prepare the stock solutions.

The sheepshead minnows were spawned and reared at the NUS laboratory. They were 29 (+2) days old, averaged 0.05 g/ individual, and ranged from 8-22mm in length at the start of testing. The grass shrimp were collected locally in Texas and acclimated to a final temperature of 22°C and 20 ppt (0/00) salinity. They averaged 0.18g/ individual and ranged from 17-38 mm in length (rostrum to telson). The oysters were purchased from a commercial lease-bed operator. The average wet-tissue weight of the oyster was 3.8 g/individual.

Rangefinder tests were conducted to determine the appropriate definitive test concentrations for all species. Six test levels were selected for the tests with sheepshead minnows (200, 112, 64, 36, 20, and 11.2 mg/l) seven for the grass shrimp (52, 36, 20, 11.2, 6.4, 3.6, and 2.0 mg/l), and seven for the American oysters (640, 200, 112, 64, 36, 20, and 11.2 mg/l.) Each definitive bioassay included a control and all treatment levels were established in duplicate. The minnows and shrimps were tested in 3 l of test solution in 1-gal widemouth jars with 10 organisms/container. The oysters were tested in 10-gal aquaria containing 30 l of test solution and 10 organisms/container. Test containers were arranged randomly in the test area and organisms were assigned randomly to the containers.

Behavior and mortality were checked daily. Temperature, D.O., and salinity were measured daily for all test solutions.

### B. Statistical Analysis

The Litchfield-Wilcoxon method was used to fit a line to mortality data plotted on log-probit paper.

## Discussion/Results Reported

### A. Cumulative Mortality

Cumulative mortality (number of individuals out of 20 at each test level) was as follows:

37