

US EPA ARCHIVE DOCUMENT

PD-470
TRK-002481

Alfred 5500
Beth
Comings
Coit

002481

November 27, 1967

File: PD # 8F0654

Pesticide Control Branch (1-13)

Dr. H. R. Gitten
Division of Toxicological Evaluation
Pesticide Review Branch (1-290)

Evaluation of toxicology data presented in support of petition for residue tolerances of Rendax-1 (mixture of o-chloro-m,m-diethyl acetamide and trichlorobenzyl chloride).

PESTICIDE REGISTRY No. 870 658

Monsanto Chemical Company
St. Louis, Missouri
(314-765)

115312


Two product formulations shown below are involved.

Rendax-1 Granular consisting of:

Active Ingredients

<u>o</u> -chloro- <u>m</u> , <u>m</u> -diethylacetamide (CSA)	11.7
Trichlorobenzyl chloride (TCBC)	23.3

Inert Ingredients




Rendax-1 Liquid consisting of:

Active Ingredients

<u>o</u> -chloro- <u>m</u> , <u>m</u> -diethylacetamide (CSA)	23.7
Trichlorobenzylchloride (TCBC)	59.3

Inert Ingredients



The petitioner requests negligible tolerances (0.5 ppm) with respect to each of the active ingredients when used, pre-emergence, on field corn, silage corn, sweet corn and popcorn as a single application at rates of 3.75 lb per acre for Rendax-1 granular and 4.5 quarts per acre for the Rendax-1 Liquid.

Received
FSA/Pest. Br.

DEC 8 1967

BEST AVAILABLE COPY

10/270

Inert ingredients

November 27, 1967

Both Dactoxol and one of its active ingredients N-methyl-2-(2-dimethylaminoethyl)-acetamide (DMEM) have been previously registered with EPA on a no residue basis.

Toxicological Data Report-1

1. Acute Oral Toxicity in Mice Younger lab Y59-40 ; 08/4/59

No. of animals used	-	22 (12 males & 10 females)
Method of administration	-	Intubation
Dosage levels	-	871, 1000, 1150, 1320 mg/kg

Results

Mortality with increasing dosage was found to be 0/3, 3/7, 3/5, 4/5. The oral LD₅₀ was determined as 1065 mg/kg with limits of 537-1780 mg/kg. Survival time was overnight to forty-eight hours, most deaths occurring overnight. Symptoms included severe diarrhea, weakness and tremors. At autopsy on selected animals which died, gross observation showed inflammation of the gastric mucosa and liver and kidney congestion.

2. Minimum Lethal Dose by Skin Absorption in Rabbits

No. of animals used	-	6 (3 male and 3 female)
Method of administration	-	By application to the clipped, intact skin then covered with plastic.
Dosage Levels	-	250, 400, 600, 800, 1000, 1500 mg/kg.

Results

Mortality with increasing dosage was found to be 0/1, 0/4, 0/1, 1/1, 1/1. 1/1. Animals died in 4 to 6 hours. The Minimum Lethal Dose by skin absorption was thus between 600 and 800 mg/kg. Symptoms included rapid onset of weakness followed by convulsions and coma. Gross findings at autopsy on animals which died was pulmonary hyperemia.

3. Skin Irritation in Rabbits

No. of animals used	-	3
Method of administration	-	By application to the clipped, intact skin, then covered with plastic for 24 hours. Washed with soap and water after 24 hours.
Dosage level	-	Not given.

BEST AVAILABLE COPY

Results

Scoring of skin irritation was that of Draize et al., Jour. of Pharm. and Exp. Therapeutics 52, 1944.

Scores at 24 and 72 hours after removal of the patches were 1.0 and 0.6 respectively. Both erythema and several foci developed. Light eschar formation was noted. Randoxel is classed as a severe skin irritant.

4. Eye Irritation in Rabbits

No. of animals used	-	3
Method of administration	-	Instillation into the conjunctival sac.
dosage level	-	0.1 ml

Results

Scoring of the eye irritation was that of Draize et al., op.cit. pp. 59, 64. Moderately severe edema and erythema as well as copious lacrimation were observed. Moderate dulling of the cornea and icteric conjunctiva were seen. Scores at 24 and 72 hours were 51.3 and 32.6 respectively. Slight conjunctival effects were still present at 7 days. Randoxel is classed as a moderately severe eye irritant.

5. Nasal Inhalation (rats)

No. of animals used	-	5 (males)
Method of administration	-	Inhalation of vapor concentration produced by bubbling air at 1 l/minute through 20 ml of Randoxel contained in a 250 ml gas washing bottle then passing the air successively through a five gallon bottle and a 70 liter exposure chamber.
Vapor concentration	-	Not determined.
Length of exposure	-	6 hours.

Results

No deaths occurred. Lacrimation was produced. Moderate ocular and nasal inflammation developed during the exposure period. No severe symptoms developed either during the exposure or the 6-day post-exposure observation period.

The Randoxel used in the foregoing studies is presumed to be the Randoxel liquid preparation.

BEST AVAILABLE COPY

③ 3

6. Study on subacute toxicity of 1,1,1-trichloro-2,2,2-trifluoroethane (TCE) in rats 187C 4810 ; 03/29/67

- Material used - A mixture of 3.1 lbs of Randox technical (p-chloro-2,2,4,4-tetrafluoroethylidene) and 0.3 lbs of trichlorobenzyl chloride.
- Brand - Randox T
- No. of animals - 24; 4 groups of 6 animals (3 males and 3 females per group)
- Method of administration - Admixture to the diet.
- Diet concentrations - 0, 20, 62 & 200 ppm.

Parameters studied were:

Mortality

Behavioral reactions

Body weights - Determined weekly

Food consumption - Determined weekly

Hematology, clinical blood chemistry and urinalysis were performed for all groups prior to the test; at 45 days for the control and high level animals; for all groups just prior to termination.

Hematology included hemoglobin, hematocrit, erythrocyte count and total and differential leucocyte counts.

Clinical blood chemistry included urea nitrogen, glucose, serum alkaline phosphatase, SGOT, SGPT.

Urinalysis included glucose, albumin, pH and microscopic elements.

Organ weights and organ to body weight ratios were obtained in each group for liver, kidney, spleen, heart, brain, gonad, adrenal and thyroid.

Pathology including microscopic examination of adrenal, thoracic aorta, bone marrow, brain (cerebrum, cerebellum, pons), caecum, colon, esophagus, eyes, gall bladder, gonad, heart, kidney, liver, lung, lymph nodes (cervical & mesenteric), nuzzle, optic nerve, pancreas, parathyroid, sciatic nerve, pituitary, prostate, submaxillary gland, skin, duodenum, jejunum, ileum, spleen, stomach (cardia, fundus, pylorus), trachea, thymus, thyroid, uterus, urinary bladder.

Results:

No significant abnormalities were found, although there is a suggestion at the 200 ppm level of a decrease in the rate of weight gain which was more evident in males than in females. Sixty two ppm is considered to

BEST AVAILABLE COPY

4

002481

PP No. 050 64

- 5 -

November 27, 1967

represent a "no effect" level. In terms of each of the ingredients this represents levels of approximately 70.7 ppm for the Roxon and 41.3 for the trichlorobenzyl chloride.

7. Two-Day Subacute Oral Toxicity Study in Rats IRT 84809 ; 04/20/67

Material used	-	A mixture of 3.1 lbs of Roxon, technical and 6.3 lbs of trichlorobenzyl chloride.
No. of animals	-	80; 4 groups of 20 animals (10M and 10F per group).
Method of Administration	-	Admixture to the diet.
Diet concentration	-	0, 20, 62 and 200 ppm.

Parameters studied were:

Mortality

Behavioral reactions

Body weights - determined weekly

Food consumption - determined weekly

Ecatology, urinalysis and clinical blood chemistry were performed at 30 days and just prior to termination.

Hematology included hemoglobin, hematocrit, erythrocyte count and total and differential leucocyte counts.

Urinalysis included glucose, albumin, pH, specific gravity, microscopic elements.

Clinical blood chemistry included urea nitrogen, serum alkaline phosphatase and isocetric dehydrogenase.

Organ, and organ to body weight ratios for liver, kidney, spleen, gonad, heart and brain.

Pathology including microscopic examination in surviving animals of the control and high level group of heart, trachea, lung, liver, pancreas, esophagus, stomach (cardia, fundus, pylorus), duodenum, jejunum, ileum, caecum, colon, spleen, lymph node, thymus, kidney, urinary bladder, testes, ovary, seminal vesicle, prostate, uterus, pituitary, adrenal, submaxillary, thyroid, parathyroid, intercostal muscle, bone marrow, peripheral nerve, spinal cord, brain (cerebrum, cerebellum, pons).

Results

No significant abnormalities were found in any of the parameters except the liver to body weight ratio. In this respect treated males showed statistically larger livers at all levels and treated females showed

BEST AVAILABLE COPY

5

slight changes at the two higher dose levels. Microscopic findings, however, were negative. The absence of any increase in isocitric dehydrogenase and serum alkaline phosphatase also suggests absence of liver effects. However, in a 3 week target organ study in rats there was considerable liver pathology at 1500 ppm. It would be desirable to obtain individual data for liver, body weight and food consumption if available.

Toxicological Data Trichlorophenyl Chloride

1. Acute Oral Toxicity in Rats

- No. of animals used - 22 (13 males & 10 females)
- Method of administration - Intubation
- Dosage levels - 2510, 2830, 3310, 3800 mg/kg

Results

Mortality with increasing dosage was found to be 1/5, 3/7, 3/3, 5/5. The oral LD₅₀ was determined as 3075 mg/kg with limits of 2800-3400 mg/kg. Survival time was overnight to five days, most deaths occurring overnight. Symptoms included salivation, diarrhea, lethargy and weakness. The animals were very irritable. Gross autopsy of those animals which died showed kidney discoloration and inflammation of the gastric mucosa.

2. Minimum Lethal Dose by Skin Absorption in Rabbits

- No. of animals used - 7 (5 male & 2 female)
- Method of administration - By application to the clipped intact skin and covered with plastic.
- Dosage levels - 1000, 2500, 3000, 4000, 5000 & 7500 mg/kg.

Results

Mortality with increasing dosage was found to be 0/1, 0/1, 0/1, 0/1, 0/1, 2/2. The Minimum Lethal Dose by Skin Absorption was thus between 5000 and 7500 mg/kg. Survival time was 2 and 3 days. Symptoms included loss of appetite, lethargy and weakness. Paralysis, if present, could not be distinguished from weakness. Gross findings at autopsy on the animals which died were pulmonary hyperemia and some evidence of liver dysfunction.

3. Skin Irritation in Rabbits

- No. of animals used - 3

BEST AVAILABLE COPY

64

Method of administration - By application to the clipped, intact skin, then covered with plastic for 24 hours. After 24 hours washed with soap and water.

Dosage level - Not given.

Results:

Scoring of the skin irritation was that of Draize *et al.*, *op.cit.* Scores at 4 and 43 hours after removal of the patches were 2.6 and 1.0 respectively. Both well defined erythema and slight edema developed. ICSC is classed as a moderate skin irritant.

4. Eye Irritation in Rabbits

No. of animals used - 3

Method of administration - Instillation into the conjunctival sac.

Dosage level - 0.1 ml.

Results:

Scoring of the eye irritation was that of Draize *et al.*, *op.cit.* In four hours edema sufficient to half close the eyelids as well as moderately severe erythema, copious lachrymation and moderate iritis was noted. These persisted through 24 hours. Some conjunctivitis remained in two animals at the seventh day. ICSC is classed as a moderately severe eye irritant.

5. Vapor Inhalation (rats)

No. of animals used - 6 (males)

Method of administration - Inhalation of vapor concentration produced by bubbling air at 14 l/minute through 100 ml of ICSC contained in a 250 ml gas washing bottle and then passing the air successively through a five gallon bottle and a 70 liter exposure chamber.

Nominal Concentration - Not determined.

Lengths of exposure - 6 hours.

Results:

No deaths occurred. Moderate nasal and ocular inflammation developed and cleared after several days. The animals did not exhibit normal activity or appetite until 3 - 4 days after exposure. No serious pulmonary complications were observed grossly. ICSC was considered to be moderately harmful to rats under these conditions.

BEST AVAILABLE COPY

7

November 27, 1967

002481

6. Chronic Toxicity Studies in Rats

Material used - Trichlorobenzyl chloride.
 No. of animals - 10; 3 groups of 70 animals (10 males and 10 females).
 Method of administration - additive to the diet.
 Diet concentrations - 0, 0.15, 0.30 per cent.

Parameters studied were:

Mortality

Behavioral reactions

Body weights - determined weekly

Food consumption - determined weekly

Hematology and urinalysis were performed initially and at termination.

Hematology included hemoglobin, hematocrit and total and differential leucocyte counts.

Urinalysis included reducing substances, albumin and microscopic elements.

Pathology including microscopic examination of heart, lung, liver, spleen, pancreas, esophagus, stomach, small intestine, large intestine, lymph nodes, adrenal, kidney, urinary bladder, thyroid, ovary and uterus, testes, brain, bone and bone marrow.

Results

Of the parameters studied effects were noted upon weight gain, food consumption and pathology. Weight gains were depressed in the male rats in both test groups and in the female rats fed the higher concentration of TCBC. Food consumption in the male test groups was somewhat lower than for the controls and was suggestive of dose correlation. No such pattern was seen for the females. There is a suggestion in the data that the body weight differences might be due to systemic toxic effects. Gross examination of the high-dose level animals revealed pale livers with distinctly granular textures. In the low-dose group degenerative liver changes were seen in 9 of the 10 males and in 5 of the 10 females. In the low-dose animals the livers were pale but were not granular. Microscopically, of the 19 organs and tissues examined, significant changes were found only in the liver. These included: (a) general derangement of hepatic chord formation; (b) irregularity of cell and nuclear size; (c) the occurrence of degenerative forms of hepatic cells characterized by acidophilic alterations of cytoplasm and pyknosis of nuclei; (d) regenerative forms of hepatic cells characterized by mitotic activity or by bi-nucleate forms; (e) hepatic cell edema. No associated inflammatory infiltrate was found. No evidence of significant fatty alteration was noted. Pigment deposition was not

BEST AVAILABLE COPY

7
8

present. These microscopic changes were found to be dose related being most severe in the 0.3% dosage group and less severe but consistently present in the 0.1% dosage group, especially in the male animals.

Toxicological data on *p*-chloro-*N*,*N*-dimethylacetamide were previously submitted in connection with Pesticide petition No. 710 617 and are summarized therein.

RESULTS:

1. Subacute oral toxicity studies in dogs with CDMA alone and in combination with TCBC did not produce any significant deviations in the parameters studied. In rats subacute feeding of CDMA alone produced no significant effects. However CDMA in combination with TCBC fed over a ninety-day period produced increased liver to body weight ratios as compared to the controls. For the males this occurred at all three feeding levels, for the females at the two higher levels. Microscopic and clinical biochemical findings, however, disclosed no abnormalities attributable to the compound. Definitive evidence of liver damage was obtained in rats fed TCBC at levels of 1500 and 3000 ppm over a three week period. The petitioner should provide liver weights, body weights and food consumption data for individual animals.

2. Acute oral toxicities (LD₅₀) obtained in rats were as follows:

- CDMA & TCBC (Ratio 3:1 to 6:3) - 1065 mg/kg
- TCBC - 1073 mg/kg
- CDMA (In corn oil) - 0.79 ml/kg

3. Minimal Lethal Dose by Skin Absorption in Rabbits were estimated to fall within limits indicated as follows:

- CDMA & TCBC (Ratio 3:1 to 6:3) - 600-900 mg/kg
- TCBC - 500-750 mg/kg
- CDMA (using a 48.8% active material) - 100-200 mg/kg

For the mixture above survival times were 4 to 9 hours. For the CDMA it was 4 hours to overnight. In the case of TCBC survival time was 2 to 3 days.

4. Topical application of Roadox-1 liquid preparation to rabbit skin and eye shows it to be a severe skin irritant and a moderately severe eye irritant. CDMA (tested as a 48.8% active material) produced severe skin irritation permanent blindness. TCBC was found to be a moderate skin irritant and moderately severe eye irritant.

5. Exposure for 6 hours to vapor concentrations produced from Roadox-1 liquid and TCBC produced no lethality and only minor effects in rats. Inhalation exposure to CDMA for 6 hours was lethal to all animals.

BEST AVAILABLE COPY

1
9

002481

FD No. 870 654

- 10 -

November 27, 1967

exposed within 24 hours. This latter result is not, however, comparable to the first two since the conditions of the exposures were quite different.

CONCLUSIONS:

The combination of *p*-chloro-2,4-diallyloacetamide and trichlorobenzyl chloride in the ratio of 1:2 is tolerated when fed subcutely at concentrations up to 62 ppm without significant signs of toxicity in dogs. For rats there is increased liver weight that requires further exploration. The petitioner is requested to provide body weight, liver weight, and food consumption data.

RECOMMENDATION:

Final evaluation of the safety of this material awaits evaluation of the requested data.

INITIALS: lumenthal

cc:

5-979

5-970 (Dr. Jacobson)

5-507

5-100

FD Nos. 870 654 & 770 617

REG: lumenthal 11/27/67

BEST AVAILABLE COPY

10

(10)