US ERA ARCHIVE DOCUMENT

U TED STATES ENVIRONMENTAL PR ECTION AGENCY

SUBJECT:

2.4.-D tolerances

DATE: December 23, 1975

FROM:

Toxicology Branch

TO:

Product Manager (Mountford)

Special Registration Section (Baker)

Petitions: 1/2E1293 5ppm on apricots IR-4 6E1678 5ppm on citrus (from/PHA) IR-4 1E1046 1ppm fish & shellfish Corps of Engineers 6H5104 o.1ppm in water TVA 3E1390 1ppm fish 3H5032 0.1ppm in water

Conclusion: CB permitting all tolerances can be toxicologically supported.

Review:

1. The toxicity of 2,4,-D has been previously reviewed, and furthermore 2,4,-D showed no data gaps when reviewed for reregistration. The NEL for 2-year rat feeding studies was shown to be 1250ppm and in a 3 generation rat reproduction study the NEL was 500ppm. 2,4,-D also showed no teratogenicity. Using a 100 fold safety factor the ADI for man would be 0.25mg/kg/day or 10ppm in his diet.

- 2. There is already a FAP of 0.1ppm in water established for the western part of the US (CFR 123.100).
- 3. There is already a tolerance established of 5ppm on citrus from preharvest use and a 5ppm tolerance on lemons from post harvest uses. The extension of the post-harvest use to all citrus (1678) thus will not change the actual exposure.

4. A tolerance of 5ppm on apricots is insignificant vis a vis the 5ppm tolerance on citrus.

Reto Engler, Ph.D Toxicology Branch

Registration Division

Initial: OE. Paynter
EPA Form 1320-6 (Rev. 6-72)

OEA 1919

Ef 19/23/75

<u>\</u>