

US EPA ARCHIVE DOCUMENT

5-29-91

030001

DATA EVALUATION REVIEW

- I. Study Type: Anaerobic Aquatic Metabolism Study
- II. Citation: Cohen, S.P. and V. V. Rama. 1990. Anaerobic Aquatic Metabolism of 2,4-Dichlorophenoxyacetic Acid. performed by Center of Hazardous Materials Research, Pgh., PA and submitted by 2,4-D Industry Task. MRID 415579-01.
- III. Reviewer:

Name: James A. Hetrick, Ph.D., Chemist
Title: Environmental Chemistry Review Section #1
Organization: EFGWB/EFED/OPP

James A. Hetrick

IV. Approved by:

6 Name: Paul J. Mastradone, Ph.D., Chief
Title: Environmental Chemistry Review Section #1
Organization: EFGWB/EFED/OPP

Paul J. Mastradone

V. Conclusions:

This study provides supplemental data on the metabolism of 2,4-dichlorophenoxyacetic acid (2,4-D) in anaerobic aquatic environments. The study cannot fulfill the anaerobic aquatic metabolism (162-3) data requirement for the following reason:

The material balance varied as a function of time. For example, poor material balances (70 to 51% of applied) were observed between the 70 and 365 sampling dates. The registrant believes poor material balances were caused by inefficient trapping of CO₂. EFGWB believes a poor material balance prevents validation of the experimental procedures.

Based on supplemental data, the half-life for 2,4-D in anaerobic (Eh=-220 mv) aquatic environments was 41 days (R²=0.91). After a 365 day incubation, [¹⁴C]-residues were distributed in the aqueous phase (1.23% of applied), sediment phase (9.00% of applied), and volatile phase (71.36% of applied). The [¹⁴C]-residue accounted for 51 to 97.5% of applied [¹⁴C]-2,4-D. The extractable soil/water residues were identified as 2,4-D, chlorophenol and 2-chlorophenol; and the volatile residue was tentatively identified as CO₂.

The reported data indicate 2,4-D appears to be moderately stable in anaerobic aquatic environments.

VI. Materials and Methods:

A subsample (1200 gm) of nonsterile, Louisiana rice paddy sediment (clay texture, 3.6% O.M., pH 7.3, CEC 28.9 meq 100⁻¹, B.D. 1.20 g cm⁻³) and water was placed into a 2 liter flask, which was connected to a series of sequential gas traps including



2011868

ethylene glycol and sulfuric acid. The sediment was then incubated under continuous, flow-through N_2 system (40-50 ml min⁻¹) at a temperature of $25 \pm 0.8^\circ C$ (Figure 5).

After a 138 day anaerobic incubation period, the sediment:water mixture was amended with radiolabeled 2,4-D solution (SA-59.7 μCi mg⁻¹, isotopic dilution factor 0.56) to produce a concentration of 4.861 μg g⁻¹. The incubation flask was then reconnected to existing gas traps; and the sediment/water system was incubated for 365 days.

Sediment/water samples (25 gm) were taken at 0, 1, 6, 13, 22, 35, 70, 85, 120, 160, 224, 281, 338, and 365 days post-treatment.

Analytical

Duplicate sediment/water samples were taken at each sampling time. Additionally, sediment/water samples were also frozen (by freezing at $-20^\circ C$) for future analyses. Prior to chemical analysis, each sample was centrifuged to separate the sediment and solution phases.

Each sediment sample was sequentially extracted with H_3PO_4 -anhydrous ethyl ether, H_3PO_4 -water, and 1N NaOH; followed by soil combustion-LSC. Prior to chemical analyses, the H_3PO_4 -ether extracts were evaporated to dryness and redissolved in D* D water.

The [^{14}C]-residues in sediment extracts were separated by HPLC with a C-18 Micro PaK Column and a linear gradient solvent system (0.1% trifluoroacetic acid and 0.1% in acetonitrile); and a UV detector. The HPLC analysis system had an level of quantification (LOQ) of 5 μg kg⁻¹. Additionally, soluble residues were also separated using TLC with a benzene:ethyl acetate:acetic acid 86:10:4 v/v/v solvent system. The separated residues were identified by co-chromatography with 2,4-D, 2,4-dichlorophenol (2,4-DC), phenoxyacetic acid, ortho-chlorophenoxyacetic acid, para-chlorophenoxyacetic acid, 1,4-dihydroxy-2-chlorophenol, 1, 2, 4-benzenetriol, 2-chlorophenol, and 4-chlorophenol.

The supernatant samples were acidified with 1% trifluoroacetic acid, filtered, and analyzed by HPLC and TLC. In addition, the total ^{14}C content in duplicate water and trapping solutions was measured using LSC.

I. Study Author's Results and/or Conclusions:

A. The applied 2,4-D was partitioned between aqueous, sediment, and volatile phases under anaerobic conditions; at time 0, the [^{14}C]-residues were distributed in the aqueous phase (66.3% of applied) and sediment phase (33.72% of applied); and at 365 days, the [^{14}C]-residues were distributed in the aqueous phase (1.23% of applied), sediment phase (9.00% of applied), and volatile phase (71.36% of applied) (Tables 1 and 2). These residues

2

accounted for 51 to 97.5% of applied [^{14}C]-2,4-D.

Note: The material balance varied as a function of time. For example, poor material balances (70 to 51% of applied) were observed between the 70 and 365 sampling dates. The registrant believes poor material balances can be attributed to inefficient trapping of CO_2 .

B. The half-life for 2,4-D in anaerobic ($\text{Eh}=-220$ mv) aquatic environments was estimated at 41 days ($R^2=0.9145$) (Figure 7).

C. The extractable soil/water residues were identified as 2,4-D (92.0% of applied at 0 days), 4-chlorophenol (21% of applied at 35 days) and 2-chlorophenol (<2% of applied at 13 days). The volatile residues were tentatively identified as CO_2 (71% of applied).

Reviewer Comments:

1. The material balance varied as a function of time. For example, poor material balances (70 to 51% of applied) were observed between the 70 and 365 sampling dates. The registrant believes poor material balances were caused by inefficient trapping of CO_2 . EFGWB believes a poor material balance prevents validation of the experimental procedures.
2. The sediment/water redox potentials were measured using a platinum electrode; the sediment/water mixtures had redox potentials of ≈ -220 mV ($\text{pe} = 3.71$). At such low redox potentials ($\text{pe} < 2$), PtS may precipitate on the Pt electrode and, therefore, measured redox potentials may only approximate ambient soil/water electron activities. EFGWB believes, however, the experiment was conducted under anaerobic conditions.

3

Page _____ is not included in this copy.

Pages 4 through 15 are not included in this copy.

The material not included contains the following type of information:

- Identity of product inert ingredients.
- Identity of product impurities.
- Description of the product manufacturing process.
- Description of quality control procedures.
- Identity of the source of product ingredients.
- Sales or other commercial/financial information.
- A draft product label.
- The product confidential statement of formula.
- Information about a pending registration action.
- FIFRA registration data.
- The document is a duplicate of page(s) _____.
- The document is not responsive to the request.
- Internal deliberative information.
- Attorney-Client work product.
- Claimed Confidential by submitter upon submission to the Agency.

The information not included is generally considered confidential by product registrants. If you have any questions, please contact the individual who prepared the response to your request.
