

US EPA ARCHIVE DOCUMENT

NAPHTHALENEACETIC ACID

Task 3: Environmental Fate Profile

Contract No. 68-01-5830

Final Report

February 27, 1981

SUBMITTED TO:

**Environmental Protection Agency
Arlington, Virginia 22202**

SUBMITTED BY:

**Enviro Control, Inc.
One Central Plaza
11300 Rockville Pike
Rockville, Maryland 20852**




This document contains commercial and/or financial information of Enviro Control, Inc. that is privileged or confidential and is therefore subject to the restrictions on disclosure outside the agency to which such document has been submitted, as provided in the Freedom of Information Act (PL 89-554, September 6, 1966, as amended).

NAPHTHALENEACETIC ACID

TASK 3

Naphthaleneacetic acid, NAA 800,
Fruitone N, Rootone, Transplantone,
Tre-Hold



1-Naphthaleneacetic acid

Environmental Fate Profile

Available data are insufficient to fully assess the environmental fate of naphthaleneacetic acid (NAA).

Aspergillus niger was implicated as metabolizing NAA to 5-hydroxy NAA. Other possible products are the 4- and 6-hydroxy NAA isomers, as well as other phenolic or naphtholic compounds. Hydroxylation thus appears to be a major route of degradation of NAA.

A coliform bacterium and two animal pathogens (Staphylococcus aureus and human type tubercle bacilli) are inhibited by NAA at 50-300 ppm, whereas avian type tubercle bacilli are not inhibited by NAA at 50 ppm. A yeast, Saccharomyces cerevisiae, is inhibited 13-58% by NAA at 100-500 ppm. In addition, large-cell mutants result from the induction of polyploidy. Two mutants of Neurospora crassa are inhibited 12-68% by NAA at 10-50 ppm, but a wild type control strain was not studied. An alga, Chlorella pyrenoidosa, is inhibited 40-80% by NAA at 46-372 ppm. Thus, a wide range of microorganisms is inhibited by NAA at 10-500 ppm which are concentrations far in excess of the highest expected level in the soil (0.08 ppm). Most studies dealt with mutants, animal pathogens, or soil microorganisms of questionable importance. None of the studies reported the source and purity of the NAA used.

In summary, although very little data are available to form a profile of the environmental fate of NAA, 5-hydroxy NAA is a suspected fungal metabolite of NAA. At very high concentrations, NAA is capable of inhibiting bacteria, fungi, and algae and also of inducing polyploid mutants. The available data suggest that currently approved NAA application rates will not have an effect on soil microorganisms.

Data Gaps

The submitted studies are not sufficient to assess the fate of NAA in the environment. The following data gaps have been identified.

Type of Data	Guideline Section
Hydrolysis Naphthaleneacetamide NAA NAA-ammonium salt NAA-ethyl ester NAA-potassium salt NAA-sodium salt	163.62-7(b)
Activated sludge metabolism Naphthaleneacetamide NAA NAA-ammonium salt NAA-ethyl ester NAA-potassium salt NAA-sodium salt	163.62-8(g)

Photolysis	163.62-7(c)
Soil metabolism	163.62-8(b,c)
Microbiological metabolism	
• Effects of microbes on NAA	163.62-8(f)(2)
• Effects of NAA on microbes	163.62-8(f)(3)
Leaching	163.62-9(b)
Volatility	163.62-9(c)
NAA	
Adsorption/desorption	163.62-9(d)
Terrestrial field dissipation	
• Tree fruit and nut crop uses	163.62-10(b)(2)

Naphthaleneacetamide
8.4% ai wettable powder
0.176 lb/gal soluble concentrate/liquid

NAA
0.2% ai dust
3.5% ai wettable powder
0.106 lb/gal emulsifiable concentrate
0.28 lb/gal soluble concentrate/liquid

NAA-ammonium salt
1.76 lb/gal soluble concentrate/liquid

NAA-potassium salt
1.76 lb/gal soluble concentrate/liquid

NAA-sodium salt
7.11% ai wettable powder
98% ai crystalline
3.5% ai soluble concentrate/solid

- Domestic outdoor, parks, ornamental and turf uses 163.62-10(b)(4)

NAA

0.12% ai soluble concentrate/liquid

NAA-ammonium salt

1.76 lb/gal soluble concentrate/liquid

NAA-ethyl ester

1% ai ready to use

1% ai pressurized liquid

NAA-sodium salt

7.11% ai wettable powder

Fish accumulation

163.62-11(d)

Label Restrictions

Current label restrictions warn against the use of NAA (acid) in combination with insecticides or fungicides. NAA-potassium salt is not to be used in the greenhouse.

References

Clifford, D.R., and D. Woodcock. 1968. Fungal detoxication-IX. Metabolism of 1-naphthaleneacetic acid by Aspergillus niger Van Tiegh. *Phytochemistry* 7(9):1499-1502. (MRID 05011538)

Doi, S., T. Takahashi, and N. Yanagishima. 1973. Auxin-induced large cell mutants in Saccharomyces cerevisiae. I. Induction, and biochemical and genetic characters. *Jap. J. Genetics*. A translation of: *Idengaku Zasshi*. 48(3):185-195. (MRID 05008826)

Gramlich, J.V., and R.E. Frans. 1964. Kinetics of Chlorella inhibition by herbicides. *Weeds*. 12(3):184-189. (MRID 05009027)

Klein, D.T. 1962. Effect of growth regulators on mutants of Neurospora crassa. *Physiologia Plantarum*. 15:239-245. (MRID 05010966)

Loveless, L.E., E. Spoerl, and T.H. Weisman. 1954. A survey of effects of chemicals on division and growth of yeast and Escherichia coli. *J. Bacteriol.* 68:637-644. (MRID 05005277)

Ukita, T., O. Tamemasa, and H. Motomatsu. 1951. Antibacterial action of fatty acids. VIII. Syntheses of fatty acids with naphthyl, naphthomethyl and benzyl groups in alpha-position and their antibacterial action. *Yakugaku Zasshi. J. Pharm. Soc. Jap.* 71(4):289-297. (MRID 05010797)