

US EPA ARCHIVE DOCUMENT

DATA EVALUATION RECORD


TRICHLORFON

METABOLISM

CITATION: Abo-Khatwa N, Hollingworth RM. 1974. Pesticidal chemicals affecting some energy-linked functions of rat liver mitochondria in vitro. Bull. Environ. Contam. Toxicol. 12:446-454.


REVIEWED BY:

Nicolas P. Hajjar, Ph.D.  
Senior Scientist  
Dynamac Corporation  
11140 Rockville Pike  
Rockville, MD 20852  
301-468-2500

Signature: 

Date: July 25, 1983


John R. Strange, Ph.D.  
Department Director  
Dynamac Corporation  
11140 Rockville Pike  
Rockville, MD 20852  
301-468-2500

Signature: 

Date: 25 July, 1983

APPROVED BY:

Irving Mauer, Ph.D.  
EPA Scientist

Signature: 

Date: 07-25-83

## DATA EVALUATION RECORD

STUDY TYPE: Metabolism, in-vitro.

CITATION: Abo-Khatwa N, Hollingworth RM. 1974. Pesticidal chemicals affecting some energy-linked functions of rat liver mitochondria in vitro. Bull. Environ. Contam. Toxicol. 12:446-454.

ACCESSION NUMBER: Not available.

MRID NUMBER: 05004940.

LABORATORY: Department of Entomology, Purdue University, West Lafayette, Indiana.

TEST MATERIAL: Trichlorfon (>95 percent purity).

### PROTOCOL:

1. Trichlorfon (>95 percent purity) was used in this assay along with 46 other pesticides.
2. Rat liver mitochondria were isolated according to the method of Katyare et al. (Arch. Biochem. Biophys. 144:209, 1971), and oxygen uptake was measured polarographically at 30°C using an assay previously described by Abo-Khatwa and Hollingworth (Pestic. Biochem. Physiol. ["in press (1973)"]).
3. The incubation mixture consisted of succinate as the substrate (concentration unspecified) and 1.2 mg of mitochondrial protein/ml of reaction mixture at a final volume of 2.5 ml. The test chemical was dissolved in ethanol and 5  $\mu$ l added to the reaction mixture at a final concentration of  $10^{-4}$  or  $10^{-3}$ M.
4. Four energy-linked functions of the mitochondria were measured:
  - (1) ADP/O ratio (efficiency of phosphorylation).
  - (2) Rate of O<sub>2</sub> uptake in the absence of a phosphate acceptor.
  - (3) Rate of O<sub>2</sub> uptake in the presence of phosphate acceptor.
  - (4) Respiratory control index (ratio of O<sub>2</sub> uptake before and after ADP is exhausted as an acceptor).

## RESULTS:

Trichlorfon was inactive in all four of the energy-linked functions at concentrations up to  $10^{-3}M$ . The compound was therefore not considered to be an uncoupler of oxidative phosphorylation.

## CONCLUSIONS:

Under the experimental conditions used trichlorfon did not uncouple oxidative phosphorylation in rat liver mitochondria. Although, it is not clear if positive controls were used in this study, several compounds (e.g., 2,5 dichloro-4-iodophenol, chlordimeform) were found to be potent uncouplers ( $10^{-6}M$ ) of oxidative phosphorylation.

CORE CLASSIFICATION: Supplementary data.

The study provides information indicating that the test material is not an uncoupler of oxidative phosphorylation in vitro.