


US EPA ARCHIVE DOCUMENT

EPA #	Date	Product Name	Use Classification	Toxicity Category
5815-GL	5/18/76	W 2910 xxx	General	III

RECOMMENDATION:

Label is satisfactory
OK to register

60% Dacthal



003294

INERT INGREDIENT INFORMATION IS NOT INCLUDED

		TECH	TECH	FORMULATION	USE DILUTION	DATA ACCEPT ABLE
acute Oral (Rat)	LD50	>3150 mg		>10,250 mg/kg		

Toxic signs: none

Comments:

acute Dermal (Rabbit)	LD50	>10,000 mg/kg				
-----------------------	------	---------------	--	--	--	--

Toxic signs:

Comments:

acute Inhalation (Rat)	LC50	>4.8 mg/L		>19.4 mg/L		
------------------------	------	-----------	--	------------	--	--

Toxic signs:

Comments:

Primary Eye Irritation (Rabbit)		mild		1.0/1.0		
---------------------------------	--	------	--	---------	--	--

Comments:

no eye irritation

Primary Skin Irritation (Rabbit)		slight irritation		0.6/8.0		
----------------------------------	--	-------------------	--	---------	--	--

Comments:

Other Studies:

Guinea Pig Sens - not a sensitizer

JF 5/18/76

003294

FRONT LABEL
Reduced 1/2 of ACTUAL SIZE

XXX

TRIPLE X PRODUCTS

HOE NO MORE WITH

GARDEN

WEED

PREVENTER

ACTIVE INGREDIENTS

Dimethyl tetrachloroterephthalate * 5.00%

INERT INGREDIENTS 95.00%

TOTAL 100.00%

*Use as a preemergence herbicide patented under U.S. Patent No. 2,923,634

E.P.A. Reg. No.
E.P.A. Est. 5815-OH-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

COVERAGE: 2500 sq. ft.
NET WT. 10 lb.

AMERICAN CHEMICAL CORPORATION

Rev

GARDEN WEED PREVENTER
Coverage: 2500 sq. ft.
Net Wt. 10 lb.

303294 BACK LABEL
Reduced to 1/2 of ACTUAL SIZE

**PRECAUTIONARY STATEMENTS
CAUTION**

- KEEP OUT OF REACH OF CHILDREN
- HARMFUL IF SWALLOWED
- AVOID CONTACT WITH EYES AND SKIN. IN CASE OF CONTACT, FLUSH EYES WITH PLENTY OF WATER. CALL A PHYSICIAN.
- WASH AFTER HANDLING OR USING
- DO NOT STORE NEAR FEED AND FOODSTUFF, SEED, FERTILIZERS OR OTHER PESTICIDES IN ORDER TO PREVENT CONTAMINATION
- DO NOT GRAZE ON TREATED AREAS OR FEED PLANT REFUSE TO LIVESTOCK

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

STORAGE AND DISPOSAL

- STORAGE:** DO NOT STORE NEAR FEED AND FOODSTUFFS, SEEDS, FERTILIZERS, OR OTHER PESTICIDES IN ORDER TO PREVENT CONTAMINATION. STORE IN A DRY PLACE.
- DISPOSAL:** DO NOT USE EMPTY CONTAINER. WRAP CONTAINER AND PUT IN TRASH COLLECTION.

THIS PRODUCT IS A SELECTIVE PREEMERGENCE HERBICIDE FOR THE CONTROL OF MANY BROADLEAFED WEEDS AND MOST ANNUAL GRASSES ON MINERAL SOILS FOR USE AROUND VEGETABLES, STRAWBERRIES, HOME FLOWER GARDENS AND NURSERY STOCK.

ANY APPLICATION SHOULD BE MADE PRIOR TO WEED SEED GERMINATION. RECOMMENDED RATE: APPLY AT A RATE OF 1 POUND PER 250 SQUARE FEET OR 1 POUND PER 250 SQUARE FEET. WEED CONTROL UP TO THREE MONTHS OR MORE CAN BE EXPECTED FOLLOWING PROPER APPLICATION. APPLICATION SHOULD BE MADE TO A UNIFORM WEED FREE SOIL WITHIN TWO TO THREE DAYS FOLLOWING CULTIVATION TO PREVENT SUBSURFACE GERMINATION WEEDS FROM BECOMING ESTABLISHED BEFORE GARDEN WEED PREVENTER IS ACTIVATED.

FOR BEST RESULTS - SPRINKLER IRRIGATION SHOULD BE USED WHEN THERE IS NO RAIN WITHIN THREE DAYS AFTER APPLICATION.

CREPS	WHEN TO APPLY	REMARKS
Broccoli, Brussels sprouts, Cauliflower, Cabbage, Garlic, Collards, Kale, Mustard Greens, and Onions	At seeding or transplanting	Apply uniformly to the soil at the rate of 1 pound per 250 square feet. If weeds have emerged, soil should be clean cultivated.
Snap beans, Southern peas (Black-eyed)	At seeding	Apply uniformly to the soil at the rate of 1 pound per 250 square feet at time of planting. Soil should be clean cultivated or weeded prior to application.
Potatoes (whole or cut pieces)	At planting or drag off	Apply uniformly to the soil at the rate of 1 pound per 250 square feet at planting or following planting. If the top of the buds are to be dragged off, Garden Weed Preventer should be applied after drag off. Soil should be weeded prior to application.
Sweet Potatoes, yam	At transplanting and anytime up to 2 weeks after transplanting	Apply uniformly to soil at the rate of 1 pound per 250 square feet at transplanting time. Soil should be weeded prior to application.
Tomatoes, Eggplant and Peppers	4 to 6 weeks after transplanting or on direct seeded plants at 4 to 6 inches in height	Uniform application to the soil should be confined to a period of 4 to 6 weeks after transplanting. Plants should be well established and growing conditions favorable for good plant growth. Do not apply to seeded plants until plants are 4 to 6 inches in height. Soil should be weeded prior to application.
Strawberries - New Plantings	At transplanting	Apply uniformly to the soil at the rate of 1 pound per 250 square feet at transplanting time. Soil should be weeded prior to application.
Strawberries - Established plantings	Fall and early spring	Apply uniformly at the rate of 1 pound per 250 square feet. DO NOT apply from first bloom through harvest. Application may be made directly over the plants without injury.
Nursery stock, Ornamentals and Flower Gardens	Early Spring	Apply uniformly to soil at the rate of 1 pound per 250 square feet. Soil should be weeded prior to application. Delayed or secondary applications can be made at any time provided beds have been cultivated and are free of weeds.

GARDEN WEED PREVENTER CAN BE USED WITH THE FOLLOWING

Abutilon	Ageratum	Alyssum	Andromeda, Jap	Artemisia
Achillea	Aster	Azalea	Aspidistra	Asplenium
Bellflower	Birch	Bleedingheart	Bluetop	Begonia
Begonia	Camellia	Candytuft	Chenopod	Chrysanthemum
Cinquefoil	Calceol	Columbine	Conflowers, Purple	Cornus
Conical bell	Coleus	Coronilla	Cottonwood	Crocus
Cubeba	Cosmos	Dahlia	Delphinium	Deutzia
Dogwood	Cypress	Daisy	Echinacea	Evening Primrose
Feverfew	Fir	Forget-me-not	Eranthis	Fuchsia
Forsythia	Gaillardia	Geranium	Galium	Gardenia
Goldenroft	Gum	Hemlock	Hebe	Holly
Hydrangea	Honey-suckle	Iris	Ivy, Boston	Ivy, Boston
Ivy, English	Ivy, Wilson	Juniper	Lantana	Lavender, candle
Lavender cotton	Lilac	Lily	Laurus	Loosestrife
Magnolia	Maple	Marguerite, golden	Marigold	Mourning Bride
Mammillaria	Mountain rose	Marshall-Thyme	Mountain Laurel	Pachysandra
Marshallia	Oak	Ornithogalum	Pachistima	Polemoni
Peony	Patunia	Pittosporum	Podocarpus	Primula
Pepper	Pink	Priest, variegated	Rhodod	Rhododendron
Rose	Sage, scarlet	Snapdragon	Soldierwort, Va.	Spiraea
Sauco	Shinerip	Strawflower	Sundrop	Sunflower
Sweet pea	Sycamore	Tree peony	Tulip tree	Viburnum
Violet	Walnut	Wiggle	Willow	Wormwood
Yarrow, farnesid	Yew	Zinnia		

DO NOT USE ON THE FOLLOWING PLANTS

Bugleweed	Butter pink	Camellia	Camellia
Mesembryanthemum	Fanny	Phlox	Sweet willow
		Comander	Telandra

Manufactured by:
Agro, Inc.
Grand Rapids, Ohio 43322

Vertical text on the left edge of the page.