

US EPA ARCHIVE DOCUMENT

7-4-79

BB-1889
TR-1889

Government of Canada
Gouvernement du Canada

MEMORANDUM

NOTE DE SERVICE

001484

Mr. D. J. Clegg
Head, Pesticides Section.

Dr. H. M. Cunningham

464
FOLPET
File

| |
|---------------------------------------|
| SECURITY CLASSIFICATION - DE SECURITE |
| OLD FILE - REFERENCE |
| YOUR FILE - REFERENCE |
| DATE July 4, 1979 |

Validation of Rat Teratogenic Study with Phaltan - IBT No. P-5397.

Overall Comment:

The audit and validation of this report indicate that the final report is not supported by the raw data. Of prime significance was a higher than actually found incidence of resorptions reported for the controls and absence of raw data on fetal skeletal and internal development.

BEST AVAILABLE COPY

1-75

Rat Teratogenic Study with Phalitan - IBT No. P-5397.

001484

A. Audit:

1. Report No. B-5397 dated October 11, 1967.
2. Date of study: Started on 21-5-67 and terminated on 27-7-67.
3. Protocol: No protocol available but procedure indicates that groups of 5-10 pregnant females (17 controls) received levels of 100 and 500 mg/kg of Phalitan suspended in corn oil (total dose = 0.1% of body weight) according to one of two dosing schedules - A = 10 doses from day 6 through 15 of gestation, or B = 3 doses from day 8 through 10. Observations were made on body weight, mortality, reactions and pathologic examination on all females sacrificed on 20th day of gestation. Number of implantation sites, resorption sites, corpora lutea, numbers of fetuses present, their weight and abnormalities in external and internal development and skeletal development were noted. Positive controls received trypan blue at 50 and 25 mg/kg.
4. Test Material: No evidence available to indicate that it was received at IBT.
5. Animal Suitability: Charles River strain albino rats obtained from Charles River Breeding Laboratory Ltd.
6. Raw Data: Initial weights of many rats missing at the 6th day of gestation when treatment started. Data absent on weighing out test material and very few data on mortality and reactions. No raw data to indicate that pathological procedures were actually conducted on gross pathology, internal development and on skeletal development.

B. Validation of Evaluation:

BEST AVAILABLE COPY

1. Dates: Dosages at the higher level (500 mg/kg) and a few rats at the lower level (100 mg/kg) appear to have been done about a month before the rest of the study and controls. Handwritten dates on summary dosage sheets indicate that the experiment started and finished on the approximate dates indicated above. None of the data sheets were signed.
2. Protocol: The protocol was not followed closely. At least three levels of trypan blue (12.5 mg/kg, 25.0 mg/kg, and 50 mg/kg) were used instead of only 50 mg/kg. The different groups of animals were dosed at different

2

001484

times and many observations were not made.

3. Test Material
& Quality
Control:

No shipping invoices or other data to show that the test material was sent to IBT. No data available on mixing the test material with vehicle for dosing although calculations showing the total amount of test material to be administered were available.

4. Personnel:

Report prepared by: Gerald Kennedy, B.S.
Staff Assistant

Report approved by:

Otis E. Fancher, Ph.D.
Director.

J. C. Calandra, M.D., Ph.D.
President

Technicians: No signatures on any of data.

C. Execution of the study:

a. Body weight:

Raw data were missing on the body weights on the first day of dosing on many of the rats but those that were available for final body weights agreed with the final report.

b. Dosing:

The use of two dosing schedules (10 doses with the lower level and 3 with the higher level) is not good scientific practice.

c. Mortality and
Reactions:

There was very little raw data on reactions. One death was reported at the high level (500 mg/kg) of Phal~~tan~~, supposedly after 3 doses but the raw data indicated that the dosing regime was complete and the rat died 1 day before the others in the same group were sacrificed.

d. Teratogenic and
Reproduction
Studies:

Table II (see appendix) has been corrected according to the raw data. It may be observed that the number of resorptions and abnormalities have been increased in the controls and changed somewhat in the positive controls. There did not appear to be any changes in the test animals.

e. Fetal Skeletal
Development:

No raw data available.

001484

f. Fetal Internal Development:

No raw data available.

g. Progeny Body Weight:

Only the average weights of the male and female fetuses in each litter were available in the raw data. The auditor, however, has taken the available data on fetal weights and calculated that statistically significant ~~significant~~ differences existed between the control and test male and female pups ($P < 0.05$) at the 100 mg/kg level, and ($P < 0.01$) at the 500 mg/kg level)

Overall Comment:

The audit and validation of this report indicate that the final report is not supported by the raw data. Of prime significance was a higher than actually found incidence of resorptions reported for the controls and absence of raw data on fetal skeletal and internal development.

H. M. Cunningham

H. M. Cunningham

D. J. Clegg

David J. Clegg

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED

Teratogenic and Reproduction Effects

Albino Rats

Summary of Results

| Group | Pregnant Females Examined | Corpora Lutea | | Implantation Sites | | Resorption Sites | | Fetuses | | Abnormal Fetuses | Par Co. |
|-------|---------------------------|---------------|---------------|--------------------|---------------|------------------|---------------|----------|---------------|------------------|------------|
| | | Total | Mean / Female | Total | Mean / Female | Total | Mean / Female | Total | Mean / Female | | |
| I | (10) 7 | (135) 96 | (13.5) 3.7 | (126) 92 | (12.6) 3.1 | (3) 0 | (0.3) 0 | (123) 92 | (12.3) 3.1 | (1) 0 | (0.8) 0 |
| II | (7) 7 | (84) 102 | (12.0) 14.7 | (80) 92 | (11.4) 13.1 | (2) 3 | (0.3) 4.2 | (78) 82 | (11.1) 11.7 | 0 | 0.0 |
| III | 10 | (129) 138 | (12.9) 13.7 | (113) 117 | (11.0) 11.8 | (5) 4 | (5.4) 5.7 | (64) 67 | (6.4) 5.7 | (17) 9 | (26.7) 5.7 |
| IV | 6 | 93 | 15.5 | 79 | 13.2 | 2 | 0.3 | 77 | 12.8 | 0 | 0.0 |
| V | 7 | 91 | 13.0 | 86 | 12.3 | 4 | 0.6 | 82 | 11.7 | 0 | 0.0 |
| VI | 5 | 49 | 9.8 | 49 | 9.8 | 4 | 0.8 | 45 | 9.0 | 0 | 0.0 |
| VII | 5 | 59 | 11.8 | 50 | 11.6 | 10 | 2.0 | 48 | 9.6 | 0 | 0.0 |
| VIII | 5 | 63 | 12.6 | 63 | 12.6 | 4 | 0.8 | 59 | 11.8 | 3 | 5.1 |
| IX | 5 | 64 | 12.8 | 60 | 12.0 | 0 | 0.0 | 60 | 12.0 | 2 | 3.3 |
| X | 9 | 132 | 14.7 | 121 | 13.4 | 4 | 0.4 | 117 | 13.0 | 0 | 0.0 |
| XI | 5 | 66 | 13.2 | 64 | 12.8 | 1 | 0.2 | 63 | 12.6 | 0 | 0.0 |
| XII | 10 | 134 | 13.4 | 124 | 12.4 | 4 | 0.4 | 120 | 12.0 | 0 | 0.0 |
| XIII | 40 | 51 | 12.7 | 51 | 12.7 | 2 | 0.5 | 49 | 12.2 | 1 | 2.1 |

One female died following the three doses. VO! 1 day since delivery was completed.

100140