

US EPA ARCHIVE DOCUMENT

# DATA REVIEW FOR ACUTE INHALATION TOXICITY TESTING (870.1300)

Product Manager: 22  
MRID No.: 45049105

Reviewer: Eugenia McAndrew  
Study Completion Date: January 20, 2000  
Study No.: 8138

Testing Facility: Product Safety Labs  
Author: George E. Moore, B.S.

Quality Assurance (40 CFR §160.12): Included

Test Material: GX-612/Equus DF; Lot #427-37; 85.94% Chlorothalonil; tan powder ground in a ball mill for 24 hours

Species: Rat; albino, Sprague-Dawley derived  
Age: Young adult  
Weight (fasted): Males: 204-242 g; Females: 181-215 g  
Source: Ace Animals, Inc., Boyertown, PA

## Conclusion:

1. **LC<sub>50</sub> (mg/L):**  
Males: > 0.056 mg/L  
Females: > 0.056 mg/L  
Combined: > 0.056 mg/L
2. **The estimated LC<sub>50</sub> is** > 0.056 mg/L
3. **Tox. Category: II**      **Classification: Acceptable**

Procedure (Deviations from 870.1300): None

Exposure Concentration mg/L (Gravimetrically Determined)	Number of Deaths/Number Tested		
	Males	Females	Combined
0.056	0/5	1/5	1/10
0.52	5/5	5/5	10/10
2.02	5/5	5/5	10/10

**Clinical Observations:** One female died on day 1 at the 0.056 mg/L exposure. Two males died during the 0.52 mg/L exposure; the remaining eight died within three days of exposure. Four animals died during the 2.02 mg/L exposure; the remaining six died within 2 hours of removal from the chamber. The following in-chamber observations were noted at all three exposure concentrations: ocular and nasal discharge, irregular respiration, dyspnea, hunched posture, and hypoactivity. All rats exhibited similar clinical signs upon removal from the chambers. At 0.056 mg/L, rales, piloerection, and reduced fecal volume were also observed but all surviving rats recovered by day 3 and gained weight. At 0.52 mg/L, one female was prone and had a reduced fecal volume. At 2.02 mg/L, the rats developed facial and ventral staining, rales and/or prone posture post-exposure.

1/2

**Necropsy Findings:** All rats that died during the study revealed discoloration of the lungs and edema of the lungs. At 0.056 mg/L, discoloration of the liver and rigor mortis were also present. No gross abnormalities were noted in the surviving rats. At 0.52 mg/L, discoloration of the liver and gastrointestinal tract, excess mucous in the trachea, gaseous distention of the GI tract and rigor mortis were also observed. At 2.02 mg/L, discoloration of the intestines and excess mucous in the trachea were observed.

Chamber Atmosphere		
Gravimetric conc. mg/L	MMAD $\mu\text{m}$	GSD
0.056	2.6	1.73
0.52	1.76	1.75
2.02	3.2	1.74

Chamber Environment <sup>a</sup>	
Chamber Volume	100 L
Airflow	50.6 LPM
Temperature	21-23°C
Relative Humidity	46-62%

<sup>a</sup>Whole body