

US EPA ARCHIVE DOCUMENT



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEB 16 1988

006597

OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

MEMORANDUM

SUBJECT: Review of a Two-Generation Reproduction Study of
Technical Bladex (Cyanazine) in Rats (Sprague-Dawley)

Caswell No.: 188C
TOX Proj. No. 8-0159

FROM: Sidney Stolzenberg, Ph.D.
Review Section V, Toxicology Branch
Hazard Evaluation Division (TS-769C)

S. Stolzenberg
2/12/88

TO: Robert J. Taylor, PM 25
Fungicide-Herbicide Branch
Registration Division (TS-767C)

and

Joan Dizikes, PM 64
Special Review Branch
Registration Division (TS-767C)

THRU: Quang Q. Bui, Ph.D., D.A.B.T.
Acting Head, Review Section V
Toxicology Branch
Hazard Evaluation Division (TS-769C)

Quang Q. Bui epr/88
W. J. Taylor
2/14/88

and

Theodore M. Farber, Ph.D., D.A.B.T.
Chief, Toxicology Branch
Hazard Evaluation Division (TS-769C)

Registrant: Agricultural Products Department
E.I. du Pont de Nemours & Company
Wilmington, DE

Action Requested:

Review the two-generation rat reproduction study with

1 *J. Taylor*

cyanazine. The data are of particular concern because of developmental toxicity in tests with this compound has been previously observed in both rats and rabbits.

Background Information:

Anophthalmia and microphthalmia were observed at 25 mg/kg/day and at 75 mg/kg/day in 2 different studies with Fischer 344 rats and at 2 mg/kg/day in a study with New Zealand rabbits. Other frank indications of developmental toxicity were seen in both species, including dilated brain ventricles, cleft palate and diaphragm abnormalities in rats, domed cranium, dilated brain ventricles, thoracoschisis, alterations in skeletal ossification sites, decreased litter size, and increased postimplantation loss in rabbits. A NOEL for developmental toxicity was set at 5mg/kg/day in the rat and at 1 mg/kg/day in the rabbit (See reports by Q-Q. Bui of July 14 and August 14, 1987).

Conclusions and Recommendations:

In the present two-generation rat reproduction study with Sprague-Dawley rats, the doses of technical Bladex, admixed in the diet, were 0, 25, 75, 150, and 250 ppm. Based on food intake, the dosages in mg/kg/day throughout the study were generally similar for nonpregnant and pregnant F₀ and F₁ parents. During these stages, they averaged 0, 1.8, 5.3, 11.1, and 18.5 mg/kg/day, respectively. During lactation, these doses, based on food intake, were about twice as high and came to 0, 3.8, 11.2, 23.0, and 37.1 mg/kg/day, respectively, averaged for all four of the F₁ and F₂ lactational periods in the study.

The LEL for reproductive toxicity is considered to be 11.2 mg/kg/day, based on pup viability and decreased mean pup body weight during lactation of dams on a diet with 75 ppm. The NOEL found was 3.8 mg/kg/day (25 ppm).

Systemic toxicity, based on decreased body weight of F₀ and F₁ adults at various time periods and final body weight of males and females of F₁ adults prior to necropsy, was seen at 1.8 mg/kg/day (25 ppm). Dose-related increased toxicity based on decreased body weight and occasionally decreased food intake at all intervals during the course of this study was observed. Therefore, the LEL for systemic toxicity to the F₀ and F₁ adults is \leq 1.8 mg/kg/day (LDT).

There was a 35% decrease in paired females that delivered a litter in the 150 ppm treated groups for both the F_{1a} and F_{1b} generations. A similar effect was not seen in the F_{2a} and F_{2b} generations nor was such an effect seen at the 250 ppm dose. The lower fertility in the 150 ppm group was considered a "random

occurrence" by the applicant but since it occurred in two generations, we should not completely ignore the possibility of an effect.

In the tables which present the Male Mating Index and Female Mating Index, data for "both mating periods", i.e. F_{1a} with F_{1b} and F_{2a} with F_{2b} were combined. In its present form, these data are obviously higher than they would be if they were calculated separately for each mating and not combined for both matings. It would also give us a better perspective on the contribution of both of these parameters to the reproduction performances in each generation. The applicant should be requested to recalculate the Male and Female Mating Indices separately for each mating and not combined for two mating periods.

There is a question of reliability of the data on "Females with Evidence of Mating". Vaginal sperm or copulation plugs were not detected for a number of animals that became pregnant. It also appears likely that there were animals considered to have mated but actually did not, particularly if evidence for mating was based on copulation plugs under the cages instead of vaginal sperm. These data and other data derived from them should be considered unreliable.

In Tables 117 and 118 (volume 2 of the report) and Tables 152 and 153 (vol 4), the footnotes indicate "does not include pups found missing or cannibalized." In the F_{1a} generation, dams 23,703 and 23,748 lost all of their 8 pups between day 7-14 of lactation but only 2 or 3 of them are listed in the necropsy data which indicates the rest of the litter could be considered as missing. Pups that are missing may be due to experimental error or loss in handling. The applicant should be requested to account for all missing pups to the extent that this is possible (See page 12 of this DER).

Core Classification: Supplementary

This may be upgraded to Minimum if the following are submitted and considered satisfactory.

- 1). Male Mating Index and Female Mating Index should be recalculated separately for each of the 4 matings.
- 2). All missing pups not included in the pathology data should be accounted for.

Primary Reviewer: Sidney Stolzenberg. Ph.D.
Review Section V, Toxicology Branch
Hazard Evaluation Division (TS-769C)
Secondary Reviewer: Quang Q. Bui. Ph.D.
Review Section V, Toxicology Branch
Hazard Evaluation Division (TS-769C)

006597

DATA EVALUATION REPORT

Study Type: Reproduction: Two-Generation
Species: Rat (Sprague-Dawley)
Guidelines: 83-4

Study Title: Two-Generation Reproduction Study of Technical
Bladex Herbicide (SD 15418) in Rats

EPA ID Nos.: 325-475, EPA Accession No. 403600-01, EPA Record
No. 206097, Caswell No. 188C, Project No. 8-0159

Sponsor: Agricultural Products Department
E.I. du Pont de Nemours & Company
Wilmington, DE

Testing Laboratory: WIL Research Laboratories, Inc.
Ashland, OH 44805-9281

Compound Submitted By: Shell Development Company
Houston, TX

Study Nos.: SRO 15-87
WIL 93001

Study Period: August 29, 1985 to October 31, 1986

Date of Report: August 12, 1987

Study Author: Mark D. Nemec, B.S., Study Director. (The
original study director was Dean Rodwell from
initiation of study to issuance of draft for this
final report, but he resigned.) Sponsorship of
the study was transferred from Shell Development
Company to DuPont on October 31, 1986.

Test Compound: Bladex technical, 100% (SD 15418, WRC 107F,
Cyanazine) CAS No. 21725-46-2. Three batches of
compound were used for this study. All received
from Shell Development Company, Westhollow Research
Center. Two were numbered TX 207 with an expira-
tion date of April 1987. The third was numbered
TX 213 with an expiration date of August 1987.

Doses: 0, 25, 75, 150, and 250 ppm.

Route of Administration: Admixed in diet.

Quality Assurance:

A statement of compliance with GLP is signed by Deborah L. Little, Supervisor of Quality Assurance, dated August 12, 1987. Three deviations from GLP are indicated:

1. Five of each sex were not screened for ectoparasites and endoparasites prior to initiation of the study.
2. "There were slight deviations from protocol in the number of animals ordered and the age of animals at receipt."
3. Temperature and humidity "occasionally varied from the specified ranges."

It was concluded that these deviations did not affect the validity of the study.

A previous "initial" two-generation reproduction study with technical Bladex was initiated on August 7, 1985 but was aborted on August 28, 1985 due to suspicion of infection, possibly with a virus. The results of gross necropsies on 10 males and 10 females in control group and serological tests for 2 males and 2 females in control and high-dose groups did not reveal the cause of poor weight gain in the controls, which was the reason for aborting the study.

A summary of the initial study was presented in Appendix A. The procedure was identical to the main study. Initial starting weight of males was about 269 g and of females about 170 g in controls and in all 4 treated groups. Control males gained an average of 60.4 g the first week but lost 50.8 g the second week. Control females gained 13.9 g the first week but lost 18.8 g the second week. In contrast, all male and female treated groups gained weight the first and second weeks of the study and weighed significantly more than controls ($p < 0.01$). The decreased body weight gain in controls was accompanied by a significantly decreased food intake.

Necropsied animals at gross pathology (21st day) revealed no apparent abnormalities. Two controls and 2 high dose per sex were tested for viruses, including SDA/RCV, Sendai, PVM, Reo-3, KRV, H-1, CD-7, MAD, and LCMV. None were found positive to any virus.

Conclusion for Aborted Study:

The reason for poor performance of both the male and female control groups is unknown.

Diet Preparation and Analysis

Each test diet preparation was made by weighing out the proper amount of Bladex, dissolving in 150 mL acetone, then adding to 5 kg powdered rat chow as a premix. Diets were freshly prepared, usually at 3-week intervals. Appendix B listed dates of preparation for each diet and details of preparation.

Analytical support data pertaining to verification of compound by spectroscopy, its purity, and its composition in each dietary preparation was given in Appendix E. The Bladex preparation used in this two-generation study was entirely Sample No. 107F, also coded as Lot No. 06 AMK-5406. It was produced in 1979 by Shell Chemical Company "Mobile Plant." Analyses in 1980, April 1985, and August 1986 (the latter 2 date during the course of this rat study) all showed purities of 98 or 97.5 percent, considered "essentially unchanged" by the analytical chemist.

During the course of the main study, 25 mixings for each dietary dose were prepared. A stability check of dietary preparations of 25 and 250 ppm at 0, 2, 4, and 6 weeks of storage at 5 °C revealed virtually no change with time. A second diet stability test in which a 25 ppm diet was analyzed at 0, 3, 7, 24, and 27.5 h after preparation and stored at both 5 and 25 °C gave results suggesting slightly greater stability at 5 °C. Conclusion by the analytical chemist was that both the 25 and 250 ppm diet preparations were stable at 5 °C, refrigerator temperature, for at least 6 weeks (Table 1 and 2 of report). Storage of diets was therefore in the refrigerator.


Of the 25 mixes for each dietary dose level prepared during the course of this study, 5 were tested for homogeneity by taking samples from 3 different layers of the stored materials labeled as "top, middle, and bottom third." Variation was small, summarized as $\pm 9\%$ at worst, "usually much less" (Tables 3, 4 and 6, 17, 26).

All 25 mixes for each dietary preparation were tested for analytical concentration of Bladex, usually after 2 to 3 days of refrigerated storage following its preparation. A number of them were analyzed 5 to 16 days after preparation. The vast majority of analyses fell below theoretical concentrations and the means came to -10.2, -11.5, -11.2, and -10.4 percent for the 25, 75, 150, and 250 ppm preparations. Occasionally, diets were as much as 25.2 or 33.2 percent below theoretical concentration.

The report on analysis of technical Bladex in rat chow is signed by J.R. Dawson, research chemist, dated February 20, 1987.

IV. STUDY DESIGN: (Figure 1)

A diagrammatic representation of the experimental design has been presented below in Figure 1.



Method

Sprague-Dawley Crl; COBS CD (SD) BR rats of Charles River Portage MI 28 males and 28 females per group. starting at about 7 weeks of age. received 0, 25, 75, 150, and 250 ppm admixed in diet for a minimum of 72 days. Freshly mixed diets were prepared every 3 weeks. Dates of mating were determined by checking each day for vaginal sperm or copulation plugs. First F₀ mating occurred when the rats were about 17 weeks old. All females were allowed to give birth and all litters were culled on day 4 to 8 pups, half of each sex if possible. Offspring of F_{1a} litters that survived were all killed when weaned at 21 days of age. After a minimum of 10 days after weaning, F₀ parents were again bred (1 male to 1 female) to produce F_{1b} litters. F_{1b} pups were weaned at 21 days of age. when 28 of each sex per dose group were randomly selected as F₁ generation parents and the remainder were discarded. F₀ adults were necropsied at time of weaning the F_{1b} litters. Control and high-dose-treated F₀ adults were subjected to histopathology evaluation.

F₁ selected parents were treated for a minimum of 70 days. First mating at 16 to 17 weeks of age was around week 39 of the study to produce F_{2a} litters. F_{2a} litters were discarded at weaning, age 21 days. and about 10 days later the F₁ parents were again mated to produce F_{2b} litters. F_{2b} pups were weaned at 21 days of age when 10 pups of each sex per group were randomly selected for necropsy and gross pathology. Those on high dose and controls were subjected to histopathology evaluation. The remainder of the pups were discarded. All F₁ parents were necropsied after weaning the F_{2b} pups but only those in control and high-dose groups were selected for histopathology evaluation.

All litters of F₁ and F₂ generations were randomly culled to 8 pups maximum, 4 of each sex when possible, on day 4 postpartum. If a female did not mate, it was paired to another male at the same dose level a week later. Offspring dying during days 0 to 4 of lactation were subjected to a necropsy similar to the Staples technique. Normal gross necropsy was performed on those dying after day 4 and only tissues that appeared abnormal were preserved in formalin. No necropsies were performed on F_{1a} weanlings and nonselected F_{1b} weanlings. All F_{2a} weanlings and nonselected F_{2b} weanlings were also discarded without necropsy examination. Macroscopic pathology was performed on all F₀ and F₁ male and female parents, also the selected F_{2b} pups, which included all orifices and organs, external and cut surfaces of brain and spinal cord. Weights of liver, right and left testes, and ovary from adults were obtained. Histopathology was limited to grossly observed lesions and reproductive organs in males testes, epididymides, prostate, and seminal vesicles. whereas in females ovaries, uterus and vagina.

Statistics included Chi-square tests with Yates correction factor for F_0 and F_1 male and female gestation indices, also for stillbirths and pup sex ratios on day 1 lactation in all 4 reproduction studies. Dunnett's test was performed on mean viable pups, gestation length, survival indices, and litter weights on days 1, 4, 7, 14, and 21 of F_{1a} , F_{1b} , F_{2a} , and F_{2b} pups, parental weight and weight gains of F_0 and F_1 adults including before mating, during mating, gestation, and lactation; also food consumption, organ weights.

It should be noted that all exposures to compound admixed in diets were continuous for all F_0 and F_1 animals throughout the entire study. This also means that pups closer in age to weaning could feed on the diet or at an earlier age, particularly before growth of fur, could possibly absorb compound through the skin.

Results:

F_0 Generation:

Mortality - One male on 75 ppm dose during week 17, 2 females on 25 ppm during weeks 13 and 24, 3 females on 75 ppm during weeks 1, 24 and 29, 1 female at 250 ppm during week 15; another female on 150 ppm was killed in moribund state during F_{1b} delivery. Deaths were not attributed to treatment.

Clinical Observations - No compound-related effects.

Body Weight and Weight Gains - (Obtained weekly). Mean body weight at the initiation of the study before treatment was the same in controls and in all 4 treated groups. At weeks 1 and 2 weighings, body weight of all 4 treated male and female groups became higher than controls ($p < 0.01$) because of a 22 g lower increase in weight gain for controls of both sexes at week 1 versus the lowest dose 25 ppm groups. Nevertheless, there was clearly a dose-related decreased body weight gain with both sexes for the 3 highest dose groups if comparisons for them were made to low-dose-treated group. By week 8 in males and week 5 in females, 250 ppm highest dose became significantly lower than respective controls. Then, by week 9 in males and week 10 in females, the 150 ppm dose group became significantly lower than controls. The 2 highest dose groups remained lower, dose-related, than controls to week 30, the time of F_0 necropsy of males and females, signifying toxicity of the 2 highest doses, 150 and 250 ppm treated males and females. The dose-related trend in body weight decrease included the 75 ppm group, especially the females, where it became significant at weeks 8 to 10. Overall, from weeks 0 to 30, both males and females in the 3 highest dose groups showed a dose-related

decreased weight. Throughout F_{1a} and F_{1b} gestation and lactation body weights were significantly lower than controls in the dams of the 2 highest dose treated groups at most time periods ($p \leq 0.05$ to 0.01). Generally, these were not due to further weight decrease in the 2 highest dose treated groups but were rather due to lower initial weights caused earlier by drug treatment, i.e., prior to gestation and lactation.

Food Consumption - (obtained weekly). Tables are presented both in terms of g/animal/day and g/kg/day, based on mean values for each dose group. During the first 2 weeks, the amount consumed by male and female controls was unusually low, reflecting poor weight gain. Therefore, food consumption, based on g/animal/day or g/kg/day, was higher in all 4 treated groups than in controls of both sexes ($p \leq 0.01$). The reason for this decreased food intake by controls during this time period could not be explained. From weeks 3 to 6 in males, all 4 treated groups had significantly lower feed intake than controls, based on g/kg/day, but in females, significantly lower feed intake by all 4 treated groups occurred during the third week, then during the fifth week. Feed consumption was usually lower in treated groups than in controls (based on mg/kg/day), especially at the 2 highest dose levels for both sexes. During the F_{1a} and F_{1b} gestation periods, there were no obvious effects of compound treatment on food intake. Diminished food intake was seen during both lactational periods, significant only during the F_{1a} lactation for the 250 ppm group.

Mean compound intake in mg/kg/day based on food intake measurements were as follows:

Dose in ppm, F ₀ Parents	0	25	75	150	250
	Compound Dosage (mg/kg/day)				
Females, excluding time of breeding, gestation, lactation	-	1.67	4.97	10.00	16.66
Females during F _{1a} gestation	-	1.83	5.25	10.65	18.50
Females during F _{1b} gestation	-	1.75	5.25	10.50	17.50
Females during F _{1a} lactation	-	2.98	9.15	19.80	26.50
Females during F _{1b} lactation	-	3.75	11.03	22.20	36.20
Males	-	1.43	4.32	8.80	15.26

Reproductive Performance

Data in the table that follows, Summary of Reproductive Performance, were compiled from 6 tables of the report (Tables 28 to 33). Males and females in the same dose groups were cohabited in pairs.

There is some questions on reliability of the data on Females with Evidence of Mating in the table. Dates of insemination were missed for a number of females. It also seems possible to have mistakenly considered an animal to be inseminated particularly if based on a copulation plug under the cage rather than vaginal sperm. 006597

Data on "Male Mating Index" and "Female Mating Index" were presented combined "for both mating periods", which we believe signifies for the F_{1a} and F_{1b} generations. In its present form these values would be higher than if they were not combined for the 2 mating periods and it is not possible to determine the contribution of these values to the reproductive performance of each group for each of the 2 generations.

The number of dams delivering a litter was reduced by 35% in the 150 ppm groups for both the F_{1a} and F_{1b} matings (See table which follows). The reason for the decreased number delivering a litter in the 150 ppm groups of both matings is unknown. It could have been due to decreased fertility, increased pregnancy terminations or other reasons. It may have been "a random occurrence" as suggested by the applicant, since it did not occur at the 250 ppm dose group, nor was it observed in the F_{2a} and F_{2b} generations. Since it occurred in 2 generations, we should not ignore the possibility of an effect.

There were no effects of treatment on length of the gestational period, sex ratio of pups at birth, or on mean gestation index (see footnote in table that follows) in the F_{1a} or F_{1b} litters. In the F_{1b} part of the study not shown in the table that follows, 1 dam on 75 ppm died on day 22 of gestation, 1 dam on 150 ppm was killed in moribund state on day 23 of gestation, both of dystocia. It was claimed that dystocia was not a compound-related effect. There was no effect of compound on live litter size at birth and no increased number of dead pups at birth in the treated groups. There was an unusually large number of pups born dead or that died within the first 24 hours of birth in the control group of the F_{1b} generation, which was responsible for significantly lower number of dead pups in the treated groups.

SUMMARY OF REPRODUCTIVE PERFORMANCE, F_{1a} and F_{1b} COMBINED

Dose Group (ppm)	0	25	75	150	250
Number of Females/GP ^o - F _{1a}	28	28	27	28	28
- F _{1b}	28	27	27	28	27
Male Mating Index (%) +	100(28/28)	96(27/28)	100(28/28)	89(25/28)	100(28/28)
Female Mating Index (%) +	96(27/28)	100(27/27)	100(27/27)	86(24/28)	89(25/27) ^c
Females with Evidence of Mating [#] - F _{1a} (%)	100	89	96	93	89
- F _{1b} (%)	96	93	96	89	96
Number Delivering a Litter, F _{1a}	26	24	26	17	24
Number Delivering a Live Litter, F _{1a}	26	24	26	17	23
Gestation Length \pm SD, F _{1a} (days)	22.2 \pm 0.6	22.0 \pm 0.7	21.9 \pm 0.6	21.9 \pm 0.3	21.9 \pm 0.5
Number Delivering a Litter, F _{1b}	25	23	23	16	22
Number Delivering a Live Litter, F _{1b}	24	22	22	15	21
Gestation Length \pm SD, F _{1b} (days)	22.8 \pm 0.8	22.0 \pm 0.7	22.0 \pm 0.7	21.9 \pm 0.6	21.8 \pm 0.6
Male Ratio in Litter - F _{1a} (%)	49.5	53.6	51.5	53.8	45.3
- F _{1b} (%)	54.1	47.0	52.1	50.2	49.0
Gestation Index - F _{1a} (%)	100	96	100	100	100
- F _{1b} (%)	96	96	96	88	95
Mean Live Litter Size at Birth - F _{1a}	13.2	13.3	13.7	14.9	13.9
- F _{1b}	12.8	13.2	13.2	13.5	12.6
Total Number of Pups Born Dead in all Litters, Combined - F _{1a}	5	8	9	4	7
- F _{1b}	21	2**	2**	8	4**

^o There were 28 males in every group during F_{1a} and F_{1b}.

+ Data for F_{1a} and F_{1b} were combined.

Evidence of mating was based on vaginal sperm.

**p < 0.01 by Chi-Square test.

Mating Index = $\frac{\text{Number mating that resulted in pregnancy} \times 100}{\text{Total number cohabited}}$

Gestation Index = $\frac{\text{Number giving birth to a live litter}}{\text{Number of pregnancies}}$

^c Includes 1 female that didn't deliver but had implantation scars at necropsy.

Litter Performance

The tables on the 2 pages that follow are summaries of viability indices and body weights during lactation of F_{1a} and F_{1b} pups as an indication of the performance of the offspring.

A reduction in viability index was seen in the F_{1a} 250 ppm treated group by day 14 ($p < 0.01$) with reductions also seen for 75 and 150 ppm groups by day 21 but not statistically significant. Six dams in the 250 ppm group and 1 in the 150 ppm group had lost their entire litters whereas entire litter loss did not occur in dams of control or in lower dose treated groups. Sharply decreased mean pup weights were seen in the 150 and 250 ppm groups ($p < 0.05 - 0.01$) starting on day 4 and continuing throughout lactation. Decreased mean pup weights were also seen in the 75 ppm treated group by day 14 (n.s.) and on day 21 ($p < 0.05$) of lactation.

In the F_{1b} generation there was an unusually high level of mortality among pups in the control group especially between day 0 to 1 and again between days 14 to 21 of lactation. Based on the poor performance of the control group in this study, there were no effects of treatment on viability index and only the 250 ppm dose caused a decrease ($p < 0.01$) in body weight between days 14 to 21 of lactation.

SUMMARY OF PUP VIABILITY INDICES

F₁A Pups
Viability Index on Lactation Day (A)

	Day 1	Day 4 Before Culling	Day 4 After Culling	Day 7	Day 14	Day 21(B)
Dose ppm	Z	Z	Z	Z	Z	Z
0	99.8	98.7	100.0	100.0	100.0	99.5
25	98.3	97.5	100.0	98.9	98.9	98.9
75	99.5	98.5	100.0	99.5	97.7	92.9
150	98.1	95.1	100.0	99.3	98.6	89.8
250	94.9	93.2	100.0	99.5*	76.2**	75.0**

F₁b Pups
Viability Index on Lactation Day (A)

	Day 1	Day 4 Before Culling	Day 4 After Culling	Day 7	Day 14	Day 21(B)
Dose ppm	Z	Z	Z	Z	Z	Z
0	95.3	94.2	100.0	99.5	98.5	94.8
25	96.5	95.9	100.0	98.9	98.3	98.3
75	99.0	97.5	100.0	98.9	98.9	98.9
150	93.9	90.9	100.0	99.2	91.7	91.7
250	98.1	96.5	100.0	99.4	98.2	94.7

(A) = Mean values calculated using individual values.

(b) = lactation index.

Pup survival compared using Dunnett's test.

**p < 0.01 (Dunnett's test).

Viability
Index (Z)

Before

Culling = $\frac{\text{No. Pups available per litter Day 4 before culling}}{\text{No. pups viable per litter Day 1}} \times 100$

Viability
Index (Z)

After

Culling = $\frac{\text{No. pups viable per litter on Day (N)}}{\text{No. pups viable per litter on Day 4 after culling}} \times 100$

SUMMARY OF MEAN PUP WEIGHTS DURING LACTATION

Dose (ppm)	0	25	75	150	250
N (on Day 0)	26	24	26	17	23
		F _{1a} Litters			
			Mean + SD		
Day 1	6.6 + 1.0	6.6 + 1.0	6.6 + 0.5	6.2 + 0.6	6.1 + 0.6
4+	9.3 + 1.5	9.3 + 1.7	9.0 + 0.8	8.1 + 1.2*	8.2 + 1.2*
4 =	9.3 + 1.6	9.3 + 1.6	9.0 + 0.9	8.1 + 1.3*	8.3 + 1.3*
7	14.9 + 2.3	15.4 + 2.1	14.9 + 1.5	12.8 + 2.6*	11.5 + 2.9*
14	24.1 + 3.6	24.4 + 3.8	24.9 + 4.5	20.6 + 3.8*	20.1 + 4.1*
21	40.0 + 6.0	37.9 + 6.2	34.8 + 6.9*	28.1 + 5.2**	29.5 + 6.0*

N (On Day 0)	24	22	22	15	21
		F _{1b} Litters			
			Mean + SD		
Day 1	6.7 + 0.8	6.6 + 1.1	6.7 + 1.1	6.3 + 0.7	6.5 + 0.8
4+	9.3 + 1.3	9.1 + 1.9	9.4 + 1.9	8.5 + 1.3	9.0 + 1.6
4 =	9.3 + 1.3	9.3 + 1.8	9.4 + 1.9	8.5 + 1.3	9.0 + 1.6
7	14.6 + 1.7	14.5 + 2.9	14.8 + 2.5	13.1 + 2.9	13.2 + 2.5
14	26.2 + 3.5	25.3 + 5.2	25.5 + 4.6	24.0 + 3.0	23.9 + 3.7
21	37.3 + 7.2	38.6 + 6.3	38.2 + 7.1	34.1 + 3.4	30.5 + 4.8*

N = Number of litters.

4+ Before culling on Day 4

4= After culling on Day 4

*p < 0.05 by Dunnett's test.

**p < 0.01.

Litter Losses During Lactation

For F_{1a}, 6 litters were entirely lost (100% dead) in 250 ppm group between days 7 to 14. For F_{1b}, 1 litter was entirely lost in 150 ppm group between days 7 to 14.

General Physical Condition and Mortality

This is a narrative summary of individual animal necropsy data from tables 117, 118, and 118A in volume 2. It includes all animals found dead in the F_{1a} and F_{1b} litters during the 21 days of lactation (Tables 117 and 118) and those found dead in F_{1b} litters between age 21 to 30 days, prior to selection of F₁ parents. These tables obviously include among the dead those that were missing and not actually found dead. For example, in the F_{1a} high dose group, dam #23703 lost all her pups between days 7-14 of lactation but only 3 of them are accounted for in the necropsy data of Table 117. Dam #23748 also lost all 8 pups between days 7-14 of lactation but only 2 of them are accounted for. These are only two examples which require accountability for missing pups.

In F_{1a} litters, there were 8 control deaths, each of them from a different litter. One male had external anasarca and brachydactyly. At 25 ppm, there were 7 deaths in 1 litter and 5 additional deaths in separate litters. One female had distended bilateral ureters. At 75 ppm there was a total of 22 deaths in 10 litters, of which 8 occurred in a single litter, 2, 3, and 3 deaths in 3 other litters, 1 in each of 6 other litters. No remarkable findings were found. At 150 ppm, there were 25 dead pups within 8 litters, of which there was 5, 7, 6, 2, and 2 deaths in litters with more than 1 death. No remarkable findings occurred. At 250 ppm, there was a total of 61 deaths within 15 litters. Multiple deaths (i.e., 2 or more) in 11 litters occurred numbering 9, 3, 2, 2, 2, 7, 3, 2, 8, 8, and 3. No remarkable findings were observed. A large number of pups on 250 ppm dose, at least 30, were small in size, but otherwise "not remarkable."

In the F_{1b} litters, controls had 29 deaths in 10 litters with multiple deaths in 7 litters, in which there were 3, 4, 3, 3, 7, 3, and 2 deaths; no remarkable findings in gross pathology. In 25 ppm, there were only 3 deaths all occurring within 3 separate litters; no remarkable findings in gross pathology. At 75 ppm, there were only 3 deaths in 3 separate litters; no remarkable findings. At 150 ppm, there were 6 deaths in 4 litters, 2 of the litters with 2 deaths. At 250 ppm, there were only 8 deaths in 5 litters, 4 of them occurring in 1 litter; no remarkable observations at gross pathology examination.

Of the F_{1b} rats that were being held to day 30 for possible use as F₁ generation adults that died, 3 controls from 3 separate litters were found. At 25 ppm, 3 from 3 litters were found. At 75 ppm, only 1 died. At 150 ppm, only 1 died. At 250 ppm, 34 from 10 litters died. No remarkable findings were reported for the necropsy results.

Pathology of F₀ Parents

Necropsy of Those Dying During the Study (From Table 36. Vol. 1 and from Table 119 in Vol 3).

No controls died. At 25 ppm 2 females died 1 on day 96 of study (day 20 estimated gestation) of renal failure the second on day 168 of undetermined cause. At 75 ppm. 1 male on day 124. 3 females on days 8. 171. and 208 of undisclosed causes. At 150 ppm. 1 female died during late pregnancy on day 132. At 250 ppm. 1 female died on day 109 of study of undetermined cause. It is claimed that no compound related deaths were evident.

Scheduled Necropsy (Week 30- days 210 and 211)

The table that follows clearly illustrates the dose-related decreased body weights in the treated 75. 150. and 250 ppm males and females on the day of scheduled necropsy.

Final Body Weight + SD at Terminal Sacrifice Week 30

Dose (ppm)	0	25	75	150	250
Males	534.0 + 40.7	550.1 + 54.5	522.4 + 45.4	474.3 + 36.5**	450.8 + 40.3**
(Number)	(28)	(28)	(27)	(28)	(28)
Females	314.9 + 24.1	309.6 + 22.2	301.3 + 25.9	282.9 + 15.2**	272.6 + 27.6**
(Number)	(28)	(26)	(25)	(27)	

**p < 0.01 Dunnett's test.

Organ Weights (from Table 38 in Vol. 1 of report)

In the table below. the decreased absolute weights of liver of both sexes occurred only in the 250 ppm dose treated groups of males and females. However. there was no effect on relative liver weight at any dose level for either sex. There were also no effects on absolute weights of testes or ovaries nor on relative weights of ovaries. The increased relative weights of testes in the 2 highest dose groups was considered by the investigators as being not biologically meaningful since it was not accompanied by macroscopic or microscopic changes. The conclusion was that there were no effects on liver or gonadal weights in either sex.

06597

Organ Weights at Necropsy in Males (From table 40 of report)

Dose (ppm)	0	25	75	150	250	0	25	75	150	250
			Absolute (g \pm SD)					Relative (g \pm SD)		
Liver	20.1 \pm 2.6	20.8 \pm 3.6	19.6 \pm 3.1	18.3 \pm 2.9	17.7 \pm 2.3*	3.76 \pm 0.40	3.78 \pm 0.50	3.75 \pm 0.43	3.86 \pm 0.53	3.93 \pm 0.37
Left Testis	1.72 \pm 0.14	1.72 \pm 0.17	1.75 \pm 0.16	1.71 \pm 0.21	1.80 \pm 0.15	0.32 \pm 0.03	0.31 \pm 0.03	0.34 \pm 0.04	0.36 \pm 0.04**	0.40 \pm 0.04**
Right Testis	1.73 \pm 0.14	1.71 \pm 0.17	1.84 \pm 0.43	1.73 \pm 0.18	1.77 \pm 0.18	0.33 \pm 0.03	0.31 \pm 0.03	0.35 \pm 0.08	0.37 \pm 0.03**	0.40 \pm 0.05**

Organ Weights at Necropsy in Females (From table 39 of report)

Dose (ppm)	0	25	75	150	250	0	25	75	150	250
			Absolute (g \pm SD)					Relative (g \pm SD)		
Liver	12.8 \pm 1.49	12.5 \pm 1.2	12.6 \pm 1.3	12.0 \pm 1.9	11.6 \pm 2.1*	4.07 \pm 0.31	4.05 \pm 0.26	4.18 \pm 0.31	4.24 \pm 0.47	4.25 \pm 0.54
Left Ovary	0.042 \pm 0.012	0.046 \pm 0.014	0.042 \pm 0.011	0.039 \pm 0.015	0.040 \pm 0.014	0.014 \pm 0.004	0.015 \pm 0.005	0.014 \pm 0.004	0.014 \pm 0.005	0.015 \pm 0.005
Right Ovary	0.042 \pm 0.010	0.040 \pm 0.012	0.042 \pm 0.012	0.037 \pm 0.014	0.038 \pm 0.012	0.014 \pm 0.003	0.013 \pm 0.003	0.013 \pm 0.003	0.013 \pm 0.004	0.014 \pm 0.004

* $P < 0.05$ by Dunnett's test

* $P < 0.01$

006597

14

Gross Necropsy of F₀ Parents (From Table 37 and 119 of the report).

A summary table for gross pathology observations at terminal kill for all males and females was presented but no effects of treatment appeared obvious in any organ.

Microscopic Pathology of F₀ Parents (From Table 41)

Lesions observed at necropsy in all animals and reproductive organs from all on highest (250 ppm) dose and controls were included. No compound related effect was observed.

Results with F₁ Parents

Mortality of F₁ Adults - One male control, 1 female 75 ppm dose, one female 150 ppm dose found dead weeks 45, 57, and 53, respectively.

Clinical Observations - No compound-related effect was obvious throughout the study at any dose level.

Body Weight and Body Weight Gains (Weeks 28 to 59 of the study). Decreased body weight was already apparent at week 28 for the 75 ppm (n.s.), 150 ppm ($p < 0.05$), and 250 ppm ($p < 0.01$) doses when F₁ parent exposure was just starting after weaning and mean control weight was 57.8 g. By week 33, mean weight of 75 ppm males and females were significantly less than controls and those on 25 ppm were also decreased (n.s.) but the decreased body weight for the 25 ppm dose became significant for males at about 45 weeks and after, for females at 34 weeks and after. Total body weight gains for the entire period between weeks 28 to 59 were significantly depressed in all 4 treated male and female groups in a dose-related manner.

During both the F_{2a} and F_{2b} gestations, mean body weights of dams were significantly lower than controls (dose related) in all 4 treated groups at every time period, including days 0, 6, 12, 15, 18, and 20. Total body weight gain throughout the F_{2a} gestational period including days 0 to 20 were significantly depressed only for the 150 and 250 ppm groups and for the F_{2b} generation only for the 250 ppm group.

During lactation, body weights were significantly depressed in a dose-related manner from day 1 of lactation on to day 21 in both F_{2a} and F_{2b} generation. However, there was generally no significant decrease in body weight gain due to treatment during lactation.

Food Consumption - Obtained weekly. When based on g/animal/day, food intake for F₁ males was decreased for the 75, 150, and 250 ppm dose groups from weeks 28 to 59 of the study ($p < 0.01$ at every time period throughout the study for the 2 highest doses). When converted to g/kg/day, mean food intake was no longer less than controls in the treated groups; in fact, it tended to be significantly higher ($p < 0.01$) at most of the time periods for the 250 ppm group, sometimes higher ($p < 0.05$ to 0.01) for the 150 ppm group. Obviously, the higher food intake based on g/kg/day is due to a correction for higher mean body weights: body weight was highest for controls and inversely dose related in the treated groups. The females, during non-gestation or non-lactation periods, similarly had decreased food intake at most of the time periods for the 75, 150, and 250 ppm doses, if based on g/animal/day. When converted to g/kg/day, there were similarly no differences in food intake for the 25, 75, and 150 ppm doses, but it was generally increased for 1/3 of the time periods with the 250 ppm dose. Mean food consumption during the F_{2a} and F_{2b} gestational and F_{2a} lactation periods were lower than controls for most time periods, especially for the 75, 150, and 250 ppm doses only when based on g/animal/day but was no longer apparent if based on g/kg/day. Differences in treated animal food intake were not apparent during F_{2b} lactation whether based on g/animal/day or g/kg/day.

The following was an estimation of dosage as mg/kg/day at various stages of the study, based on feed consumption measurements.

Feed Content of Bladex (ppm)	0	25	75	150	250
	Compound Dosage (mg/kg/day)				
Males and females, excluding breeding, gestation, lactation	-	2.01	6.04	12.26	21.08
During F _{2a} gestation (mg/kg/d)	-	1.90	5.55	11.25	20.25
During F _{2b} gestation (mg/kg/d)	-	1.73	5.10	12.20	17.25
During F _{2a} lactation (mg/kg/d)	-	4.23	12.60	24.30	42.00
During F _{2b} lactation (mg/kg/d)	-	4.13	11.93	25.80	43.75

Reproductive Performance, F₁ Rats

Males were cohabited with females that had received the same dosage of compound for about 70 days, starting after weaning. F_{2a} mating occurred during weeks 40 to 43, F_{2b} mating occurred at 50 to 53 weeks of the study.

Data in the table which follows were compiled from Tables 69-73 of the report, which basically is similar in presentation as the previous data on Reproductive Performance. We are similarly

requesting that the data which were combined for the Male and Female Mating Indices be separated for the F_{2a} and F_{2b} generations. There is again the question of results on Females with Evidence of Mating and the data on any Reproductive Indices which may have been derived from these results.

Based on the data in the table on Reproductive Performance which follows, there was no reduction in females delivering a litter in any treated group, no evidence of increased number of dead litters or of dystocia at parturition. Although pregnancy duration was 22.2 days for control groups of both the F_{2a} and F_{2b} pregnancies, gestation length was apparently reduced ($p < 0.05$ to 0.01) in all 4 treated groups by 0.5 to 0.8 days of the F_{2a} but no such effect for the F_{2b} pregnancy. The applicant considered the decreased gestation period in the F_{2a} pregnancies as not biologically meaningful. Gestation index was 100 percent in virtually all control and treated groups for F_{2a} and F_{2b} pregnancies. In addition, there were no effects on live births per litter. Although total number of pups born dead was increased for the 75 ppm group of the F_{2b} litters ($p < 0.05$), it was considered a sporadic effect due to 2 litters with all or almost all pups dead on day 0. Sex ratio at birth was similar in control and treated groups for litters of both generations.

SUMMARY OF REPRODUCTIVE PERFORMANCE; F_{2a} and F_{2b}

Dose Group (ppm)	0	25	75	150	250
Number in Group; each sex	28	28	28	28	27=
Mating Index, Males (%) ⁺	96(27/28)	96(27/28)	100(28/28)	100(28/28)	93(25/27)
Mating Index, Females (%) ⁺	96(27/28)	96(27/28)	96(27/28)	96(27/28)	93(26/28)
Females with Evidence of Mating [#] - F _{2a} (%)	82	93	96	100	96
- F _{2b} (%)	93	93	96	96	100
Number Females Delivering (F _{2a})	27	25	27	25	23
Number Delivering Live Litter, (F _{2a})	23	24	26	25	23
Gestation Length \pm SD, (F _{2a})	22.1 \pm 0.6	21.5 \pm 0.6**	21.4 \pm 0.5**	21.3 \pm 0.5**	21.6 \pm 0.5*
Number Females Delivering (F _{2b})	23	22	25	25	23
Number Delivering Live Litter, (F _{2b})	22	21	24	24	23
Gestation Length \pm SD, (F _{2b})	22.1 \pm 0.4	22.1 \pm 0.6	21.9 \pm 0.7	22.0 \pm 0.4	21.8 \pm 0.5
Gestation Index - F _{1a} (%)	100	100	100	96	100
- F _{1b} (%)	100	100	100	100	100
Live Litter Size F _{2a}	12.4	13.7	12.7	12.3	11.0
Live Litter Size F _{2b}	13.0	13.2	12.8	13.4	13.2
Total Dead Pups at Birth in all Litters - F _{2a}	11	5	7	21	5
- F _{2b}	10	4	28\$	4	7
Male Ratio in F _{2a} (%)	45.6	51.6	55.6	48.5	43.4
Male Ratio in F _{2b} (%)	50.7	50.9	50.8	50.0	50.3

+ Includes both mating periods. Apparently, F_{2a} and F_{2b} were combined.

= 27 males, 28 females were included.

Based on vaginal sperm in paired female or copulation plug.

\$ p < 0.05 by the Chi-Square test (comparing total) number of dead in all litters combined per group.

*p < 0.05 by Dunnett's test.

**p < 0.01 by Dunnett's test.

Live litter size = total number of pups/number of litters.

Mating Index = $\frac{\text{Number mating that became pregnant} \times 100}{\text{Total number cohabited}}$

Gestation Index = $\frac{\text{Number giving birth to live litter} \times 100}{\text{Number of pregnancies}}$

Litter Performance

Summaries of mean live litter size based on viability indices and of mean pup body weights throughout lactation are given in the 2 tables that follow as indications of performance of the offspring.

Viability index was decreased only on days 1 and 4 before culling in the 150 and 250 ppm groups of the F_{2a} generation. No significant reduction in viability index was seen after culling in any treated groups of both the F_{2a} and F_{2b} generations. However, decreased pup weights were clearly evident during lactation even after culling on day 4 in a dose-related manner for the 2 highest doses in the F_{2a} generation and for the 3 highest doses in the F_{2b} generation.

There was no effect of compound treatment at any dose level on number of lactating dams that lost the entire litters.

006597

SUMMARY OF PUP VIABILITY INDICES
F₂A Pups
Viability Index on Lactation Day (A)

Dose ppm	Day 1 %	Day 4 Before Selection %	Day 4 After Selection %	Day 7 %	Day 14 %	Day 21(B) %
0	98.9	98.6	100.0	99.6	97.7	97.7
25	99.7	98.9	100.0	99.5	99.5	99.5
75	98.0	97.3	100.0	100.0	100.0	99.6
150	83.2**	82.1**	100.0	99.5	99.5	99.5
250	85.6*	83.6*	100.0	98.4	96.3	90.9

F₂b Pups
Viability Index on Lactation Day (A)

Dose ppm	Day 1 %	Day 4 Before Selection %	Day 4 After Selection %	Day 7 %	Day 14 %	Day 21(B) %
0	99.7	99.7	100.0	100.0	100.0	99.0
25	97.3	96.1	100.0	98.3	98.3	98.3
75	95.0	92.3	100.0	98.5	92.3	90.2
150	97.2	96.4	100.0	99.5	98.0	92.6
250	94.7	89.6	100.0	96.8	95.7	94.6

(A) = Mean values calculated using individual values.

(B) = lactation index.

Pup survival compared using Dunnett's test.

*p < 0.05.

**p < 0.01 (Dunnett's test).

Viability

Index (%)

Before

Culling = $\frac{\text{No. pups viable per litter Day 4 before culling}}{\text{No. pups viable per litter Day 1}} \times 100$

Viability

Index (%)

After

Culling = $\frac{\text{No. pups viable per litter on Day (N)}}{\text{No. pups viable per litter on Day 4 after culling}} \times 100$

SUMMARY OF MEAN PUP WEIGHTS DURING LACTATION

Dose (ppm)	0	25	75	150	250
N (on day 0)	27	25	27	22	22
	F _{2a} Litters: Mean + SD				
Day 1	6.6 ± 0.7	6.1 ± 0.5	6.1 ± 0.8	6.0 ± 0.9	6.1 ± 1.1
4+	9.5 ± 1.2	8.4 ± 0.8*	8.5 ± 1.2	8.3 ± 1.6	8.1 ± 1.8**
4 =	9.5 ± 1.2	8.4 ± 0.8*	8.5 ± 1.2	8.3 ± 1.7*	8.1 ± 1.9**
7	15.1 ± 1.5	14.1 ± 1.3	13.9 ± 1.8	13.0 ± 2.8*	12.2 ± 3.1**
14	28.8 ± 3.1	28.2 ± 2.6	28.7 ± 3.7	26.9 ± 3.7	23.6 ± 5.0**
21	43.6 ± 3.8	41.1 ± 2.8	41.7 ± 3.6	39.4 ± 5.3	36.4 ± 6.7**
N (On day 0)	23	22	25	25	23
	F _{2b} Litters: Mean + SD				
Day 1	6.6 ± 0.7	6.4 ± 0.7	6.2 ± 0.8	6.4 ± 0.6	5.7 ± 0.9**
4+	9.6 ± 1.3	9.1 ± 1.2	8.7 ± 1.2	8.6 ± 1.0	7.5 ± 1.5**
4 =	9.6 ± 1.3	9.1 ± 1.2	8.7 ± 1.2	8.6 ± 1.0*	7.5 ± 1.5**
7	15.7 ± 1.7	15.0 ± 1.6	13.7 ± 2.6**	13.7 ± 1.9**	11.0 ± 2.5**
14	31.6 ± 3.1	29.6 ± 3.0	27.4 ± 5.1**	25.4 ± 5.3**	23.8 ± 4.9**
21	46.3 ± 4.9	44.2 ± 3.5	42.8 ± 4.1	38.5 ± 7.0**	36.1 ± 6.9**

N = Number of litters.

4+ Before culling

4= After culling

*p < 0.05 by Dunnett's test.

**p < 0.01.

Litters Lost During Lactation

For F_{2a}, 1 litter was entirely lost (100% dead) in 250 ppm group between days 7 to 14 of lactation. For F_{2b}, 1 litter was entirely lost in 75 ppm group only between days 14 to 21.

Physical Appearance and Mortality of Pups

This is summarized narratively from the individual animal necropsy tables of pups. It includes only those found dead and "does not include pups found missing or cannibalized," as stated in tables 151 and 152, pages 1515 to 1529 of the report. Examination of the tables on Individual Litter Viability (Tables 145 and 147) and the data on necropsy in Tables 151 and 152 revealed no serious problem in accounting for missing pups.

In the narration, it is claimed that the pups in the 75, 150, and 250 ppm groups of the F_{2a} generation were small. Surprisingly, this small appearance of pups in treated groups is not claimed for F_{2b} pups even though the table above for mean pup weights clearly shows decreased weight of pups in the 3 highest dose treated groups.

In the F_{2a} generation, the only discovered malformation was in a 25 ppm group female pup that died on day 0, which had an agnathia. In the F_{2b} generation, 1 male on 75 ppm had hydrocephaly. With 150 ppm, 1 male had hydrocephaly, a female in the same litter had an ocular defect thought to be anophthalmia and also had anasarca.

In the F_{2a} generation control group, 18 dead pups were found in 7 litters with 4 deaths in 2 litters, 3 deaths in 2 litters, 2 deaths in 1 litter, and 2 litters each with 1 death. At 25 ppm, there were 9 deaths occurring in 5 litters with 3 deaths occurring in 2 of them. At 75 ppm, there were also 9 deaths in 5 litters. At 150 ppm, there were 40 dead pups in 13 litters, 1 of them with 12, another with 8 dead pups. In the 250 ppm group, there were 31 deaths in 12 litters, multiple deaths of 5 or more occurring in 2 of them.

Mortality of pups in the F_{2b} generation included 10 controls in 8 litters, 5 at 25 ppm all in the same litter. At 75 ppm there were 40 deaths in 14 litters, 2 of which had 5 or more deaths. At 150 ppm there were 19 deaths in 9 litters, 2 of them with 5 deaths. At 250 ppm there were 18 deaths in 12 litters, all with 3 deaths or less in a litter.

Gross Necropsy of Pups that Died:

Complete gross necropsies were obviously performed, most with "no remarkable observations," a number of them autolyzed. No discernible compound effect was evident.

Pathology of F₁ Adults

Final Body Weights

Dose-related decreases were seen in males and females.

Final Body Weights + SD at Terminal Sacrifice. Week 30

Dose (ppm)	0	25	75	150	250
Males (g)	611.7 + 65.4	575.7 + 63.8	533.4 + 50.6**	488.3 + 62.4**	447.4 + 46.9**
(Number/group)	(27)	(28)	(28)	(28)	(27)
Females (g)	337.9 + 32.4	310.8 + 28.2**	301.7 + 31.1**	278.7 + 21.6**	271.4 + 17.3**
(Number/group)	(28)	(28)	(27)	(27)	(28)

**p < 0.01 by Dunnett's test.

Organ Weights

Liver weights in both males and females were decreased ($p < 0.01$) in a dose-related manner when based on absolute weight, but these decreases were no longer evident when based on relative weight (corrected for body weight). In fact, relative liver weights were increased in the 150 and 250 ppm female groups. The biological significance of this organ change is unknown. There were no changes in absolute weight of testes or ovaries but on a relative basis there was a tendency for a dose related increase in both the male and female gonads.

ORGAN WEIGHTS AT NECROPSY IN F1 ADULTS

Males	Absolute Weight(g)						Relative to Body Weight					
	GROUP: 0 PPM	25 PPM	75 PPM	150 PPM	250 PPM	0 PPM	25 PPM	75 PPM	150 PPM	250 PPM	0 PPM	25 PPM
LIVER	MEAN	22.36	20.55	19.69**	16.24**	3.660	3.564	3.689	3.741	3.623		
	S.D.	3.325	3.211	2.596	2.214	0.4176	0.3531	0.3127	0.2999	0.2345		
	N	27	28	28	27	27	28	28	28	27		
LEFT TESTIS	MEAN	1.78	1.83	1.84	1.86	0.294	0.321	0.346**	0.390**	0.419**		
	S.D.	0.198	0.170	0.160	0.242	0.0399	0.0350	0.0306	0.0518	0.0612		
	N	27	28	28	27	27	28	28	28	27		
RIGHT TESTIS	MEAN	1.83	1.84	1.85	1.86	0.301	0.321	0.348	0.391**	0.419**		
	S.D.	0.169	0.160	0.156	0.255	0.0345	0.0329	0.0357	0.0505	0.0656		
	N	27	28	28	27	27	28	28	28	27		
Females												
	Absolute Weight(g)						Relative to Body Weight					
	GROUP: 0 PPM	25 PPM	75 PPM	150 PPM	250 PPM	0 PPM	25 PPM	75 PPM	150 PPM	250 PPM	0 PPM	25 PPM
LIVER	MEAN	13.14	12.00*	11.58**	11.70**	3.899	3.860	3.839	4.189**	4.213**		
	S.D.	1.493	1.452	1.438	1.098	0.3662	0.3051	0.3211	0.4040	0.3157		
	N	28	28	27	27	28	28	27	27	28		
LEFT OVARY	MEAN	0.0458	0.0428	0.0515	0.0475	0.014	0.014	0.017*	0.017*	0.016		
	S.D.	0.01426	0.01232	0.01323	0.01168	0.0041	0.0045	0.0044	0.0045	0.0047		
	N	28	28	27	27	28	28	27	27	28		
RIGHT OVARY	MEAN	0.0438	0.0426	0.0471	0.0490	0.013	0.014	0.016	0.018**	0.017**		
	S.D.	0.01424	0.01020	0.01328	0.01268	0.0040	0.0040	0.0041	0.0046	0.0055		
	N	28	28	27	27	28	28	27	27	28		

* p<0.05 by Dunnett's test
 ** p<0.01

006597

Z 4

28

Pathology F₁ Male and Female Adults

Gross pathology was performed on all animals on test (Tables 77, 78 and 155). Histopathology included lesions found in organs of all groups when seen at necropsy. Reproductive organ histopathology included all animals in the control and the 250 ppm treated groups. No compound related effect was evident.

Pathology of F_{2b} Pups

Gross pathology and limited histopathology was performed for 10 of each sex per group in controls and 250 ppm dose. No compound-related effect was apparent.

Final body weights were as follows.

Dose (ppm)	0	25	75	150	250
Males	46.2 ± 6.8	44.2 ± 5.5	43.2 ± 5.8	37.5 ± 7.2**	35.9 ± 8.5**
(g ± SD)					
Females	43.6 ± 2.3	42.7 ± 4.3	41.4 ± 3.6	38.4 ± 7.7	35.7 ± 6.6**
(g ± SD)					

P < 0.01 by Dunnett's test.

This indicates generally dose-related decreases in final body weights of the pups at the 2 highest dose levels.

There were also no effects on liver weights or female reproductive organs. There was a decrease in absolute and relative testes weight at the 150 and 250 ppm dose groups.

Summary and Evaluation

Doses selected in the present two-generation reproduction rat study were 0, 25, 75, 150, and 250 ppm in the diet which came to an average of 0, 1.8, 5.3, 11.1, and 18.5 mg/kg/day, respectively, for the 5 dose levels in nonpregnant animals and during gestation. These were about twice as high in lactating females; i.e., an average of 0, 3.8, 11.2, 23.0, and 37.1 mg/kg/day, respectively. F₀ and F₁ parents were treated for at least 70 to 72 days prior to the first pairing. Histopathology was performed only on control and high-dose-treated F₀ and F₁ adults but only 10 of each sex of control and high-dose-treated F_{2b} pups at weaning. Histopathology was limited to reproductive organs of both sexes, but also included areas of grossly observed lesions in organs of animals of all treated groups. Organ weights included liver and reproductive organs.

Manifestations of toxicity to male and female F₀ and F₁ parents were seen as dose-related decreased body weight gain and decreased food intake at 75, 150, and 250 ppm doses. In addition, final body weights at the time of terminal sacrifice was lower in the 25 ppm treated male (n.s.) and female (p < 0.01)

F₁ parents. There was a tendency for decreased liver weights particularly at the 3 highest dose levels. No effects on gross pathology or histopathology due to treatment were evident in the F₀ and F₁ males or females.

In both the F_{1a} and F_{1b} generations there was a 35% decrease in number of paired females that delivered a litter in the 150 ppm treated group. Such an effect was not seen for either generation of the 250 ppm treated group nor for any treated group of the F_{2a} and F_{2b} generations. The lower fertility in the 150 ppm group was considered a "random occurrence" by the investigator but since it occurred in two successive matings we should not ignore the possibility of an effect.

For the "Male Mating Index" and "Female Mating Index", data for "both mating periods"; i.e. F_{1a} with F_{1b} and F_{2a} with F_{2b}, were combined. These data should be recalculated and presented separately for each mating period. In its present form, the data appear misleading and are higher than they should really be.

There is also the question on the reliability of the data on "Females with Evidence of Mating". Vaginal sperm or copulation plugs were missed for a number of animals on test. It also appears likely that a similar number of animals were considered to have mated but did not, particularly if evidence of mating was based on copulation plugs under the cages rather than vaginal sperm.

In the 4 tables on pathology of the pups (Tables 117, 118, 152 and 153), there are footnotes "Does not include pups found missing or cannibalized". In the F_{1a} generation, dams 23,703 and 23,748 each lost all 8 of their pups between days 7-14 of lactation but on 2 or 3 or them are accounted for with each of the 2 litters. The applicant should account for all missing pups.

There were slight decreases in gestation length in all 4 treated groups if the F_{2a} litters ($P < 0.05-0.01$; 22.1 days for controls 21.3 to 21.6 days for treated). The latter effect is also difficult to explain but it did not occur in the F_{2b}, F_{1a} or F_{1b} generations. It was not accompanied by dystocia or other pregnancy irregularity and was considered not a compound effect.

There were obviously no compound related decreases in live litters at birth, no decreases in live litter size or increase in dead pups at birth, no effect on sex ratio or on Gestation Index (live litter births/total pregnancies). Gestation Index was slightly decreased in all 4 treated groups of the F_{1a} generation, but since it did not occur with any other generation, it was considered sporadic by the applicant.

Viability of pups during lactation was decreased ($p = 0.05$ to 0.01) as early as days 1 and 4 of lactation in the 150 and 250 ppm treated groups of the F_{2a} generation. There was a decrease in pup viability in the 150 and 250 ppm dosed groups of the F_{1a} generation, which became evident by day 14 of lactation. However, dose-related decreased pup weight by day 21 of lactation was evident in the 75, 150, and 250 ppm treated groups in the F_{1a} and F_{2b} generations ($p < 0.01$). Decreased pup weight may be a clearer indication of toxicity to the pups than decrease in viability index.

Results of necropsies of pups found dead are given. Although anomalies were found in a few pups, all of which were subjected to necropsy if not autolyzed, there is no indication of increased incidence of anomalies with dosing of compound. A large number of pups in the 250 ppm dose treated group of the F_{1a} generations prior to discarding them were observed to be small in size, but otherwise "not remarkable." In the F_{2a} generation, pups were visually observed to be small in the 75, 150, and 250 ppm treated groups prior to discarding after weaning, but otherwise "not remarkable."

Ten pups of each sex per group in the F_{2b} generation were examined by gross and microscopic pathology at time of weaning. No compound effect was noted. Final body weights were lower in the pups from the 150 and 250 ppm treated male and female groups. No effect was seen on liver weights of pups, but a decrease was noted in testes weight in the 150 and 250 ppm groups based both on absolute and relative weights, which could suggest retarded testicular development of the pups in the 2 highest dose groups. However, there was obviously no impairment in reproduction function in males based on reproduction indices of F₁ males.

At the terminal killing of adults, parent body weights were decreased for F₀ males and females at the 3 highest doses and in F₁ males and females at all 4 doses, dose-related. There were no effects on relative liver weights of either sex nor on absolute testis or ovarian weights. No effects on gross pathology performed on organs of all animals on test was observed. A histopathology report, signed by Robert G. Geil, D.V.M., Diplomate, American College of Veterinary Pathologists, dated January 19, 1987, indicates that no microscopic lesions related to treatment were seen in F₀ and F₁ parental rats or F_{2b} weanling rats from the 250 ppm group. Histopathology was limited to organs of the reproductive tract of both the males and females of controls and 250 ppm group and lesions of other organs in animals of all groups.

In conclusion, systemic toxicity was observed in the present two-generation rat study with Bladex at 25 ppm in the diet (LDT), based on decreased body weight of males (n.s.) and females

006597

($p < 0.01$) of F_1 adults at the time of terminal killing, around week 57 of the study and at various time periods throughout the study. This dosage is equivalent to about 1.8 mg/kg/day based on food intake. Thus, 1.8 mg/kg/day (LDT) is considered the LEL for systemic toxicity in rats due to long-term oral intake of Bladex.

The LEL for Reproductive Toxicity is considered to be 11,2 mg/kg/day based on pup viability indices and decreased body weight gain of pups during lactation with a diet containing 75 ppm Bladex or higher. The NOEL is 3.8 mg/kg/day (25 ppm).