

US EPA ARCHIVE DOCUMENT



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OK

OFFICE OF  
PESTICIDES AND TOXIC SUBSTANCES

FEB 6 1981

MEMORANDUM

16-2440

SUBJECT: Glyphosate; PP#~~107240~~; EUP and temporary tolerance for Glyphosate in or on cottonseed at 15 ppm. CASWELL#661A; Accession#099720

FROM: William Dykstra, Toxicologist  
Toxicology Branch, HED (TS-769)

WSD JDC  
2/11/81

TO: Robert Taylor (25)  
Registration Division (TS-767)  
and  
Residue Chemistry Branch  
Hazard Evaluation Division (TS-769)

WJB

Recommendations:

- 1) The requested tolerance can be toxicologically supported.
  - A. EUP Program: A total of 2688 lbs of Roundup is to be used on 1280 acres in eight states. A two-year testing program is proposed, due to the cropping systems involved and the long-term efficacy determinations that need to be made.
  - B. Formulation to be used in Roundup (EPA Reg.#524-308). Inerts are cleared under 180.1001.

Section F: Proposed Temporary Tolerance

Request is made to establish temporary tolerances for the combined residues of the herbicide N-phosphono-methylglycine (Glyphosate) and its metabolite, aminomethylphosphonic acid, in or on the raw agricultural commodity as follows:

cottonseed ----- 15.0 ppm

Review:

1) Memo of 8/22/78 from R. Engler to R. Taylor. Toxicology Branch has reviewed the validated studies in support of Glyphosate.

a) Data considered

- °Oral LD<sub>50</sub> (rabbit): 3.8 gm/kg (valid)
- °90-Day Rat Feeding: NOEL = 2000 ppm (valid)
- °90-Day Dog Feeding: NOEL = 2000 ppm (valid)
- °Teratology (2 studies) Rabbit: Negative at 30 mg/kg/day (highest dose); repeat studies with a higher dose.
- °2-Year Dog Feeding: NOEL = 300 ppm (valid)
- °3-Generation Rat Reproduction: NOEL = 100 ppm (valid)
- °18-Month Mouse Feeding: No carcinogenic potential at 300 ppm (highest dose). Study must be repeated since too many animals are missing.
- °2-Year Rat Feeding: NOEL = 100 ppm (valid) Study is adequate to determine the toxic effects, but only marginal with respect to oncogenic evaluation since too few animals examined. As reported the study shows no oncogenic potential.
- °Neurotoxicity (hen): Negative at 7.5 gm/kg (cumulative for 3 days) (valid)
- °Dominant Lethal (mice): Negative at 10 mg/kg (highest dose), supplemental study, no records of positive control.
- °Host-Mediated Assay: Negative (supplemental study) no raw data available.
- °Rec-Assay: Negative (supplemental study) no raw data available.

2) Memo of 9/22/79 from M.L. Alexander to Product Manager #25. Glyphosate was not mutagenic in the following test systems:

- a) Rec-assay in two strains of B. subtilis up to 2000 ug test material/disk.
- b) Reverse mutation in five histidine-requiring strains of S. typhimurium and one tryptophan-requiring strain of E. coli with or without metabolic activation.
- c) Ames test in four strains of Salmonella, with or without metabolic activation.

- 3) Memo of 1/16/81 from W. Dykstra to R. Taylor.
  - a) Rat Teratology: Severe maternal toxicity at 3500 mg/kg/day;  
negative at 3500 mg/kg/day.  
Fetotoxic NOEL = 1000 mg/kg/day
  - b) Rabbit Teratology: Negative at 350 mg/kg/day  
Fetotoxic NOEL = 175 mg/kg/day
  - c) Mouse Dominant Lethal: Negative at 2000 mg/kg
- 4) No new toxicity data were submitted with this petition.
- 5) Evaluation of the ADI.

The ADI is based on the NOEL of 100 ppm (5 mg/kg/day) in a 2-year rat feeding study. This is the most sensitive species for which chronic toxicity data are available.

$$\text{ADI} = \text{NOEL} \times \frac{1}{100}$$
$$\text{ADI} = 5 \text{ mg/kg/day} \times \frac{1}{100} = 0.05 \text{ mg/kg/day}$$

The MPI for a 60 kg person is 3 mg/day

- 6) Tolerances have been established under 40 CFR 180.364.
- 7) No regulatory actions are pending against the pesticide.
- 8) The published tolerances utilize 7.21% of the ADI. Unpublished, Tox approved tolerances utilize the ADI to 19.05%.

The current action utilizes 1.12% of the ADI. All tolerances on Glyphosate utilize 20.17% of the ADI. (computer printout attached).

#### Conclusions and Recommendations:

The requested tolerance can be toxicologically supported.

The oncogenic potential of Glyphosate is not fully elucidated. The chronic rat and mouse feeding studies, however, provide assurance that Glyphosate has a relatively low oncogenic potential. A further assurance of low risk with Glyphosate is found in the fact that on a theoretical basis the exposure via the diet is about one-fifth of the ADI at present.