

US EPA ARCHIVE DOCUMENT

TEXT SEARCHABLE DOCUMENT

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

Data Requirement:	PMRA Data Code	{.....}
	EPA DP Barcode	Not Provided
	OECD Data Point	{.....}
	EPA MRID	43334710
	EPA Guideline	OPPTS 850.2200 (71-2a)

Test material: Aminomethyl Phosphonic Acid (Glyphosate Degradate) **Purity:** 87.8%

Common name: AMPA

Chemical name: IUPAC: Not Reported

CAS name: Not Reported

CAS No. Not Reported

Synonyms: None Reported

Primary Reviewer: John Marton
Staff Scientist, Cambridge Environmental Inc.

Signature:

Date: 1/12/07

Secondary Reviewer: Teri S. Myers
Senior Scientist, Cambridge Environmental Inc.

Signature:

Date: 2/12/07

Primary Reviewer: Stephen Carey
EPA Biologist, OPP, EFED, ERBIII

Date: 7/18/07

Secondary Reviewer(s): {.....}
{EPA/OECD/PMRA}

Date: {.....}

Reference/Submission No.: {.....}

Company Code {.....} [For PMRA]

Active Code {.....} [For PMRA]

Use Site Category: {.....} [For PMRA]

EPA PC Code 417300 (Parent Compound) & 207800 (Degradate Compound)

Date Evaluation Completed: July 18 2007

CITATION: Long, R.D., G.J. Smith, J.B. Beavers and S.P. Lynn. 1994. AMPA: A Dietary LC₅₀ Study with the Northern Bobwhite. Unpublished study performed by Wildlife International Ltd., Easton, MD. Laboratory report number 139-275. Study sponsored by Monsanto Agricultural Company, St. Louis, Missouri. Study completed October 28, 1991.

DISCLAIMER: This document provides guidance for EPA and PMRA reviewers on how to complete a data evaluation record after reviewing a scientific study concerning the acute dietary toxicity of a pesticide to avian species. It is not intended to prescribe conditions to any external party for conducting this study nor to establish absolute criteria regarding the assessment of whether the study is scientifically sound and whether the study satisfies any applicable data requirements. Reviewers are expected to review and to determine for each study, on a case-by-case basis, whether it is scientifically sound and provides sufficient information to satisfy applicable data requirements. Studies that fail to meet any of the conditions may be accepted, if appropriate; similarly, studies that meet all of the conditions may be rejected, if appropriate. In sum, the reviewer is to take into account the totality of factors related to the test methodology and results in determining the acceptability of the study.

US EPA ARCHIVE DOCUMENT



2080479

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

EXECUTIVE SUMMARY:

The subacute dietary toxicity of AMPA (Glyphosate Degradate) to 10-d-old Northern Bobwhite Quail (*Colinus virginianus*) was assessed over 8 days. AMPA was administered to the birds in the diet at 0 (vehicle control), 493, 878, 1563, 2774 and 4934 mg ai/kg dw of diet; verification of these dose levels was provided in MRID 43334712 and showed that all values were within 93-109% of target (see Reviewer's Comments section for details). The 8-day acute dietary LC₅₀ was >4934 mg ai/kg diet. The 8-day NOAEC of AMPA based on the lack of treatment-related mortality and sub-lethal effects was 4934 mg ai/kg diet. According to the US EPA classification, AMPA (Glyphosate Degradate) would be classified as practically non-toxic to Northern bobwhite quail on a subacute dietary basis at a nominal concentration of 4934 mg ai/kg diet.

One mortality was observed on Day 2 in the 493 mg ai/kg diet treatment group; however, this was not considered to be treatment related. Furthermore, the test material did not significantly impact weight gain or food consumption.

This toxicity study is classified as scientifically sound and does satisfy the guideline requirement for a subacute dietary toxicity study with Northern bobwhite quail.

Results Synopsis

Test Organism Size/Age(Mean Weight): 10 Days; 20 (18-22) g

LC ₅₀ :	>4934 mg ai/kg diet	95% C.I.:	N/A
Probit slope:	N/A	95% C.I.:	N/A
NOAEC:	4934 mg ai/kg diet		
Endpoint(s) affected:	None		

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

I. MATERIALS AND METHODS:

GUIDELINE FOLLOWED: This study was conducted following guidelines outlined in Section 71-2 of the Environmental Protection Agency Registration Guidelines, *Pesticide Assessment Guidelines, FIFRA Subdivision E, Hazard Evaluation, Wildlife and Aquatic Organisms*; and upon ASTM Standard E857-87, "Standard Practice for Conducting Subacute Dietary Toxicity Tests with Avian Species." The following deviation from OPPTS 850.2200 was noted:

The physiochemical properties of the test material were not reported.

This deviation did not impact the acceptability of the study.

COMPLIANCE: Signed and dated No Data Confidentiality, GLP and Quality Assurance statements were provided. This study was conducted in compliance with GLP standards as published by the U.S. EPA in 40 CFR, Part 160; OECD, ISBN 92-84-12367-9; and Japan MAFF, 59 NohSan, Notification No. 3850, Agricultural Production Bureau.

A. MATERIALS:

1. Test Material Aminomethyl Phosphonic Acid (Glyphosate Degradate)

Description: White Powder

Lot No./Batch No. : PIT-9008-2407T

Purity: 87.8%

Stability of Compound Under Test Conditions: Samples of the test diets were taken to verify the test concentrations administered and to confirm the stability and homogeneity of the test substance in the diets. Samples were frozen and transferred to Monsanto Environmental Health Laboratories for analysis. The results of this analysis are provided in a separate report, MRID 43334712. They revealed that the test material levels taken on days 0 and 5 ranged from 88-105% of target.

Storage Conditions of Test Chemicals: Stored at room temperature.

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

Physicochemical properties of AMPA.

Parameter	Values	Comments
Water solubility at 20EC	Not Reported	
Vapor pressure	Not Reported	
UV absorption	Not Reported	
pKa	Not Reported	
Kow	Not Reported	

(OECD recommends water solubility, stability in water and light, pKa, Pow, and vapor pressure of test compound)

2. Test organism:

Species (common and scientific names): Northern bobwhite quail (*Colinus virginianus*)
(EPA recommends using either bobwhite quail or mallard duck.)

Age at study initiation: 10 Days
(EPA recommends: 10-14 days old)

Weight at study initiation (mean and range): 20 (18-22) g

Source: On-site Production Flock.

B. STUDY DESIGN:

1. Experimental Conditions

- a. Range-finding Study: No range-finding studies were reported.
- b. Definitive Study:

Table 1: Experimental Parameters

Parameter	Details	Remarks
		Criteria
<u>Acclimation</u> Period: Conditions: (same as test or not) Feeding: Health: (any mortality observed)	Continuous Same as test Game bird ration formulated to Wildlife International Ltd.'s specifications Not reported; however, birds exhibiting abnormal behavior or physical injury were not used in the definitive test.	Food and water from the town of Easton public water supply were provided <i>ad libitum</i> .

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

Parameter	Details	Remarks
		Criteria
Pen size and construction materials	Thermostatically controlled brooding pens had floors that measured approximately 72 x 90 cm and ceiling height was approximately 23 cm. External walls, ceilings and floors were constructed of galvanized steel wire and sheeting.	<i>Recommended pen size is about 35 x 100 x 24 cm</i>
Test duration	5 days with treated feed and 3 days with untreated feed	<i>Recommended test duration is 5 days with treated feed and at least 3 days observation with "clean" feed.</i>
<u>Test concentrations</u> nominal: measured:	0 (vehicle control), 493, 878, 1563, 2774 and 4934 mg ai/kg dw of diet Measured samples of dose levels showed them to be 93-109% of target.	The reviewer corrected the nominal concentrations for the purity of the test material (87.8%). The results from the analysis of the test material in the feed were provided in MRID 43334712. <i>Five or six test concentrations should be used in a geometric scale, unless the LC₅₀ > 5000 mg ai/kg diet.</i>
<u>Solvent/vehicle, if used</u> type: amount:	Corn Oil 2%	<i>Recommended solvents include distilled water, corn oil, propylene glycol, 1% carboxymethylcellulose, or gum arabic. The solvent should not be more than 2%.</i>
Diet preparation and feeding	Test diets were prepared by mixing the test substance into the diet with corn oil. Mixing was done with a Hobart mixer. Diets were prepared on the day of test initiation and sufficient feed was prepared for the duration of the treated feed period.	The control was treated with 2% corn oil only. <i>The control group should be tested with a diet containing the maximum amount of vehicle used in treated diets.</i>
Feed withholding period	None reported	
Stability and homogeneity of test material in the diet determined (Yes/No)	Samples were shipped to Monsanto Environmental Health Laboratories for analysis; the results of these analyses are described in another study report, MRID 43334712 (see Reviewer's Comments section for	

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

Parameter	Details	Remarks
		Criteria
	details).	
<u>Number of birds per replicate/groups</u> for negative control: for vehicle control: for treated:	N/A 10 10	<i>The recommended number of birds per replicate is a minimum of ten.</i>
<u>Number of replicates/group (if used)</u> for negative control: for vehicle control: for treated:	N/A 3 1	
<u>Test conditions</u> temperature: relative humidity(%): photoperiod:	Brooding Compartment 35±2°C Room 25±2°C 41±1% 16L:8D; 130 lux	<i>Recommended brooder temperature is about 35°C (95°F)</i> <i>Recommended room temperature is 22-27°C (71-81°F)</i> <i>Recommended relative humidity is 30-80%</i> <i>Recommended photoperiod is a minimum of 14 hours of light.</i>
Reference chemical, if used	N/A; a reference chemical was not used	

2. Observations:

Table 2: Observations

Parameters	Details	Remarks
Parameters measured (mortality/body weight/ mean feed consumption/ others)	-Mortality -Average Weight Gain -Feed Consumption	Feed consumption was reported as an estimate due to the unavoidable wastage of the by the birds.
Indicate the stability and homogeneity of test chemical in the diet	Not reported	
Indicate if the test material was regurgitated	No regurgitation was reported	
Treatments on which necropsies were performed	No necropsies were performed	
Observation intervals	Daily	
Were raw data included?	Sufficient summarized data tables	

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

Parameters	Details	Remarks
	were provided.	

II. RESULTS AND DISCUSSION:

A. MORTALITY:

Throughout the duration of the test, a single mortality occurred in the nominal 493 mg ai/kg diet treatment group on Day 2. This one mortality was not attributed to the test material. Therefore, the resulting NOAEC and LC₅₀ values were 4934 and >4934 mg ai/kg diet, respectively.

Table 3: Effect of AMPA on Mortality of *Colinus virginianus*

Treatment (mg ai/kg diet)	No. of Birds	Cumulative Mortality					
		Day 1	Day 2	Day 3	Day 5	Day 8	
Vehicle Control	30	0	0	0	0	0	
493	10	0	1	1	1	1	
878	10	0	0	0	0	0	
1563	10	0	0	0	0	0	
2774	10	0	0	0	0	0	
4934	10	0	0	0	0	0	
NOAEC	4934 mg ai/kg diet						
LC ₅₀	>4934 mg ai/kg diet						
Reference chemical	mortality	N/A	N/A	N/A	N/A	N/A	N/A
	LC ₅₀	N/A					
	NOAEC	N/A					

N/A- Not Applicable

B. SUB-LETHAL TOXICITY ENDPOINTS:

No differences in food consumption of body weight gain were noted at any treatment level. Furthermore, no behavioral abnormalities were noted. The resulting NOAEC and EC₅₀ values were 4934 and >4934 mg ai/kg diet, respectively.

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

Table 4: Sub-lethal Effect of AMPA on *Colinus virginianus*

Treatment (mg ai/kg diet)	Observation						
	Body Weight (g)			Food Consumption (g/bird/day)			
	Day 0	Day 5	Day 8	Days 0-5	Days 6-8		
Vehicle Control	18	30	39	8	10		
493	20	30	39	8	10		
878	20	32	42	8	11		
1563	22	35	45	9	10		
2774	21	34	43	7	8		
4934	19	33	41	10	9		
NOAEC	4934 mg ai/kg diet						
EC ₅₀	>4934 mg ai/kg diet						
Reference chemical	effect	N/A	N/A	N/A	N/A	N/A	N/A
	NOAEL	N/A					
	LC ₅₀	N/A					

N/A- Not Applicable

C. REPORTED STATISTICS:

All toxicity values were determined by visual inspection of the data.

D. VERIFICATION OF STATISTICAL RESULTS:

Statistical Method(s): The lack of treatment-related mortality precluded the statistical analysis of the data. As values for weight gain and food consumption at all treatment levels were similar to the control values, the reviewer visually determined the toxicity values. All values were determined based on the nominal concentrations which the reviewer corrected for the purity of the test material (87.8%).

LC₅₀: >5620 mg ai/kg diet 95% C.I.: N/A
 NOAEC: 5620 mg ai/kg diet
 Probit Slope: N/A 95% C.I.: N/A

Adjusted for active ingredient: (Optional if over 80% ai)

LC₅₀: >4934 mg ai/kg diet 95% C.I.: N/A
 NOAEC: 4934 mg ai/kg diet
 Probit Slope: N/A 95% C.I.: N/A

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

E. STUDY DEFICIENCIES:

There were no study deficiencies.

F. REVIEWER'S COMMENTS:

The reviewer's results were identical to those of the study author.

Samples of the test diets were taken to verify the test concentrations administered and to confirm the stability and homogeneity of the test substance in the diets. Samples were frozen upon transfer to and for storage at Monsanto Environmental Health Laboratories. Results of these analyses are described in a separate report:

MRID 43334712. Schermes, S. and K.W. McCune. (1994). Results of the Analyses of Avian Diet Samples for AMPA (aminomethyl phosphonic acid). Guideline 71-2. Submitted by The Agricultural Group of Monsanto Company, St. Louis, MO.

Data from this report showed that mixing was uniform (coefficient of variations <3%), as determined by high and low level mixtures of avian diet and test material. Stability of AMPA in avian diet over 5 days at room temperature was also shown (day 0 and 5 samples ranged from 88-105% of target). Adequate diet homogeneity was observed and concentrations of AMPA in avian diet were shown to accurately reflect target levels (93-109%). The analytical report additionally noted a positive diet interference problem (resulting in higher recoveries), where co-extracted amino acids from the diet matrix were possibly reacting with the derivitization agent used in the LC-UV process. To minimize this interference, some sample recovery values were corrected using the concurrently analyzed mean QC sample recoveries.

The in-life portion of the definitive toxicity test was conducted from September 6 to September 14, 1990.

G. CONCLUSIONS:

This study is scientifically sound and does satisfy the guideline requirements for a subacute dietary toxicity test with the Northern bobwhite quail. The NOAEC and LC₅₀ values were 4934 and >4934 mg ai/kg diet, respectively.

LC₅₀: >4934 mg ai/kg diet 95% C.I.: N/A
NOAEC: 4934 mg ai/kg diet
Endpoint(s) affected: None

III. REFERENCES:

Environmental Protection Agency. 1982. Pesticide Assessment Guidelines, FIFRA Subdivision E, Hazard Evaluation: Wildlife and Aquatic Organisms, Subsection 71-2. Office of Pesticide Programs. Washington, DC. 86 pp.

ASTM Standard E857-81. 1982 "Standard Practice for Conducting Subacute Dietary Toxicity Tests with Avian Species". American Society for Testing and Materials.

National Institutes of Health. 1985. Guide for the care and use of laboratory animals. NIH Pub. No. 85-23. 83 pp.

Stephan, C.E. 1978. U.S. EPA, Environmental Research Laboratory, Duluth, Minnesota. Personal Communication.

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the Acute Dietary Toxicity of AMPA to Avian Species, *Colinus virginianus*

PMRA Submission Number {.....}

EPA MRID Number 43334710

Finney, D.J. 1971. Statistical Methods in Biological Assay, 2nd edition, Griffin Press, London.

Thompson, W.R. 1947. Bacteriological Reviews. Vol II, 2:115-145.

Stephan, C.E. 1977. Methods for Calculating an LC50. Aquatic Toxicology and Hazard Evaluations, Amer. Soc. Test Mat., Pub. No. STP 634: 65-84.