

US EPA ARCHIVE DOCUMENT



OFFICIAL RECORD  
HEALTH EFFECTS DIVISION  
SCIENTIFIC DATA REVIEWS  
EPA SERIES 361

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES, AND  
TOXIC SUBSTANCES

TXR No. 0054586

MEMORANDUM

DATE: May 18, 2007

SUBJECT: **Pyroxsulam:** Qualitative Risk Assessment Based On CD-1  
(CrI:CD1(ICR)) Mouse Carcinogenicity Dietary Study

P.C. Code: 108702

TO: Kimberly Harper, Toxicologist  
Registration Action Branch 2  
Health Effects Division (7509P)

FROM: Lori L. Brunsman, Statistician  
Science Information Management Branch  
Health Effects Division (7509P)

THROUGH: Jessica Kidwell, EPS   
and  
Jess Rowland, Branch Chief   
Science Information Management Branch  
Health Effects Division (7509P)

BACKGROUND

A carcinogenicity study in CD-1 (CrI:CD1(ICR)) mice was conducted by Dow Chemical Company, Midland, Michigan, for Dow AgroSciences, Indianapolis, Indiana, and dated December 15, 2005 (Study ID No. 031015, MRID No. 46908406).

The study design allocated groups of 50 mice per sex to nominal dose levels of 0, 10, 100 and 1000 mg/kg/day of Pyroxsulam in the diet for 79 weeks. Actual doses were 0, 10, 100 and 932 mg/kg/day for males. There were no compound-related tumors in the females so only analyses of the males are presented in this document.

MAY 22 REC'D  
2007  
R/RC  
mcs

US EPA ARCHIVE DOCUMENT

## ANALYSES

### **Survival Analyses**

There were no statistically significant incremental changes in mortality with increasing doses of Pyroxsulam in male mice (Table 1).

### **Tumor Analyses**

Male mice had a statistically significant trend for liver carcinomas at  $p < 0.05$ . There were statistically significant pair-wise comparisons of the 10 and 1000 mg/kg/day dose groups with the controls for liver adenomas, and liver adenomas and/or carcinomas combined, all at  $p < 0.05$ . The statistical analyses of the tumors in the male mice were based upon Fisher's Exact Test for pair-wise comparisons and the Exact Test for trend (Table 2)

Table 1. Pyroxsulam – CD-1 (CrI:CD1(ICR)) Mouse Study (MRID 46908406)

Male Mortality Rates<sup>e</sup> and Cox or Generalized K/W Test Results

Dose (mg/kg/day)	Weeks			Total
	1-26	27-52	53-79 <sup>f</sup>	
0	0/50	1/50	9/49	10/50 (20)
10	1/50	3/49	6/46	10/50 (20)
100	0/50	2/50	8/48	10/50 (20)
1000	0/50	2/50	10/48	12/50 (24)

<sup>e</sup>Number of animals that died during interval/Number of animals alive at the beginning of the interval.

<sup>f</sup>Final sacrifice at week 79.

( )Percent.

Note: Time intervals were selected for display purposes only.  
 Significance of trend denoted at control.  
 Significance of pair-wise comparison with control denoted at dose level.  
 If \*, then  $p < 0.05$ . If \*\*, then  $p < 0.01$ .

Table 2. Pyroxsulam – CD-1 (CrI:CD1(ICR)) Mouse Study (MRID 46908406)

Male Liver Tumor Rates<sup>†</sup>  
and Fisher's Exact Test and Exact Test for Trend Results

	Dose (mg/kg/day)			
	0	10	100	1000
Adenomas (%)	5/49 (10)	13/46 (28)	9 <sup>a</sup> /49 (18)	14/48 (29)
p =	0.06696	0.02300*	0.19363	0.01716*
Carcinomas (%)	1/49 (2)	0/46 (0)	2 <sup>b</sup> /49 (4)	4/48 (8)
p =	0.02622*	1.00000	0.50000	0.17451
Combined (%)	6/49 (12)	13/46 (28)	10 <sup>c</sup> /49 (20)	15 <sup>d</sup> /48 (31)
p =	0.05737	0.04462*	0.20651	0.02067*

<sup>†</sup>Number of tumor bearing animals/Number of animals examined, excluding those that died before week 52.

<sup>a</sup>First adenoma observed at week 53, dose 100 mg/kg/day.

<sup>b</sup>First carcinoma observed at week 73, dose 100 mg/kg/day.

<sup>c</sup>One animal in the 100 mg/kg/day dose group had both an adenoma and a carcinoma.

<sup>d</sup>Three animals in the 1000 mg/kg/day dose group had both an adenoma and a carcinoma.

Note: Significance of trend denoted at control.  
Significance of pair-wise comparison with control denoted at dose level.  
If \*, then p < 0.05. If \*\*, then p < 0.01.

## References

Cox, D.R. (1972) Regression Models and Life Tables (with discussion). J. Royal Stat. Soc. Ser. B. 34, 187-220.

Gart, J.J., D. Krewski, P.N. Lee, R.E. Tarone, and J. Wahrendorf (1986) The Design and Analysis of Long-Term Animal Experiments. In: Statistical Methods in Cancer Research, Volume III. IARC Scientific Publications No. 79. Lyon, France: International Agency for Research on Cancer, p. 18.

Peto, R., M. Pike, N. Day, R. Gray, P. Lee, S. Parish, J. Peto, S. Richard, and J. Wahrendorf (1980) Guidelines for Simple, Sensitive, Significant Tests for Carcinogenic Effects in Long-Term Animal Experiments. In: Monographs on the long-term and short-term screening assays for carcinogens: a critical appraisal. IARC Monographs, Supplement 2. Lyon, France: International Agency for Research on Cancer, pp. 311-426.

Thomas, D.G., N. Breslow, and J.J. Gart (1977) Trend and Homogeneity Analyses of Proportions and Life Table Data. Computers and Biomedical Research 10, 373-381



13544

# R144885

Chemical: Benzene hexachloride gamma isomer of ----- % other  
isomers of ----- %  
Lindane

PC Code:

008901

009001

HED File Code: 13100 Other Tox Documents

Memo Date: 2/26/1970

File ID: 00000000

Accession #: 000-00-0118

HED Records Reference Center

5/16/2007