

US EPA ARCHIVE DOCUMENT


Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID Number ⁴⁶⁹⁰⁸⁴⁻³⁴ {.....}

Data Requirement: PMRA DATA CODE: 9.8.5
 EPA DP Barcode:
 OECD Test Guideline: IIA 8.6
 EPA Guideline: 123-2 (OPPTS 850.4400 (Draft April 1996))

Test material: ADTP Metabolite of pyroxsulam **Purity (%):** 98%
Common name: ADTP Metabolite of XDE-742
Chemical name: [1,2,4]triazolo[1,5-a]pyrimidin-2-amine,5,7-dimethoxy
ID No: TSN 103822
Lot No. 104425
IUPAC name: 5,7-dimethoxy[1,2,4]triazolo[1,5-a]pyrimidin-2-amine
CAS Name: [1,2,4]triazolo[1,5-a]pyrimidin-2-amine, 5,7-dimethoxy
CAS No.: Not available
Synonyms: ADTP
Chemical structure



Primary Reviewer: Chris Lee-Steere **Date:** 5 July 2007 *22/08/07*
 Australian Government Department of the Environment, Water, Heritage and the Arts

Secondary Reviewers: Jack Holland **Date:** 23 July 2007 *22/2/08*
 Australian Government Department of the Environment, Water, Heritage and the Arts
 Brian Kiernan **Date:** 22 August 2007 *7/10/08*

US Environmental Protection Agency
PMRA Reviewer: Émilie Larivière **Date:** 30 July 2007 *05/03/08*
Environmental Assessment Directorate, PMRA

Reference/Submission No.: APVMA ATS 40362 NCRIS 61286

Company Code: DWE
Active Code: JUA
Use Site Category: 13, 14
EPA PC Code:



US EPA ARCHIVE DOCUMENT

**Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID Number{.....}

CITATION: Hoberg, James R. (2006): ADTP Metabolite of XDE-742: Toxicity to Duckweed, *Lemna gibba*. Springborn Smithers Laboratories, 790 Main Street, Wareham, Massachusetts, Study No. 12550.6415. Dow AgroSciences, Report No. 050123, 3 April 2006. Unpublished.


Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID Number ⁴⁶⁷⁰⁸⁴⁻³⁴ {.....}

Data Requirement: PMRA DATA CODE: 9.8.5
EPA DP Barcode:
OECD Test Guideline: IIA 8.6
EPA Guideline: 123-2 (OPPTS 850.4400 (Draft April 1996))

Test material: ADTP Metabolite of pyroxsulam **Purity (%):** 98%
Common name: ADTP Metabolite of XDE-742
Chemical name: [1,2,4]triazolo[1,5-a]pyrimidin-2-amine,5,7-dimethoxy
ID No.: TSN 103822
Lot No.: 104425
IUPAC name: 5,7-dimethoxy[1,2,4]triazolo[1,5-a]pyrimidin-2-amine
CAS Name: [1,2,4]triazolo[1,5-a]pyrimidin-2-amine, 5,7-dimethoxy
CAS No.: Not available
Synonyms: ADTP
Chemical structure



Primary Reviewer: Chris Lee-Steere **Date:** 5 July 2007 *22/08/07*
Australian Government Department of the Environment, Water, Heritage and the Arts

Secondary Reviewers: Jack Holland **Date:** ~~23 July 2007~~ *22/2/08*
Australian Government Department of the Environment, Water, Heritage and the Arts
Brian Kiernan **Date:** 22 August 2007 *7/10/08*

US Environmental Protection Agency

PMRA Reviewer: Émilie Larivière
Environmental Assessment Directorate, PMRA

Date: 30 July 2007 *Emilie Lariviere*
05/03/08

Reference/Submission No.: APVMA ATS 40362 NCRIS 61286

Company Code: DWE
Active Code: JUA
Use Site Category: 13, 14
EPA PC Code:

**Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID Number{.....}

CITATION: Hoberg, James R. (2006): ADTP Metabolite of XDE-742: Toxicity to Duckweed, *Lemna gibba*. Springborn Smithers Laboratories, 790 Main Street, Wareham, Massachusetts, Study No. 12550.6415. Dow AgroSciences, Report No. 050123, 3 April 2006. Unpublished.


Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Data Requirement: PMRA DATA CODE: 9.8.5
EPA DP Barcode:
OECD Test Guideline: IIA 8.6
EPA Guideline: 123-2 (OPPTS 850.4400 (Draft April 1996))

Test material: ADTP Metabolite of pyroxsulam **Purity (%):** 98%
Common name: ADTP Metabolite of XDE-742
Chemical name: [1,2,4]triazolo[1,5-a]pyrimidin-2-amine,5,7-dimethoxy
ID No: TSN 103822
Lot No. 104425
IUPAC name: 5,7-dimethoxy[1,2,4]triazolo[1,5-a]pyrimidin-2-amine
CAS Name: [1,2,4]triazolo[1,5-a]pyrimidin-2-amine, 5,7-dimethoxy
CAS No.: Not available
Synonyms: ADTP
Chemical structure



Primary Reviewer: Chris Lee-Steere **Date:** 5 July 2007
Australian Government Department of the Environment and Water Resources

Secondary Reviewers: Jack Holland **Date:** 23 July 2007
Australian Government Department of the Environment and Water Resources
Brian Kiernan Date: 22 August 2007

US Environmental Protection Agency
PMRA Reviewer: Émilie Larivière **Date:** 30 July 2007
Environmental Assessment Directorate, PMRA

Reference/Submission No.: APVMA ATS 40362 NCRIS 61286

Company Code: DWE
Active Code: JUA
Use Site Category: 13, 14
EPA PC Code:

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

CITATION: Hoberg, James R. (2006): ADTP Metabolite of XDE-742: Toxicity to Duckweed, *Lemna gibba*. Springborn Smithers Laboratories, 790 Main Street, Wareham, Massachusetts, Study No. 12550.6415. Dow AgroSciences, Report No. 050123, 3 April 2006. Unpublished.

EXECUTIVE SUMMARY:

In a 7-day acute toxicity study, freshwater floating aquatic vascular plants (duckweed, *Lemna gibba*) were exposed to pyroxsulam ADTP metabolite at nominal concentrations of 0, 6.3, 13, 25, 50 and 100 mg ac/L. Mean measured concentrations were 0, 5.7, 12, 23, 46 and 93 mg ac/L. The study was conducted under static renewal conditions at days 3 and 5 in accordance with the guidelines, OECD 221 "Lemna sp. Growth Inhibition Test" (draft, 2002) and US EPA guidelines including U.S. Environmental Protection Agency (1996). Ecological Effects Test Guidelines. OPPTS 850.4400 Aquatic Plant Toxicity Test using Lemna sp., Tiers I and II. Draft April 1996.

The % inhibition was determined for frond number, mean specific growth rate and biomass (frond dry weight). No biological endpoint showed statistically significant inhibition at any concentration tested. Inhibition compared to controls ranged between 6% to -32% (stimulation) for all endpoints. The 7-day NOECs for all endpoints was 93 mg ac/L, the highest rate tested, and the 7-day LOECs for all endpoints was >93 mg ac/L (mean measured concentration). The 7-day EC50s for all endpoints was >93 mg ac/L.

At the end of the study, fronds at all exposure concentrations were observed to be normal.

This toxicity study is classified as acceptable and satisfies the guideline requirement for an acute toxicity study with the aquatic vascular plants *Lemna gibba* (duckweed).

Results Synopsis

Test Organism: Duckweed (*Lemna gibba*)

Test Type (Flow through, Static, Static Renewal): Static renewal

Endpoint	Frond No.	Growth rate	Frond dry weight
NOEC (mg ac/L)	93	93	93
EC05 (mg ac/L) (95% C.I.)	>93	>93	>93
LOEC (mg ac/L)	>93	>93	>93
IC50 or EC50 (mg ac/L) (95% C.I.)	>93	>93	>93

No 95% confidence intervals associated with EC05 and EC50 values as these were determined empirically.

Endpoint(s) Effected: No biological endpoint tested in this study was adversely affected.

I. MATERIALS AND METHODS

GUIDELINE FOLLOWED:

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

OECD, 2000. OECD Guideline for Testing of Chemicals. *Lemna* sp. Growth Inhibition Test. Proposed Guideline #221. Revised Draft, October 2000.

U.S. EPA, 1996. Office of Prevention, Pesticides and Toxic Substances. Ecological Effects Test Guideline, OPPTS 850.4400. Aquatic Plant Toxicity Test Using *Lemna* spp., Tiers I and II. "Public Draft" EPA 712-C-96-156 April 1996. U.S. Environmental Protection Agency. Washington, D.C.

The following protocol deviations are noted in the study report:

The protocol states that the light intensity will range from 6500 to 10,000 lux and the photosynthetically-active radiation (PAR) will range from 85 to 120 $\mu\text{E}/\text{m}^2/\text{s}$. During the definitive test, the light intensity ranged from 6500 to 9100 lux and the PAR ranged from 119 to 141 $\mu\text{E}/\text{m}^2/\text{s}$. Since the light intensity was within the appropriate range, the PAR was not adjusted. This deviation did not impact the results of this study.

The study protocol requires the light intensity of the test area be measured daily. The light intensity was inadvertently not measured on day 1. Light intensity measurements were recorded on all other test days and were consistent within the required range.

These deviations are not expected to influence the study results.

This DER has assessed the study report against the OECD 221 and US EPA OPPTS 850.4400 requirements.

COMPLIANCE: All phases of the study were reported as conducted in compliance with the following Good Laboratory Practice Standards:

- OECD Series on Principles of Good Laboratory Practice and Compliance Monitoring, Number 1. OECD Principles on Good Laboratory Practice (as revised in 1997) ENVIMCICHEM (98) 17; and
- U.S. Environmental Protection Agency - FIFRA GLPs, Title 40 CFR, Part 160- Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), Good Laboratory Practice Standards, Final Rule.

Signed and dated Compliance with Good Laboratory Practice Standards, Quality Assurance and No Data Confidentiality Claims statements were provided.

A. MATERIALS:

1. Test Material

ADTP Metabolite of pyroxsulam

Description:

Solid

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Lot No./Batch No.: 104425

Purity: 98% (Certificate of analysis provided)

Stability of Compound Under Test Conditions:

At the beginning and end of two renewal periods (i.e., days 0 and 3 and days 5 and 7), one sample was removed from each test solution and the control and analyzed for ADTP metabolite of pyroxsulam. Samples analyzed from newly prepared solutions were removed from the volumetric flasks prior to division into the replicate vessels. Test solution samples analyzed at the end of the renewal periods (days 3 and 7) were removed from composited solutions of each treatment level and the control. The following results were found:

Table 1: Measured concentrations of ADTP Metabolite of pyroxsulam

Nominal Concentration (mg ac/L)	Measured Concentration (mg ac/L)					% of Nominal
	0 hour (new)	Day 3 (aged)	Day 5 (new)	Day 7 (aged)	Mean (SD)	
Control	<0.20	<0.22	<0.22	<0.22	Not applicable	Not applicable
6.3	6.1	5.9	5.3	5.4	5.7	90
13	12	12	11	12	12	91
25	23	23	23	23	23	92
50	48	48	45	45	46	93
100	94	95	90	93	93	93

There are no stability data for the test substance under light.

Storage conditions of test chemicals:

Stored at in the original container in a dark ventilated cabinet.

Physicochemical properties of ADTP metabolite of pyroxsulam:

None available at the time of testing.

2. Test organism:

Name: Duckweed (*Lemna gibba*)

Strain, if provided: 310

Source: Obtained from the University of Toronto, Toronto, Canada and maintained in stock culture at Springborn Smithers Laboratories, Wareham, MA.

Method of cultivation: Stock cultures were grown in 270-mL covered crystallizing dishes containing 100 mL of medium. The cultures were maintained in an

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

environmental chamber within the following conditions: a temperature of $24 \pm 2^\circ\text{C}$ and continuous illumination of approximately 600 to 930 footcandles (6500 to 10,000 lux). Lighting was supplied by Premira VitaLux® fluorescent bulbs.

Age of inoculum: The fronds used to initiate the test were taken from a stock culture that had been transferred to fresh medium two days prior to testing.

B. STUDY DESIGN:

1. Experimental Conditions

a) Range-finding Study:

A 7-day preliminary range-finding exposure was conducted at Springborn Smithers at nominal ADTP metabolite of pyroxsulam concentrations of 0.010, 0.10, 1.0, 10 and 100 mg ac/L and a control. Two exposure vessels were established for each concentration and the control. Test solutions were renewed on days 2 and 4. All test solutions were clear and colorless with no visible undissolved test substance following solution preparation. Following 7 days of exposure, frond densities in the 0.010, 0.10, 1.0, 10 and 100 mg ac/L treatment levels averaged 379, 377, 330, 363 and 383 fronds/replicate, respectively. Frond density in the control averaged 317 fronds/replicate. Fronds exposed to the treatment levels tested and the control were normal. Based on these data, nominal concentrations of 6.3, 13, 25, 50 and 100 mg ac/L were selected for the definitive exposure.

b) Definitive Study

Table 2: Experimental Parameters

Parameter	Details	Remarks <i>Criteria</i>
------------------	----------------	------------------------------------

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Parameter	Details	Remarks Criteria
<p><u>Acclimation</u> Period:</p> <p>Culturing media and conditions: (same as test or not)</p> <p>Health: (any toxicity observed)</p>	<p>The fronds used to initiate the test were taken from a stock culture that had been transferred to fresh medium two days prior to testing.</p> <p>Stock cultures were grown in 270-mL covered crystallizing dishes containing 100 mL of 20X Algal Assay Procedure (AAP) medium. The cultures were maintained under test conditions.</p> <p>Not reported. However, control plants demonstrated satisfactory growth indicating healthy plants.</p>	<p>It is unclear how long stock cultures were maintained under test conditions.</p> <p><i>OECD: at least seven days before testing, sufficient colonies are transferred aseptically into fresh sterile medium and cultured for 7-10 days under the conditions of the test.</i></p> <p><i>EPA: axenic stock cultures should be grown in the aquariums for 2 weeks (with necessary transfers) prior to being used in a test. Inocula should be taken from cultures which are less than 2 weeks old.</i></p>
<p><u>Test system</u></p> <p>Static/static renewal/ Renewal rate for static renewal:</p>	<p>Static renewal Renewals on days 3 and 5.</p>	<p>Requirement considered met.</p> <p><i>EPA: Renewals (transfer of colonies to test solution) should occur on days 3 and 5.</i></p> <p><i>OECD: When using a semi-static test regime the colonies should be exposed to freshly prepared test and control solutions on at least two occasions during the test (e.g. days 3 and 5). The frequency of exposure to fresh medium will depend on the stability of the test substance.</i></p>
<p>Incubation facility</p>	<p>Environmental chamber within the following conditions: a temperature of $24 \pm 2^\circ\text{C}$ and continuous illumination of approximately 600 to 930 footcandles (6,500 to 10,000 lux). Lighting was supplied by Premira VitaLux® fluorescent bulbs.</p>	<p>Requirement considered met.</p> <p><i>OECD: temperature in the test vessels should be $24 \pm 2^\circ\text{C}$ and refers to use of a growth chamber incubator.</i></p> <p><i>EPA: temperature should be maintained at $25 \pm 2^\circ\text{C}$ and that a controlled environment growth chamber or an enclosed area</i></p>

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Parameter	Details	Remarks Criteria																																
		capable of maintaining the specified number of test chambers and test parameters is required.																																
Duration of the test	7 days.																																	
<u>Test vessel</u> Material: (glass/polystyrene) Size: Fill volume:	Glass crystallizing dishes. 270 mL 100 mL	Requirement considered met. <i>OECD: glass beakers, crystallising dishes or glass petri dishes of appropriate dimensions have all proved suitable. This guideline also states the test vessels must be covered. A minimum depth of 20 mm and minimum volume of 100 mL in each test vessel.</i> <i>EPA: test containers being glass beakers or Erlenmeyer flasks. Containers should be large enough to contain 150 mL of test solution, or enough test solution to result in a volume to-vessel size ratio of 2:5</i>																																
<u>Details of growth medium</u> Name:	20 Algal Assay Procedure (AAP) medium as follows: <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Compound</th> <th>Final concentration (mg/L)</th> </tr> </thead> <tbody> <tr><td>NaNO₃</td><td>510</td></tr> <tr><td>MgCl₂•6H₂O</td><td>240</td></tr> <tr><td>CaCl₂•2H₂O</td><td>90</td></tr> <tr><td>MgSO₄•7H₂O</td><td>290</td></tr> <tr><td>K₂HPO₄•3H₂O</td><td>30</td></tr> <tr><td>NaHCO₃</td><td>300</td></tr> <tr><td>H₃BO₃</td><td>3.7</td></tr> <tr><td>Na₂SeO₄^a</td><td>0.0376</td></tr> <tr><td>MnCl₂•4H₂O</td><td>8.3</td></tr> <tr><td>ZnCl₂</td><td>0.066</td></tr> <tr><td>CoCl₂•6H₂O</td><td>0.029</td></tr> <tr><td>CuCl₂•2H₂O</td><td>0.00024</td></tr> <tr><td>Na₂MoO₄•2H₂O</td><td>0.145</td></tr> <tr><td>FeCl₃•6H₂O</td><td>3.2</td></tr> <tr><td>Na₂EDTA•2H₂O</td><td>6.0</td></tr> </tbody> </table> <p>a) additional nutrient required</p>	Compound	Final concentration (mg/L)	NaNO ₃	510	MgCl ₂ •6H ₂ O	240	CaCl ₂ •2H ₂ O	90	MgSO ₄ •7H ₂ O	290	K ₂ HPO ₄ •3H ₂ O	30	NaHCO ₃	300	H ₃ BO ₃	3.7	Na ₂ SeO ₄ ^a	0.0376	MnCl ₂ •4H ₂ O	8.3	ZnCl ₂	0.066	CoCl ₂ •6H ₂ O	0.029	CuCl ₂ •2H ₂ O	0.00024	Na ₂ MoO ₄ •2H ₂ O	0.145	FeCl ₃ •6H ₂ O	3.2	Na ₂ EDTA•2H ₂ O	6.0	The growth medium used was based on that recommended in the OECD guideline. The only variation from this guideline was the addition of 0.0376 mg/L Na ₂ SeO ₄ . <i>OECD: For <i>L. gibba</i>, the OECD guideline recommends use of 20 AAP growth medium with the composition well defined in the guideline.</i>
Compound	Final concentration (mg/L)																																	
NaNO ₃	510																																	
MgCl ₂ •6H ₂ O	240																																	
CaCl ₂ •2H ₂ O	90																																	
MgSO ₄ •7H ₂ O	290																																	
K ₂ HPO ₄ •3H ₂ O	30																																	
NaHCO ₃	300																																	
H ₃ BO ₃	3.7																																	
Na ₂ SeO ₄ ^a	0.0376																																	
MnCl ₂ •4H ₂ O	8.3																																	
ZnCl ₂	0.066																																	
CoCl ₂ •6H ₂ O	0.029																																	
CuCl ₂ •2H ₂ O	0.00024																																	
Na ₂ MoO ₄ •2H ₂ O	0.145																																	
FeCl ₃ •6H ₂ O	3.2																																	
Na ₂ EDTA•2H ₂ O	6.0																																	
pH at test initiation: pH at test termination: Chelator used:	7.6 (control medium) 8.7 (control medium) EDTA.	<i>OECD: The pH of the control medium should initially be 7.5±1 and should not increase by more</i>																																

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Parameter	Details	Remarks Criteria
Carbon source:	Not reported.	than 1.5 units through the test.
If non-standard nutrient medium was used, detailed composition provided (Yes/No)	Standard AAP medium was used with the addition of 0.0376 mg/L Na ₂ SeO ₄ . Full details were provided.	Requirement considered met.
<u>Dilution water</u> Source/type: pH: Total Organic Carbon: Particulate matter: Metals: Pesticides: Chlorine: Water pretreatment (if any): Intervals of water quality measurement	 The 20X AAP medium used to prepare the exposure solutions was formulated in the same manner as the culture medium, prepared using sterile, deionized water and equilibrated to test temperature. Reported above for control medium. Typically 3 mg/L. Not reported. Not reported. Not reported. Not reported. Sterile, deionized water used. Measurements (pH) were made on days 0, 3 and 5 (new solutions) and days 3, 5 and 7 (old solutions).	 Requirements considered met. OECD 221 and US EPA OPPTS 850.4400 do not address these parameters specifically (other than pH). <i>OECD: The pH should be measured in each batch of 'fresh' test solution prior to each renewal and also in the corresponding 'spent' solutions.</i>
Indicate how the test material is added to the medium (added directly or used stock solution)	A 100 mg ac/L primary stock solution was prepared on the day of test initiation by placing 0.2042 g of test substance (0.20 g as active ingredient) in a 2000-mL volumetric flask and bringing it to volume with 20X AAP medium. The solution was sonicated for 30 minutes, stirred for around 2.5 hours. The stock solution was then observed to be clear and colorless with no visible undissolved test substance. Nominal test solutions were	

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Parameter	Details	Remarks <i>Criteria</i>
	<p>prepared from the primary stock solution.</p> <p>The 100 mg a.i./L primary stock solution and nominal test solutions were prepared at each solution renewal (days 3 and 5) following similar procedures.</p>	
Aeration or agitation	No aeration or agitation was reported.	Requirements considered met. OECD 221 and US EPA OPPTS 850.4400 do not specifically refer to aeration or agitation. OECD 221 notes that test vessels must be covered to minimise evaporation and accidental contamination, while allowing necessary air exchange.
<u>Sediment used (for rooted aquatic vascular plants)</u>	Not applicable.	
<u>Number of replicates</u>		Requirement considered met.
Control:	3	<p><i>OECD: at least 3 replicates should be used for each test concentration. The number of replicate control vessels should be at least equal to, and ideally twice, the number of vessels used for each test concentration.</i></p> <p><i>US EPA: for each concentration and control at least three replicate containers should be used.</i></p>
Solvent control:	Not applicable.	
Treatments:	3	
Number of plants/replicate	Expected around 5 (see below)	Requirements considered met. <i>OECD: each test vessel should contain a total of 9 to 12 fronds. The number of fronds and colonies should be the same in each test vessel.</i> <i>EPA: 3 to 5 plants per replicate.</i>
Number of fronds/plant	An inoculum of fifteen fronds was aseptically introduced into each test vessel by placing plants with two to four fronds each into each vessel until each	Requirements considered met. <i>OECD: colonies consisting of 2 to 4 visible fronds are transferred from the inoculum culture and randomly assigned to the test</i>

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Parameter	Details	Remarks Criteria
	vessel contained 15 fronds.	<i>vessels under aseptic conditions. Each test vessel should contain a total of 9 to 12 fronds. EPA: 3 to 4 fronds per plant.</i>
<u>Test concentrations</u> Nominal (mg ac/L): Measured (mg ac/L):	 0, 6.3, 13, 25, 50, 100 0, 5.7, 12, 23, 46, 93	Six concentrations were tested in a geometric series of (nominal) ~2.5. <i>OECD: in the definitive test, there should normally be at least five test concentrations arranged in a geometric series. Preferably the separation factor between test concentrations should not exceed 3.2.</i> <i>EPA at least five concentrations of chemical, exclusive of controls, should be used in the definitive test and chosen in a geometric series in which the ratio is between 1.5 and 2.0 (e.g. 2, 4, 8, 16, 32, 64 mg/L).</i>
Solvent (type, percentage, if used)	Not applicable.	
Method and interval of analytical verification: Limit of Quantitation: Limit of Detection:	All exposure solutions and QC samples were analyzed for test substance using high performance liquid chromatography with ultraviolet detection (HPLC/UV). Fresh solutions were analysed on days 0, 3 and 5. Old exposure solutions were analysed on days 3, 5 and 7. 0.0120 mg ac/L Not reported.	The requirements are considered met. The method validation study was conducted prior to the initiation of definitive testing and established an average recovery of 100±2.73% from 20X AAP medium.
<u>Test conditions</u> Temperature: Photoperiod: Light intensity and quality:	 23 to 24 °C Continuous illumination. 6500 to 9700 lux	Light intensity within OECD range but largely outside the range provided in the US EPA guideline. <i>OECD: temperature in the test vessels should be 24 ± 2°C with light intensity equivalent to 6500 to 10000 lux. Photoperiod not specified.</i>

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Parameter	Details	Remarks Criteria
		<i>EPA: environmental conditions should be maintained at 25 ± 2°C. Continuous warm-white fluorescent lighting should be used to provide a light intensity in the range of 4,200 and 6,700 lux</i>
<u>Reference chemical (if used)</u>	Not applicable.	Requirement considered met. There is no specific requirement to run a reference chemical test in conjunction with the test substance. It is unlikely that provision of the results from the most recent reference chemical study would have added any further value to interpretation of this test report. <i>OECD: a reference substance such as 3,5-dichlorophenol used in the international ring test may be tested as a means of checking the test procedure. It is advisable to test a reference substance at least twice a year or, where testing is carried out at a lower frequency, in parallel to the determination of the toxicity of a test substance.</i> <i>EPA: positive controls using zinc chloride as a reference chemical should be run periodically.</i>
Other parameters, if any	None.	

2. Observations:

Table 3: Observation parameters

Parameters	Details	Remarks Criteria
Parameters measured (eg: number of fronds, plant dry weight or other toxicity symptoms)	Frond production; growth rate; frond dry weight.	Requirement considered met. <i>OECD: Frond numbers are the primary parameter measured. In addition to determinations of frond number during the test, effects of the test substance on one or more of total frond area, dry weight or fresh weight</i>

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyrosulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

		<p><i>are also assessed.</i></p> <p><i>EPA: observations of frond numbers and appearance should be made of the colonies on day 0, 3, 5, and 7 and refers to other (optional) growth inhibition endpoints such as chlorophyll values and biomass (dry weight at 60°C) at the end of the test.</i></p>
Measurement technique for frond number and other end points	<p>On days 3, 5 and at test termination (day 7), fronds were counted (visual observation). At test termination (day 7), after frond density determinations were complete, the fronds were removed from each vessel, blotted dry and transferred to preweighed aluminum pans. Fronds were dried in an oven at 60-64°C for five days prior to dry weight determination.</p>	<p>Requirement considered met.</p> <p><i>OECD: Dry weight measurement - All colonies are collected from each of the test vessels and rinsed with distilled or deionised water. They are blotted to remove excess water and then dried at 60°C to a constant weight.</i></p>
Observation intervals	<p>Days 0, 3, 5 and 7 (frond numbers); Days 0-3; 0-5; and 0-7 (growth rate); Day 7 (frond dry weight).</p>	Requirement considered met.
Other observations, if any	<p>Frond appearance was observed at intervals measuring frond numbers. Throughout the exposure, fronds exposed to all treatment levels tested and the control were normal.</p>	
Indicate whether there was an exponential growth in the control	<p>Yes. Mean frond numbers in the control groups 349 after 7 days, or around a 23 fold increase over the test period. The average growth rate over the 7 day test period in the control was 0.46 d⁻¹.</p>	<p>Requirement considered met.</p> <p><i>OECD: For the test to be valid, the doubling time of frond number in the control must be less than 2.5 days (60 h), corresponding to approximately a seven-fold increase in seven days and an average specific growth rate of 0.275 d⁻¹.</i> <i>EPA: No specific requirements identified.</i></p>
Water quality was acceptable (Yes/No)	<p>Not specifically recorded in the test report but the successful control growth indicates the</p>	Requirement considered met.

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

<p>Were raw data included?</p>	<p>quality was acceptable. Yes, in transcribed form. Results for test end-points were provided in tabulated form and included results for individual replicates. Test condition parameters (pH, temperature and light intensity) were provided in tabulated form.</p>	<p>Requirement considered met. The transcribed data provided correspond to the OECD description of raw data requirements. While the EPA guideline does not comment on raw data, the reporting requirements outlined in this guideline were met.</p> <p><i>OECD: raw data: number of fronds and other measurement variables in each test and control vessel at each observation and occasion of analysis.</i></p> <p><i>EPA: No comment on raw data.</i></p>
--------------------------------	--	---

II. RESULTS AND DISCUSSION:

A. INHIBITORY EFFECTS:

Data sets for all end-points were shown statistically to pass requirements for normality and homogeneity of variance and therefore, Williams' Test was used to determine treatment-related effects.

No inhibitory effects were found for any biological endpoints evaluated in this study. For all biological end-points, the 7-day NOEC and LOEC were determined to be 93 and >93 mg ac/L respectively. At test termination, fronds exposed to all treatment levels were observed to be normal.

The pH of the newly formulated test and control solutions (day 0, 3 and 5) ranged from 7.6 to 8.2. The aged test and control solution pH (day 3, 5 and 7) ranged from 8.4 to 9.2. No other observations were made in the test report.

The frond counts from days 0 to 7 plus the calculated % inhibition compared to control counts, as given in the study report, are shown in Table 4. The growth rates for days 0-3, 0-5 and 0-7 plus the calculated % inhibition compared to control growth rates as given in the study report, are shown in Table 5. The day 7 frond dry weights plus the calculated % inhibition compared to control frond dry weight, as given in the study report, are shown in Table 6.

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Table 4: Effect of ADTP metabolite of pyroxsulam on frond number of Duckweed (*Lemna gibba*).

Mean Measured Concentration (mg ac/L)	Replicate No.	Frond number at:				7 Day Inhibition (%) compared to control.
		Day 0	Day 3	Day 5	Day 7	
Control	A	15	72	174	351	Not applicable
	B	15	60	149	317	
	C	15	68	169	380	
	Mean	15	67	164	349	
	SD	0	6	13	32	
5.7	A	15	69	171	359	-8
	B	15	60	168	359	
	C	15	75	195	410	
	Mean	15	68	178	376	
	SD	0	8	15	29	
12	A	15	84	169	460	-16
	B	15	64	191	413	
	C	15	63	161	344	
	Mean	15	70	174	406	
	SD	0	12	16	58	
23	A	15	68	196	499	-22
	B	15	72	184	405	
	C	15	68	181	379	
	Mean	15	69	187	428	
	SD	0	2	8	63	
46	A	15	67	199	425	-19
	B	15	58	172	415	
	C	15	55	160	405	
	Mean	15	60	177	415	
	SD	0	6	20	10	
93	A	15	64	172	411	-32
	B	15	78	196	498	
	C	15	74	202	470	
	Mean	15	72	190	460	
	SD	0	7	16	44	

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Table 5: Effect of ADTP metabolite of pyroxsulam on Growth Rate of Duckweed (*Lemna gibba*).

Mean Measured Concentration (mg ac/L)	Average Growth Rate (days ⁻¹)				
	Replicate No.	Observation interval (days)			7 Day Inhibition (%) compared to control.
		Day 0-3	Day 0-5	Day 0-7	
Control	A	0.54	0.51	0.47	Not applicable
	B	0.48	0.47	0.45	
	C	0.52	0.50	0.48	
	Mean	0.52	0.49	0.46	
	SD	0.03	0.02	0.01	
5.7	A	0.53	0.50	0.47	-4
	B	0.48	0.50	0.47	
	C	0.56	0.53	0.49	
	Mean	0.52	0.51	0.48	
	SD	0.04	0.01	0.01	
12	A	0.60	0.50	0.51	-7
	B	0.50	0.53	0.49	
	C	0.50	0.49	0.46	
	Mean	0.53	0.51	0.49	
	SD	0.06	0.02	0.02	
23	A	0.52	0.53	0.52	-7
	B	0.54	0.52	0.49	
	C	0.52	0.51	0.48	
	Mean	0.53	0.52	0.49	
	SD	0.01	0.01	0.02	
46	A	0.52	0.53	0.49	-7
	B	0.47	0.50	0.49	
	C	0.45	0.49	0.49	
	Mean	0.48	0.51	0.49	
	SD	0.04	0.02	0.00	
93	A	0.50	0.50	0.49	-9
	B	0.57	0.53	0.52	
	C	0.55	0.54	0.51	
	Mean	0.54	0.52	0.50	
	SD	0.04	0.02	0.01	

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Table 6: Effect of ADTP metabolite of pyroxsulam on Frond Dry Weight Duckweed (*Lemna gibba*).

Mean Measured Concentration (mg ac/L)	Frond weight (mg)		
	Replicate No.	Day 7	7 Day Inhibition (%) compared to control
Control	A	44.70	Not applicable
	B	39.20	
	C	52.50	
	Mean	45.47	
	SD	6.7	
5.7	A	40.50	6.4
	B	42.40	
	C	45.00	
	Mean	42.57	
	SD	2.3	
12	A	51.70	0.3
	B	44.70	
	C	39.60	
	Mean	45.33	
	SD	6.1	
23	A	61.90	-9.1
	B	44.20	
	C	42.70	
	Mean	49.60	
	SD	10.7	
46	A	47.10	-2.9
	B	49.40	
	C	43.80	
	Mean	46.77	
	SD	2.8	
93	A	49.20	-15.0
	B	55.50	
	C	52.20	
	Mean	52.30	
	SD	3.2	

Table 7: Statistical endpoint values.

Statistical Endpoint	Frond No.	Growth rate	Frond dry weight
NOEC (mg ac/L)	93	93	93
EC05 (mg ac/L) (95% C.I.)	>93	>93	>93
LOEC (mg ac/L)	>93	>93	>93
IC50 or EC50 (mg ac/L) (95% C.I.)	>93	>93	>93

EC05 and EC50 values estimated empirically so no confidence intervals calculated.

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

B. REPORTED STATISTICS:

Means and standard deviations of frond densities and growth rate were calculated for each treatment level and the control at each observation interval. Means and standard deviations for dry weight were also calculated for each treatment level and the control and were based on the dry plant weight determined after 7 days of exposure.

The growth rate (μ) for each replicate flask was calculated for the period from test initiation to each observation time.

Based on the results of statistical analysis performed for 7-day frond density, growth rate and biomass, the No-Observed-Effect Concentration (NOEC), the highest test concentration which demonstrated no statistically adverse effect ($p < 0.05$) when compared to the control data, was determined. Additionally, the Lowest-Observed-Effect Concentration (LOEC), the lowest concentration tested with a statistically significant reduction relative to the control data, was determined. The data were first checked for normality using Shapiro-Wilks' Test (Weber, et al., 1989) and for homogeneity of variance using Bartlett's Test (Horning and Weber, 1985). If the data sets passed the tests for homogeneity and normality, then Williams' Test (Williams, 1971, 1972) was used to determine the NOEC and LOEC. If the data did not pass the tests for homogeneity and normality, then Kruskal-Wallis' Test was used to determine the NOEC. All statistical determinations were made at the 95% level of certainty, except in the case of Shapiro-Wilks' and Bartlett's Tests, where the 99% level of certainty was applied.

The EC05, EC50 and EC90 values were calculated, when possible, for frond densities, average growth rate and biomass at test termination. TOXSTAT® version 3.5 (Gulley et al., 1996) was used to perform both the statistical (LOEC and NOEC determinations) and EC05, EC50 and EC90 calculations. If no concentration resulted in a 5% or 50%, the EC values were empirically estimated to be greater than the highest concentration tested.

C. VERIFICATION OF STATISTICAL RESULTS BY THE REVIEWER:

Due to a lack of inhibition observed in this study to all biological endpoints, no verification of statistical results is required.

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

D. STUDY DEFICIENCIES:

Study component	Deficiency
Details of Growth Medium:	The growth medium used was essentially the AAP medium recommended in OECD TG 221. However, in addition, 0.0376 mg/L selenate (Na_2SeO_4) was added as an additional nutrient, noted in the test report as based on personal communication. The need for this additional nutrient was not given. However, given the satisfactory growth of control plants this is not considered to have resulted in an impact on the study outcomes.
Test concentrations:	6 test concentrations were used in a geometric series of (nominal) ~2.5. This is within OECD guidance (separation factor between test concentrations should not exceed 3.2), but does not comply with EPA guidance where it states the geometric series should have a ratio between 1.5 and 2.0.
Test conditions.	Light intensity within OECD range but largely outside the range provided in the US EPA guideline. The protocol states that the light intensity will range from 6500 to 10,000 lux and the photosynthetically-active radiation (PAR) will range from 85 to 120 $\mu\text{E}/\text{m}^2/\text{s}$ (with the OECD guideline requirement being 85 to 135 $\mu\text{E}/\text{m}^2/\text{s}$). During the definitive test, the light intensity ranged from 6500 to 9700 lux and the PAR ranged from 102 to 147 $\mu\text{E}/\text{m}^2/\text{s}$. Since the light intensity was within the appropriate range, the PAR was not adjusted. This is not expected to have resulted in an impact on the study outcomes.
Acclimation period:	It is unclear how long stock cultures were maintained at the test facility prior to transferring fronds from the stock culture to fresh medium for use in the test. Given the strong growth of control plants, this is not expected to have resulted in an impact on the study outcomes.
Test monitoring	The study protocol requires the light intensity of the test area be measured daily. The light intensity was inadvertently not measured on day 1. Light intensity measurements were recorded on all other test days and were consistent within the required range. This is not expected to have resulted in an impact on the study outcomes.

E. REVIEWERS COMMENTS: Nothing additional.

The PMRA reviewer agrees with the conclusions of the Australian reviewer. This study is acceptable to the PMRA.

F. CONCLUSIONS: The study is acceptable. Based on this test, ADTP Metabolite of pyroxsulam is practically non-toxic to duckweed, *Lemna gibba*.

EC50/IC50: >93 mg ac/L (all endpoints); EC05: 93 mg ac/L (all endpoints).

III. REFERENCES:

Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

ASTM. 2002. Standard Guide for Conducting acute toxicity tests with fishes, macroinvertebrates and amphibians. Standard E729-96. American Society for Testing and Materials, 100 Barr Harbor Drive, West Conshohocken, PA 19426.

Environmental Monitoring and Support Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio.

Gulley, D.D., Boetler, A.M. and Bergman, H.L. 1996 TOXSTAT® Release 3.5. University of Wyoming, Laramie, Wyoming. Horning, W.B. and C.I. Weber. 1985. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms. EPA/600/4-85/014.

Hoberg J, 2006. Study Profile Template (SPT) for ADTP Metabolite of XDE-742 – Toxicity to Duckweed, *Lemna gibba*. Study No. 12550.6415.SPT. Springborn Smithers Laboratories, Massachusetts. 3 April 2006.

OECD, 1998. OECD Series on Principles of Good Laboratory Practice and Compliance Monitoring. Number 1. OECD Principles on Good Laboratory Practice (as revised in 1997). Environment Directorate Chemicals Group and Management Committee. ENV/MC/CHEM(98)17. OECD Paris. France. 41 pp.

OECD, 2000. OECD Guideline for Testing of Chemicals. *Lemna* sp. Growth Inhibition Test. Proposed Guideline #221. Revised Draft, October 2000.

U.S. EPA, 1996. Office of Prevention, Pesticides and Toxic Substances. Ecological Effects Test Guideline, OPPTS 850.4400. Aquatic Plant Toxicity Test Using *Lemna* spp., Tiers I and II. "Public Draft" EPA 712-C-96-156 April 1996. U.S. Environmental Protection Agency. Washington, D.C.

U.S. EPA. 1989. Federal Insecticide, Fungicide and Rodenticide Act (FIFRA); Good Laboratory Practice Standards; Final Rule (40 CFR, Part 160); FR: 8/17/89; pp. 34052. U.S. Environmental Protection Agency, Washington, D.C.

Weber, C.I., W.H. Peltier, T.J. Norberg-King, W.B. Horning II, F.A. Kessler, J.R. Menkedick, T.W. Neiheisel, P.A. Lewis, D.J. Klemm, Q.H. Pickering, E.L. Robinson, J.M. Lazorchak, L.J. Wymer and R.W. Freyberg (eds.). 1989. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms. 2nd ed. EPA/600/4/89/001. Environmental Monitoring Systems Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio.

Williams, D.A. 1971. A test for differences between treatment means when several dose levels are compared with a zero dose control. *Biometrics* 27: 103-117.

Williams, D.A. 1972. A comparison of several dose levels with a zero control. *Biometrics* 28: 519-531.

**Data Evaluation Report on the acute toxicity of ADTP metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283261

EPA MRID 469084-34

Approved 04/01/01 C.K.

US EPA ARCHIVE DOCUMENT