

US EPA ARCHIVE DOCUMENT

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID Number{.....}

469084-35

Data Requirement:

PMRA DATA CODE: 9.8.5

EPA DP Barcode:

OECD Test Guideline: IIA 8.6

EPA Guideline: 123-2 (OPPTS 850.4400 (Draft April 1996))

Test material:

Purity (%):

98%

Common name: 5,7-Di-OH Metabolite of pyroxsulam

Chemical name:

ID No: TSN 105233

Lot No. XN8-33938-53


IUPAC: Not available

IUPAC name: N-(5,7-dihydroxy[1,2,4]triazolo [1,5-a]pyrimidin-2-yl)-2-methoxy-4-(trifluoromethyl)-3- pyridinesulfonamide

CAS No.: Not available

Synonyms: 5,7-dihydroxy-XDE-742

Chemical structure



Primary Reviewer:

Chris Lee-Steere

Date: 5 July 2007

Australian Government Department of the Environment, Water, Heritage and the Arts

Secondary Reviewers:

Jack Holland

Date: 23 July 2007

Australian Government Department of the Environment, Water, Heritage and the Arts

Brian Kiernan

Date: 22 August 2007

US Environmental Protection Agency

PMRA Reviewer:

Émilie Larivière

Date: 30 July 2007

Environmental Assessment Directorate, PMRA

Reference/Submission No.: APVMA ATS 40362 NCRIS 61286

Company Code: DWE

Active Code: JUA

Use Site Category: 13, 14

EPA PC Code:

CITATION: Hoberg J, 2006. 5,7-Di-OH Metabolite of XDE-742 – Toxicity to Duckweed,

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257 EPA MRID Number{.....}

Lemna gibba. Springborn Smithers Study No. 12550.6418. Springborn Smithers
Laboratories, Wareham, Massachusetts. 30 March 2006. Unpublished.

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID Number 469084-35 {.....}

Data Requirement:

PMRA DATA CODE: 9.8.5

EPA DP Barcode:

OECD Test Guideline: IIA 8.6

EPA Guideline: 123-2 (OPPTS 850.4400 (Draft April 1996))

Test material:

Purity (%):

98%

Common name: 5,7-Di-OH Metabolite of pyroxsulam

Chemical name:

ID No: TSN 105233

Lot No. XN8-33938-53


IUPAC: Not available

IUPAC name: N-(5,7-dihydroxy[1,2,4]triazolo [1,5-a]pyrimidin-2-yl)-2-methoxy-4-(trifluoromethyl)-3- pyridinesulfonamide

CAS No.: Not available

Synonyms: 5,7-dihydroxy-XDE-742

Chemical structure



Primary Reviewer:

Chris Lee-Steere

Date: 5 July 2007

Australian Government Department of the Environment, Water, Heritage and the Arts

Secondary Reviewers:

Jack Holland

Date: 23 July 2007

Australian Government Department of the Environment, Water, Heritage and the Arts

Brian Kiernan

Date: 22 August 2007

US Environmental Protection Agency

PMRA Reviewer:

Émilie Larivière

Date: 30 July 2007

Environmental Assessment Directorate, PMRA

Reference/Submission No.: APVMA ATS 40362 NCRIS 61286

Company Code: DWE

Active Code: JUA

Use Site Category: 13, 14

EPA PC Code:

CITATION: Hoberg J, 2006. 5,7-Di-OH Metabolite of XDE-742 – Toxicity to Duckweed,

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257 EPA MRID Number{.....}

Lemna gibba. Springborn Smithers Study No. 12550.6418. Springborn Smithers
Laboratories, Wareham, Massachusetts. 30 March 2006. Unpublished.

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Data Requirement: PMRA DATA CODE: 9.8.5
EPA DP Barcode:
OECD Test Guideline: IIA 8.6
EPA Guideline: 123-2 (OPPTS 850.4400 (Draft April 1996))

Test material: **Purity (%):** 98%

Common name: 5,7-Di-OH Metabolite of pyroxsulam

Chemical name:

ID No: TSN 105233

Lot No. XN8-33938-53


IUPAC: Not available

IUPAC name: N-(5,7-dihydroxy[1,2,4]triazolo [1,5-a]pyrimidin-2-yl)-2-methoxy-4-(trifluoromethyl)-3- pyridinesulfonamide

CAS No.: Not available

Synonyms: 5,7-dihydroxy-XDE-742

Chemical structure



Primary Reviewer: Chris Lee-Steere **Date:** 5 July 2007
Australian Government Department of the Environment and Water Resources

Secondary Reviewers: Jack Holland **Date:** 23 July 2007
Australian Government Department of the Environment and Water Resources

Brian Kiernan **Date:** 22 August 2007

US Environmental Protection Agency

PMRA Reviewer: Émilie Larivière **Date:** 30 July 2007

Environmental Assessment Directorate, PMRA

Reference/Submission No.: APVMA ATS 40362 NCRIS 61286

Company Code: DWE

Active Code: JUA

Use Site Category: 13, 14

EPA PC Code:

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

CITATION: Hoberg J, 2006. 5,7-Di-OH Metabolite of XDE-742 – Toxicity to Duckweed, *Lemna gibba*. Springborn Smithers Study No. 12550.6418. Springborn Smithers Laboratories, Wareham, Massachusetts. 30 March 2006. Unpublished.

EXECUTIVE SUMMARY:

In a 7 day acute toxicity study, freshwater floating aquatic vascular plants (duckweed, *Lemna gibba*) were exposed to 5,7-Di-OH metabolite of pyroxsulam at nominal concentrations of 0, 1.0, 2.6, 6.0, 16, 40 and 100 mg ac/L. Mean measured concentrations were 0, 0.75, 2.1, 5.4, 14, 37 and 95 mg ac/L. The study was conducted under static renewal conditions at days 3 and 5 in accordance with the guidelines, OECD 221 "Lemna sp. Growth Inhibition Test" (draft, 2002) and US EPA guidelines including U.S. Environmental Protection Agency (1996). Ecological Effects Test Guidelines. OPPTS 850.4400 Aquatic Plant Toxicity Test using Lemna sp., Tiers I and II. Draft April 1996.

The % growth inhibition was determined for frond number, mean specific growth rate and biomass (frond dry weight). The only statistically significant inhibitory effects found in the study were for frond numbers at the highest tested rate (20% inhibition compared to control) and growth rates (6% inhibition compared to control). The corresponding EC05 values for frond numbers and growth rates were calculated as 24 and 69 mg ac/L respectively. No differences between control plants and treatment rates were found for frond dry weights. The 7-day NOECs based on frond number, specific growth rates and biomass (dry weight at 7 days) were 37, 37 and 95 mg ac/L respectively (mean measured concentration). The EC50 values for all endpoints were >95 mg ac/L (mean measured concentration), determined empirically.

Curled fronds on plants exposed to 37 and 95 mg ac/L were observed at day 7. No other abnormalities were observed or recorded.

This toxicity study is classified as acceptable and satisfies the guideline requirement for an acute toxicity study with the aquatic vascular plants *Lemna gibba* (duckweed).

Results Synopsis

Test Organism: Duckweed (*Lemna gibba*)

Test Type (Flowthrough, Static, Static Renewal): Static renewal

Endpoint	Frond No.	Growth rate	Frond dry weight
NOEC (mg ac/L)	37	37	95
EC05 (mg ac/L) (95% C.I.)	24 (2.9-52)	69 (18-86)	57 (1.4-85)
LOEC (mg ac/L)	95	95	>95
IC50 or EC50 (mg ac/L) (95% C.I.)	>95	>95	>95

No 95% confidence intervals associated with EC50 values as these were determined empirically.

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Endpoint(s) Effected: 7 day frond number and growth rate.

I. MATERIALS AND METHODS

GUIDELINE FOLLOWED:

OECD, 2000. OECD Guideline for Testing of Chemicals. *Lemna* sp. Growth Inhibition Test. Proposed Guideline #221. Revised Draft, October 2000.

U.S. EPA, 1996. Office of Prevention, Pesticides and Toxic Substances. Ecological Effects Test Guideline, OPPTS 850.4400. Aquatic Plant Toxicity Test Using *Lemna* spp., Tiers I and II. "Public Draft" EPA 712-C-96-156 April 1996. U.S. Environmental Protection Agency. Washington, D.C.

The following protocol deviation is noted in the study report:

The protocol states that light intensity will range from 6500 to 10,000 lux and the photosynthetically-active radiation (PAR) will range from 85 to 120 $\mu\text{E}/\text{m}^2/\text{s}$. During the definitive test, the light intensity ranged from 6500 to 9400 lux and the PAR ranged from 112 to 138 $\mu\text{E}/\text{m}^2/\text{s}$. Since the light intensity was within the appropriate range, the PAR was not adjusted.

This deviation is not expected to influence the study results.

This DER has assessed the study report against the OECD 221 and US EPA OPPTS 850.4400 requirements.

COMPLIANCE: All phases of the study were reported as conducted in compliance with the following Good Laboratory Practice Standards:

- OECD Series on Principles of Good Laboratory Practice and Compliance Monitoring, Number 1. OECD Principles on Good Laboratory Practice (as revised in 1997) ENVIMCICHEM (98) 17; and
- U.S. Environmental Protection Agency - FIFRA GLPs, Title 40 CFR, Part 160- Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), Good Laboratory Practice Standards, Final Rule.

Signed and dated Compliance with Good Laboratory Practice Standards, Quality Assurance and No Data Confidentiality Claims statements were provided.

A. MATERIALS:

1. Test Material

5,7-Di-OH Metabolite of pyroxsulam

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Description: Solid

Lot No./Batch No.: XN8-33938-53

Purity: 98% (Certificate of analysis provided)

Stability of Compound Under Test Conditions:

At the beginning and end of two renewal periods (i.e., days 0 and 3 and days 5 and 7), one sample was removed from each test solution and the control and analyzed for 5,7-di-OH metabolite of pyroxsulam. Samples analyzed from newly prepared solutions were removed from the volumetric flasks prior to division into the replicate vessels. Test solution samples analyzed at the end of the renewal periods (days 3 and 7) were removed from composited solutions of each treatment level and the control. The following results were found:

Table 1: Measured concentrations of 5,7-di-OH Metabolite of pyroxsulam

Nominal Concentration (mg ac/L)	Measured Concentration (mg ac/L)					% of Nominal
	0 hour (new)	Day 3 (aged)	Day 5 (new)	Day 7 (aged)	Mean (SD)	
Control	<0.028	<0.040	<0.039	<0.043	Not applicable	Not applicable
1.0	0.87	0.41	0.97	0.76	0.75 (0.24)	75
2.6	2.3	1.4	2.5	2.3	2.1 (0.49)	81
6.4	5.8	4.1	6.1	5.7	5.4 (0.87)	85
16	15	13	16	15	14 (1.2)	91
40	35	35	39	39	37 (2.4)	92
100	91	94	98	97	95 (3.3)	95

There are no stability data for the test substance under light.

Storage conditions of test chemicals:

Stored refrigerated (1 to 10°C) in the original container

Physicochemical properties of 5,7-di-OH metabolite of pyroxsulam: None available at the time of testing.

2. Test organism:

Name: Duckweed (*Lemna gibba*)

Strain, if provided: 310

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Source: Obtained from the University of Toronto, Toronto, Canada and maintained in stock culture at Springborn Smithers Laboratories, Wareham, MA.

Method of cultivation: Stock cultures were grown in 270-mL covered crystallizing dishes containing 100 mL of medium. The cultures were maintained in an environmental chamber within the following conditions: a temperature of $24 \pm 2^\circ\text{C}$ and continuous illumination of approximately 600 to 930 footcandles (6500 to 10,000 lux). Lighting was supplied by Premira VitaLux® fluorescent bulbs.

Age of inoculum: The fronds used to initiate the test were taken from a stock culture that had been transferred to fresh medium two days prior to testing.

B. STUDY DESIGN:

1. Experimental Conditions

a) **Range-finding Study:** A 7-day range-finding study was conducted at nominal 7-OH metabolite of pyroxsulam concentrations of 0.010, 0.10, 1.0, 10 and 100 mg ac/L and a control. Two exposure vessels were established for each concentration and the control. Test solutions were renewed on days 3 and 5. All test solutions were clear and colorless with no visible undissolved test substance following solution preparation. Following 7 days of exposure, frond densities in the 0.010, 0.10, 1.0, 10 and 100 mg ac/L treatment levels averaged 395, 296, 332, 282 and 198 fronds/replicate, respectively. Frond density in the control averaged 334 fronds/replicate. Fronds exposed to the 100 mg ac/L treatment level were observed to be slightly chlorotic and curled. Fronds exposed to the remaining treatment levels (0.010, 0.10, 1.0 and 10 mg ac/L) and the control were normal. Based on these data, nominal concentrations of 1.0, 2.6, 6.4, 16, 40 and 100 mg ac/L were selected for the definitive exposure.

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

b) Definitive Study

Table 2: Experimental Parameters

Parameter	Details	Remarks Criteria
<u>Acclimation</u> Period: Culturing media and conditions: (same as test or not) Health: (any toxicity observed)	<p>The fronds used to initiate the test were taken from a stock culture that had been transferred to fresh medium two days prior to testing.</p> <p>Stock cultures were grown in 270-mL covered crystallizing dishes containing 100 mL of 20X Algal Assay Procedure (AAP) medium. The cultures were maintained under test conditions.</p> <p>Not reported. However, control plants demonstrated satisfactory growth indicating healthy plants.</p>	<p>It is unclear how long stock cultures were maintained under test conditions.</p> <p><i>OECD: at least seven days before testing, sufficient colonies are transferred aseptically into fresh sterile medium and cultured for 7-10 days under the conditions of the test.</i></p> <p><i>EPA: axenic stock cultures should be grown in the aquariums for 2 weeks (with necessary transfers) prior to being used in a test. Inocula should be taken from cultures which are less than 2 weeks old.</i></p>
<u>Test system</u> Static/static renewal/ Renewal rate for static renewal:	<p>Static renewal</p> <p>Renewals on days 3 and 5.</p>	<p><i>EPA: Renewals (transfer of colonies to test solution) should occur on days 3 and 5.</i></p> <p><i>OECD: When using a semi-static test regime the colonies should be exposed to freshly prepared test and control solutions on at least two occasions during the test (e.g. days 3 and 5). The frequency of exposure to fresh medium will depend on the stability of the test substance.</i></p>
Incubation facility	<p>Environmental chamber within the following conditions: a temperature of $24 \pm 2^\circ\text{C}$ and continuous illumination of approximately 600 to 930 footcandles (6,500 to 10,000 lux). Lighting was supplied by</p>	<p>Requirement considered met.</p> <p><i>OECD: temperature in the test vessels should be $24 \pm 2^\circ\text{C}$ and refers to use of a growth chamber incubator.</i></p>

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Parameter	Details	Remarks Criteria																																
	Premira VitaLux® fluorescent bulbs.	EPA: temperature should be maintained at 25 ± 2°C and that a controlled environment growth chamber or an enclosed area capable of maintaining the specified number of test chambers and test parameters is required.																																
Duration of the test	7 days.																																	
<u>Test vessel</u> Material: (glass/polystyrene) Size: Fill volume:	Glass crystallizing dishes. 270 mL 100 mL	Requirement considered met. OECD: glass beakers, crystallising dishes or glass petri dishes of appropriate dimensions have all proved suitable. This guideline also states the test vessels must be covered. A minimum depth of 20 mm and minimum volume of 100 mL in each test vessel. EPA: test containers being glass beakers or Erlenmeyer flasks. Containers should be large enough to contain 150 mL of test solution, or enough test solution to result in a volume to-vessel size ratio of 2:5																																
<u>Details of growth medium</u> Name:	20 Algal Assay Procedure (AAP) medium as follows: <table><tr><th>Compound</th><th>Final concentration (mg/L)</th></tr><tr><td>NaNO₃</td><td>510</td></tr><tr><td>MgCl₂•6H₂O</td><td>240</td></tr><tr><td>CaCl₂•2H₂O</td><td>90</td></tr><tr><td>MgSO₄•7H₂O</td><td>290</td></tr><tr><td>K₂HPO₄•3H₂O</td><td>30</td></tr><tr><td>NaHCO₃</td><td>300</td></tr><tr><td>H₃BO₃</td><td>3.7</td></tr><tr><td>Na₂SeO₄^a</td><td>0.0376</td></tr><tr><td>MnCl₂•4H₂O</td><td>8.3</td></tr><tr><td>ZnCl₂</td><td>0.066</td></tr><tr><td>CoCl₂•6H₂O</td><td>0.029</td></tr><tr><td>CuCl₂•2H₂O</td><td>0.00024</td></tr><tr><td>Na₂MoO₄•2H₂O</td><td>0.145</td></tr><tr><td>FeCl₃•6H₂O</td><td>3.2</td></tr><tr><td>Na₂EDTA•2H₂O</td><td>6.0</td></tr></table> a) additional nutrient required 7.8 (control medium)	Compound	Final concentration (mg/L)	NaNO ₃	510	MgCl ₂ •6H ₂ O	240	CaCl ₂ •2H ₂ O	90	MgSO ₄ •7H ₂ O	290	K ₂ HPO ₄ •3H ₂ O	30	NaHCO ₃	300	H ₃ BO ₃	3.7	Na ₂ SeO ₄ ^a	0.0376	MnCl ₂ •4H ₂ O	8.3	ZnCl ₂	0.066	CoCl ₂ •6H ₂ O	0.029	CuCl ₂ •2H ₂ O	0.00024	Na ₂ MoO ₄ •2H ₂ O	0.145	FeCl ₃ •6H ₂ O	3.2	Na ₂ EDTA•2H ₂ O	6.0	The growth medium used was based on that recommended in the OECD guideline. The only variation from this guideline was the addition of 0.0376 mg/L Na ₂ SeO ₄ . OECD: For <i>L. gibba</i> , the OECD guideline recommends use of 20 AAP growth medium with the composition well defined in the guideline.
Compound	Final concentration (mg/L)																																	
NaNO ₃	510																																	
MgCl ₂ •6H ₂ O	240																																	
CaCl ₂ •2H ₂ O	90																																	
MgSO ₄ •7H ₂ O	290																																	
K ₂ HPO ₄ •3H ₂ O	30																																	
NaHCO ₃	300																																	
H ₃ BO ₃	3.7																																	
Na ₂ SeO ₄ ^a	0.0376																																	
MnCl ₂ •4H ₂ O	8.3																																	
ZnCl ₂	0.066																																	
CoCl ₂ •6H ₂ O	0.029																																	
CuCl ₂ •2H ₂ O	0.00024																																	
Na ₂ MoO ₄ •2H ₂ O	0.145																																	
FeCl ₃ •6H ₂ O	3.2																																	
Na ₂ EDTA•2H ₂ O	6.0																																	

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Parameter	Details	Remarks Criteria
pH at test initiation: pH at test termination: Chelator used: Carbon source:	9.0 (control medium) EDTA. Not reported.	<i>OECD: The pH of the control medium should initially be 7.5±1 and should not increase by more than 1.5 units through the test.</i>
If non-standard nutrient medium was used, detailed composition provided (Yes/No)	Standard AAP medium was used with the addition of 0.0376 mg/L Na ₂ SeO ₄ . Full details were provided.	Requirement considered met.
<u>Dilution water</u> Source/type: pH: Total Organic Carbon: Particulate matter: Metals: Pesticides: Chlorine: Water pretreatment (if any): Intervals of water quality measurement	The 20X AAP medium used to prepare the exposure solutions was formulated in the same manner as the culture medium. Several liters of 20X AAP medium were prepared using sterile, deionized water and equilibrated to test temperature. Reported above for control medium. Typically 3 mg/L. Not reported. Not reported. Not reported. Not reported. Sterile, deionized water used. Measurements (pH) were made on days 0, 3 and 5 (new solutions) and days 3, 5 and 7 (old solutions).	Requirements considered met. OECD 221 and US EPA OPPTS 850.4400 do not address these parameters specifically (other than pH). <i>OECD: The pH should be measured in each batch of 'fresh' test solution prior to each renewal and also in the corresponding 'spent' solutions.</i>
Indicate how the test material is added to the medium (added directly or used stock solution)	A 100 mg ac/L primary stock solution was prepared on the day of test initiation by placing 0.1020 g of test substance (0.10 g as active ingredient) in a 1000-mL volumetric flask and bringing it to volume with 20X AAP medium. Following 15 minutes of sonication and stirring for an	

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Parameter	Details	Remarks Criteria
	additional 15 minutes, the stock solution was observed to be clear and slightly yellow in color with no visible undissolved test substance present. Nominal test solutions were prepared from the primary stock solution.	
Aeration or agitation	No aeration or agitation was reported.	Requirements considered met. OECD 221 and US EPA OPPTS 850.4400 do not specifically refer to aeration or agitation. OECD 221 notes that test vessels must be covered to minimise evaporation and accidental contamination, while allowing necessary air exchange.
<u>Sediment used (for rooted aquatic vascular plants)</u>	Not applicable.	
<u>Number of replicates</u> Control: Solvent control: Treatments:	3 Not applicable. 3	Requirement considered met. <i>OECD: at least 3 replicates should be used for each test concentration. The number of replicate control vessels should be at least equal to, and ideally twice, the number of vessels used for each test concentration.</i> <i>US EPA: for each concentration and control at least three replicate containers should be used.</i>
Number of plants/replicate	5	Requirements considered met. <i>OECD: each test vessel should contain a total of 9 to 12 fronds. The number of fronds and colonies should be the same in each test vessel.</i> <i>EPA: 3 to 5 plants per replicate.</i>
Number of fronds/plant	3	Requirements considered met. <i>OECD: colonies consisting of 2 to 4 visible fronds are transferred from the inoculum culture and randomly assigned to the test</i>

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Parameter	Details	Remarks <i>Criteria</i>
		vessels under aseptic conditions. Each test vessel should contain a total of 9 to 12 fronds. EPA: 3 to 4 fronds per plant.
<u>Test concentrations</u> Nominal (mg ac/L): Measured (mg ac/L):	 0, 1.0, 2.6, 6.4, 16, 40, 100 0, 0.75, 2.1, 5.4, 14, 37, 95	Six concentrations were tested in geometric series of (nominal) ~2.5. <i>OECD: in the definitive test, there should normally be at least five test concentrations arranged in a geometric series. Preferably the separation factor between test concentrations should not exceed 3.2.</i> <i>EPA at least five concentrations of chemical, exclusive of controls, should be used in the definitive test and chosen in a geometric series in which the ratio is between 1.5 and 2.0 (e.g. 2, 4, 8, 16, 32, 64 mg/L).</i>
Solvent (type, percentage, if used)	Not applicable.	
Method and interval of analytical verification: Limit of Quantitation: Limit of Detection:	All exposure solutions and QC samples were analyzed for test substance using high performance liquid chromatography with ultraviolet detection (HPLC/UV). Fresh solutions were analysed on days 0, 3 and 5. Old exposure solutions were analysed on days 3, 5 and 7. 0.0203 mg ac/L Not reported.	The requirements are considered met. The method validation study was conducted prior to the initiation of definitive testing and established an average recovery of $96.1 \pm 6.13\%$ from 20X AAP medium.
<u>Test conditions</u> Temperature: Photoperiod: Light intensity and quality:	 22 to 24 °C Continuous illumination. 6500 to 9400 lux	Light intensity within OECD range but largely outside the range provided in the US EPA guideline. <i>OECD: temperature in the test vessels should be $24 \pm 2^{\circ}\text{C}$ with light intensity equivalent to 6500 to 10000 lux. Photoperiod not specified.</i>

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Parameter	Details	Remarks Criteria
		<i>EPA: environmental conditions should be maintained at $25 \pm 2^{\circ}\text{C}$. Continuous warm-white fluorescent lighting should be used to provide a light intensity in the range of 4,200 and 6,700 lux</i>
<u>Reference chemical (if used)</u>	Not applicable.	Requirement considered met. There is no specific requirement to run a reference chemical test in conjunction with the test substance. It is unlikely that provision of the results from the most recent reference chemical study would have added any further value to interpretation of this test report. <i>OECD: a reference substance such as 3,5-dichlorophenol used in the international ring test may be tested as a means of checking the test procedure. It is advisable to test a reference substance at least twice a year or, where testing is carried out at a lower frequency, in parallel to the determination of the toxicity of a test substance.</i> <i>EPA: positive controls using zinc chloride as a reference chemical should be run periodically.</i>
Other parameters, if any	None.	

2. Observations:

Table 3: Observation parameters

Parameters	Details	Remarks Criteria
Parameters measured (eg: number of fronds, plant dry weight or other toxicity symptoms)	Frond density; growth rate; frond dry weight.	Requirement considered met. <i>OECD: Frond numbers are the primary parameter measured. In addition to determinations of frond number during the test, effects of the test substance on one or more of total frond area, dry weight or fresh weight</i>

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

		<p><i>are also assessed.</i></p> <p><i>EPA: observations of frond numbers and appearance should be made of the colonies on day 0, 3, 5, and 7 and refers to other (optional) growth inhibition endpoints such as chlorophyll values and biomass (dry weight at 60°C) at the end of the test.</i></p>
Measurement technique for frond number and other end points	<p>On days 3, 5 and at test termination (day 7), fronds were counted (visual observation). At test termination (day 7), after frond density determinations were complete, the fronds were removed from each vessel, blotted dry and transferred to preweighed aluminum pans. Fronds were dried in an oven at 61°C for three days prior to dry weight determination.</p>	<p>Requirement considered met.</p> <p><i>OECD: Dry weight measurement - All colonies are collected from each of the test vessels and rinsed with distilled or deionised water. They are blotted to remove excess water and then dried at 60°C to a constant weight.</i></p>
Observation intervals	<p>Days 0, 3, 5 and 7 (frond numbers);</p> <p>Days 0-3; 0-5; and 0-7 (growth rate);</p> <p>Day 7 (frond dry weight).</p>	Requirement considered met.
Other observations, if any	<p>Frond appearance was observed at intervals measuring frond numbers. At test termination, fronds exposed to treatment levels ≥ 37 mg ac/L were curled.</p>	
Indicate whether there was an exponential growth in the control	<p>Yes.</p> <p>Mean frond numbers in the control groups 425 after 7 days, or around a 28 fold increase over the test period.</p> <p>The average growth rate over the 7 day test period in the control was 0.48 d^{-1}.</p>	<p>Requirement considered met.</p> <p><i>OECD: For the test to be valid, the doubling time of frond number in the control must be less than 2.5 days (60 h), corresponding to approximately a seven-fold increase in seven days and an average specific growth rate of 0.275 d^{-1}.</i></p> <p><i>EPA: No specific requirements identified.</i></p>
Water quality was acceptable (Yes/No)	<p>Not specifically recorded in the test report but the successful control growth indicates the quality was acceptable.</p>	Requirement considered met.

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Were raw data included?	Yes, in transcribed form. Results for test end-points were provided in tabulated form and included results for individual replicates. Test condition parameters (pH, temperature and light intensity) were provided in tabulated form.	Requirement considered met. The transcribed data provided correspond to the OECD description of raw data requirements. While the EPA guideline does not comment on raw data, the reporting requirements outlined in this guideline were met. <i>OECD: raw data: number of fronds and other measurement variables in each test and control vessel at each observation and occasion of analysis.</i> <i>EPA: No comment on raw data.</i>
-------------------------	--	--

II. RESULTS AND DISCUSSION:

A. INHIBITORY EFFECTS:

Based on statistical analysis, inhibitory effects were only found at the highest test concentration for frond numbers and growth rate. No inhibitory effects were found in terms of frond dry weight. No dose-response relationship could therefore be determined.

At test termination, fronds exposed to the 37 and 95 mg ac/L treatment levels tested were observed to be curled. Fronds exposed to the remaining treatment levels tested (0.75, 2.1, 5.4 and 14 mg ac/L) and the control were observed to be normal. The pH of the newly formulated test and control solutions (day 0, 3 and 5) ranged from 7.2 to 7.9. The aged test and control solution pH (day 3, 5 and 7) ranged from 8.3 to 9.1. No other observations were made in the test report.

The frond counts from days 0 to 7 plus the calculated % inhibition compared to control counts, as given in the study report, are shown in Table 4. The growth rates for days 0-3, 0-5 and 0-7 plus the calculated % inhibition compared to control growth rates as given in the study report, are shown in Table 5. The day 7 frond dry weights plus the calculated % inhibition compared to control frond dry weight, as given in the study report, are shown in Table 6.

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Table 4: Effect of 5,7-di-OH metabolite of pyroxsulam on frond number of Duckweed (*Lemna gibba*).

Mean Measured Concentration (mg ac/L)	Replicate No.	Frond number at:				7 Day Inhibition (%) compared to control.
		Day 0	Day 3	Day 5	Day 7	
Control	A	15	78	230	510	NA ^a
	B	15	74	187	384	
	C	15	68	183	381	
	Mean	15	73	200	425	
	SD	0	5	26	74	
0.75	A	15	70	210	492	-1
	B	15	74	192	397	
	C	15	58	179	402	
	Mean	15	67	194	450	
	SD	0	8	16	53	
2.1	A	15	66	193	438	-7
	B	15	63	193	438	
	C	15	78	209	486	
	Mean	15	69	198	454	
	SD	0	8	9	28	
5.4	A	15	74	193	405	-7
	B	15	78	223	556	
	C	15	62	197	400	
	Mean	15	71	204	454	
	SD	0	8	16	89	
14	A	15	78	206	463	0
	B	15	61	185	402	
	C	15	68	198	411	
	Mean	15	69	196	425	
	SD	0	9	11	33	
37	A	15	68	192	368	4
	B	15	78	213	402	
	C	15	70	187	458	
	Mean	15	72	197	409 ^b	
	SD	0	5	14	45	
95	A	15	62	165	339	20
	B	15	63	181	353	
	C	15	72	180	330	
	Mean	15	66	175	341 ^{bc}	
	SD	0	6	9	12	

a) Not applicable; b) Curled fronds were observed; c) Significantly reduced compared to the control, based on Williams' Test.

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Table 5: Effect of 5,7-di-OH metabolite of pyroxsulam on Growth Rate of Duckweed (*Lemna gibba*).

Average Growth Rate (days ⁻¹)					
Mean Measured Concentration (mg ac/L)	Replicate No.	Observation interval (days)			7 Day Inhibition (%) compared to control.
		Day 0-3	Day 0-5	Day 0-7	
	A	0.56	0.54	0.51	NA ^a
	B	0.54	0.50	0.47	
	C	0.51	0.50	0.47	
	Mean	0.53	0.52	0.48	
	SD	0.02	0.03	0.02	
0.75	A	0.52	0.53	0.51	-2
	B	0.54	0.51	0.48	
	C	0.46	0.49	0.48	
	Mean	0.50	0.51	0.49	
	SD	0.04	0.02	0.02	
2.1	A	0.50	0.51	0.49	-2
	B	0.48	0.51	0.49	
	C	0.56	0.53	0.50	
	Mean	0.51	0.52	0.49	
	SD	0.04	0.01	0.01	
5.4	A	0.54	0.51	0.48	-2
	B	0.56	0.54	0.52	
	C	0.48	0.51	0.48	
	Mean	0.52	0.52	0.49	
	SD	0.04	0.02	0.03	
14	A	0.56	0.52	0.50	-2
	B	0.47	0.50	0.48	
	C	0.51	0.51	0.48	
	Mean	0.51	0.51	0.49	
	SD	0.04	0.01	0.01	
37	A	0.51	0.51	0.46	0
	B	0.56	0.53	0.48	
	C	0.52	0.50	0.50	
	Mean	0.53	0.51	0.48	
	SD	0.02	0.01	0.02	
95	A	0.48	0.48	0.45	6
	B	0.48	0.50	0.46	
	C	0.53	0.50	0.45	
	Mean	0.50	0.49	0.45 ^b	
	SD	0.03	0.01	0.0	

a) Not applicable; b) Significantly reduced compared to the control, based on Williams' Test.

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Table 6: Effect of 5,7-di-OH metabolite of pyroxsulam on Frond Dry Weight Duckweed (*Lemna gibba*).

Mean Measured Concentration (mg ac/L)	Frond dry weight (g)		
	Replicate No.	Day 7	7 Day Inhibition (%) compared to control.
Control	A	0.0514	Not applicable
	B	0.0449	
	C	0.0414	
	Mean	0.0459	
	SD	0.0051	
0.75	A	0.0645	-11.0
	B	0.0418	
	C	0.0465	
	Mean	0.0509	
	SD	0.012	
2.1	A	0.0467	-5.0
	B	0.0448	
	C	0.0531	
	Mean	0.0482	
	SD	0.0043	
5.4	A	0.0431	-20.4
	B	0.0801	
	C	0.0426	
	Mean	0.0553	
	SD	0.022	
14	A	0.0533	-1.2
	B	0.0436	
	C	0.0424	
	Mean	0.0464	
	SD	0.0060	
37	A	0.0429	-11.3
	B	0.0426	
	C	0.0678	
	Mean	0.0511	
	SD	0.015	
95	A	0.0423	1.4
	B	0.0518	
	C	0.0417	
	Mean	0.0453	
	SD	0.0057	

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Table 7: Statistical endpoint values.*

Statistical Endpoint	Frond No.	Growth rate	Frond dry weight
NOEC (mg ac/L)	37	37	95
EC05 (mg ac/L) (95% C.I.)	24 (2.9-52)	69 (18-86)	57 (1.4-85)
LOEC (mg ac/L)	95	95	>95
IC50 or EC50 (mg ac/L) (95% C.I.)	>95	>95	>95

No 95% confidence intervals associated with IC50 values as these were determined empirically.

B. REPORTED STATISTICS:

Means and standard deviations of frond densities and growth rate were calculated for each treatment level and the control at each observation interval. Means and standard deviations for dry weight were also calculated for each treatment level and the control and were based on the dry plant weight determined after 7 days of exposure.

The growth rate (μ) for each replicate flask was calculated for the period from test initiation to each observation time.

Based on the results of statistical analysis performed for 7-day frond density, growth rate and biomass, the No-Observed-Effect Concentration (NOEC), the highest test concentration which demonstrated no statistically adverse effect ($p < 0.05$) when compared to the control data, was determined. Additionally, the Lowest-Observed-Effect Concentration (LOEC), the lowest concentration tested with a statistically significant reduction relative to the control data, was determined. The data were first checked for normality using Shapiro-Wilks' Test (Weber, et al., 1989) and for homogeneity of variance using Bartlett's Test (Horning and Weber, 1985). If the data sets passed the tests for homogeneity and normality, then Williams' Test (Williams, 1971, 1972) was used to determine the NOEC and LOEC. If the data did not pass the tests for homogeneity and normality, then Kruskal-Wallis' Test was used to determine the NOEC. All statistical determinations were made at the 95% level of certainty, except in the case of Shapiro-Wilks' and Bartlett's Tests, where the 99% level of certainty was applied.

The EC05, EC50 and EC90 values were calculated, when possible, for frond densities, average growth rate and biomass at test termination. TOXSTAT® version 3.5 (Gulley et al., 1996) was used to perform both the statistical (LOEC and NOEC determinations) and EC05, EC50 and EC90 calculations. Due to a lack on inhibition in this study, the EC50 and EC90 values were empirically estimated to be greater than the highest concentration tested.

C. VERIFICATION OF STATISTICAL RESULTS BY THE REVIEWER:

To verify the statistics, the data have been re-analysed using TOXCALC – Toxicity Data

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Analysis Software v5.0.26. Given the general lack of inhibition found in this study, only the EC05 values for frond numbers and growth rates were verified using non-linear interpolation. The following results were found:

Statistical Endpoint	Frond No.	Growth rate
EC05 (mg ac/L) (95% C.I.)	23.5 (0.0-82.25)	68.6 (no CI generated)

The results are in good agreement with those found in the study.

D. STUDY DEFICIENCIES:

Study component	Deficiency
Details of Growth Medium:	The growth medium used was essentially the AAP medium recommended in OECD TG 221. However, in addition, 0.0376 mg/L selenate (Na_2SeO_4) was added as an additional nutrient, noted in the test report as based on personal communication. The need for this additional nutrient was not given. However, given the satisfactory growth of control plants this is not considered to have resulted in an impact on the study outcomes.
Test concentrations:	6 test concentrations were used in a geometric series of (nominal) ~2.5. This is within OECD guidance (separation factor between test concentrations should not exceed 3.2), but does not comply with EPA guidance where it states the geometric series should have a ratio between 1.5 and 2.0.
Test conditions.	Light intensity within OECD range but largely outside the range provided in the US EPA guideline. The protocol states that the light intensity will range from 6500 to 10,000 lux and the photosynthetically-active radiation (PAR) will range from 85 to 120 $\mu\text{E}/\text{m}^2/\text{s}$ (with the OECD guideline requirement being 85 to 135 $\mu\text{E}/\text{m}^2/\text{s}$). During the definitive test, the light intensity ranged from 6500 to 9700 lux and the PAR ranged from 102 to 147 $\mu\text{E}/\text{m}^2/\text{s}$. Since the light intensity was within the appropriate range, the PAR was not adjusted. This is not expected to have resulted in an impact on the study outcomes.
Acclimation period:	It is unclear how long stock cultures were maintained at the test facility prior to transferring fronds from the stock culture to fresh medium for use in the test. Given the strong growth of control plants, this is not expected to have resulted in an impact on the study outcomes.

E. REVIEWERS COMMENTS: Nothing additional

F. CONCLUSIONS: The study is acceptable. Based on this study, 3,5-di-OH-XDE-742 is practically non-toxic to duckweed, *Lemna gibba*.

EC50/IC50: >95 mg ac/L; NOEC: 37 mg ac/L (7 day frond number).

III. REFERENCES:

Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742 (Pyroxsulam) to aquatic vascular plants *Lemna gibba*

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

ASTM. 2002. Standard Guide for Conducting acute toxicity tests with fishes, macroinvertebrates and amphibians. Standard E729-96. American Society for Testing and Materials, 100 Barr Harbor Drive, West Conshohocken, PA 19426.

Gulley, D.D., Boetler, A.M. and Bergman, H.L. 1996 TOXSTAT® Release 3.5. University of Wyoming, Laramie, Wyoming. Horning, W.B. and C.I. Weber. 1985. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms. EPA/600/4-85/014.

Hoberg J, 2006. Study Profile Template (SPT) for 5,7-di-OH Metabolite of XDE-742 – Toxicity to Duckweed, *Lemna gibba*. Study No. 12550.6418.SPT. Springborn Smithers Laboratories, Massachusetts. 30 March 2006.

Horning, W.B. and C.I. Weber. 1985. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms. EPA/600/4-85/014. Environmental Monitoring and Support Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio.

OECD, 1998. OECD Series on Principles of Good Laboratory Practice and Compliance Monitoring. Number 1. OECD Principles on Good Laboratory Practice (as revised in 1997). Environment Directorate Chemicals Group and Management Committee. ENV/MC/CHEM(98)17. OECD Paris. France. 41 pp.

OECD, 2000. OECD Guideline for Testing of Chemicals. *Lemna* sp. Growth Inhibition Test. Proposed Guideline #221. Revised Draft, October 2000.

U.S. EPA, 1996. Office of Prevention, Pesticides and Toxic Substances. Ecological Effects Test Guideline, OPPTS 850.4400. Aquatic Plant Toxicity Test Using *Lemna* spp., Tiers I and II. "Public Draft" EPA 712-C-96-156 April 1996. U.S. Environmental Protection Agency. Washington, D.C.

U.S. EPA. 1989. Federal Insecticide, Fungicide and Rodenticide Act (FIFRA); Good Laboratory Practice Standards; Final Rule (40 CFR, Part 160); FR: 8/17/89; pp. 34052. U.S. Environmental Protection Agency, Washington, D.C.

Weber, C.I., W.H. Peltier, T.J. Norberg-King, W.B. Horning II, F.A. Kessler, J.R. Menkedick, T.W. Neihsel, P.A. Lewis, D.J. Klemm, Q.H. Pickering, E.L. Robinson, J.M. Lazorchak, L.J. Wymer and R.W. Freyberg (eds.). 1989. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms. 2nd ed. EPA/600/4/89/001. Environmental Monitoring Systems Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio.

Williams, D.A. 1971. A test for differences between treatment means when several dose levels are compared with a zero dose control. *Biometrics* 27: 103-117.

**Data Evaluation Report on the acute toxicity of 5,7-Di-OH metabolite of XDE-742
(Pyroxsulam) to aquatic vascular plants *Lemna gibba***

PMRA Submission Number 2006-4727; ID 1283257

EPA MRID 469084-35

Williams, D.A. 1972. A comparison of several dose levels with a zero control. *Biometrics* 28: 519-531.

Approved 04/01/01 C.K.