

US EPA ARCHIVE DOCUMENT

258018,-19
Record No.

22
Review No.

109303
SHAUGHNESSEY NO.

EEB REVIEW

DATE: IN 1/19/90 OUT JUN 11 1990

FILE OR REG. NO. 352-485,-515

PETITION OR EXP. NO. _____

DATE OF SUBMISSION 1-11-90

DATE RECEIVED BY EFED 1-18-90

RD REQUESTED COMPLETION DATA 3-19-90

EEB ESTIMATED COMPLETION DATE 3-19-90

RD ACTION CODE/TYPE OF REVIEW 331

TYPE PRODUCTS(S): I, D, H, F, N, R, S I

MRID NO(S). _____

PRODUCT MANAGER NO. G. LaRocca (15)

PRODUCT NAME(S) ASANA XL Insecticide

COMPANY NAME DuPont

SUBMISSION PURPOSE Review Amendment to add Rangeland and
Pasture Grass to Label

SHAUGHNESSEY NO.	CHEMICAL AND FORMULATION	% A.I.
<u>109303</u>	<u>Esfenvalerate</u>	_____
_____	_____	_____
_____	_____	_____

100.0 Chemical: ASANA

100.1 Submission: DuPont is requesting registration of ASANA XL Insecticide for use on rangeland and pasture grass.

100.2 Application Rate: A rate of 0.006-0.015 lb ai/A is proposed for rangeland and 0.03-0.05 lb ai/A for improved pasture grass. The maximum yearly rates are 0.04 lb ai/A for range and 0.10 lb ai/A for pasture.

100.4 Precautionary Labeling:

This pesticide is toxic to wildlife and extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

100.5 Hazard Assessment: The uses proposed by DuPont are large acreage that result in about 1 billion acres for rangeland and pastures.

101.1 Likelihood of Adverse Effects to Nontarget Organisms

Although the acute/chronic fish and wildlife data base for ASANA is not complete, studies have shown that this isomer of fenvalerate appears to have similar fate and toxicity parameters as the parent compound. Therefore, the Agency will rely upon the fenvalerate data base in evaluating the potential hazard of ASANA use to nontarget terrestrial and aquatic organisms.

Aquatic Toxicity

Fenvalerate, a second generation pyrethroid, degrades in soil with a half-life of six months and undergoes hydrolysis after 24 days at ph 7.2. Fenvalerate can strongly bind to sediment/particulate and result in a soil/water partition coefficient of greater than 15,000.

Fenvalerate is a neurotoxicant and effector of ion permeability (Miller and Adams 1982) and appears to interact with sodium gates (Lawrence and Casida 1983). Laboratory testing has shown that fenvalerate is very highly toxic to freshwater aquatic organisms as noted in acute toxicity

values that ranged from 0.032 ug/L (Daphnia magna) to 2.35 ug/L (fathead minnow) (Mayer and Ellersieck 1986). This very high toxicity has also been documented in acute marine studies. Schimmel et al. (1983) found that fenvalerate was acutely toxic to mysid shrimp, Mysidopsis bahia at 0.008 (0.005 - 0.01) ug/L and pink shrimp, Penaeus duorarum, at 0.84 (0.66 - 1.2) ug/L. They further found that acute toxicity values for estuarine fish ranged from 5.0 (0.55 - 5.3) ug/L for sheepshead minnow, Cyprinodon variegatus, and 0.31 (0.21 - 0.40) ug/L for Atlantic silversides, Menidia menidia.

An evaluation of sublethal fenvalerate exposure to aquatic invertebrate larval development and metabolism was conducted by McKenney and Hamaker (1984). They concluded that exposure to 0.0001 and 0.0002 ug/L can result in alterations in metabolic-salinity patterns of larval grass shrimp Palaemonetes pugio. This alteration reduces ecological fitness at a critical life stage by limiting the capacity of the organism to adapt to fluctuating salinity conditions that are normally encountered in estuarine waters.

The use of ASANA on rangeland and pastures represents substantial acreage, as well as, the potential for increase exposure to aquatic systems via runoff and/or drift. In order to evaluate potential exposure, the Ecological Effects Branch (EEB) has calculated estimated environmental concentrations (EEC) of ASANA residues following ground or aerial application (Appendix I, II). These calculations suggest that at the maximum rate of 0.05 lb ai/A for pasture grass, ASANA residues resulting from drift or runoff to an adjacent aquatic system can be expected to be 0.03 ug/L (ground) and 0.154 ug/L (aerial). Similar calculations for a maximum application to rangeland of 0.015 ug/L can result in 0.009 ug/L (ground) and 0.046 ug/L (aerial). A comparison of these estimated residue levels with toxicity values suggests that ASANA use may result in runoff or drift that exceeds aquatic toxicity concerns, especially to aquatic invertebrates. Smith et al. (1983) noted that fenvalerate concentration in runoff from a sugarcane-insect IPM system could present a toxicity problem to aquatic organisms. They concluded that although the toxicity of fenvalerate may be reduced as a result of sorption to sediment, this reduction was only 2-fold, and does not eliminate aquatic hazard.

Avian Toxicity

The available data suggests that although fenvalerate is practically non-toxic to birds at an acute level (Mallard LC₅₀ = 10,000 ppm), avian reproductive effects were found at 25 ppm. In assessing acute toxicity of ASANA to avian wildlife, EEB has estimated the potential exposure from

residues by using Hoerger and Kenaga (1972) table of typical maximum residues on differing categories of vegetation (Table 1).

Table 1. Maximum Expected Fenvalerate Residues on Avian Food and Dietary Intake (ppm) after an Application of 0.05 lb ai/A

<u>Food Type</u>	<u>Residues (ppm)</u>
Short Grass	14.0
Dense Foliage/ Small Insects	2.8
Large Insects	0.1

The maximum expected residues from the consumption of vegetation and insects (application rate of 0.05 lb ai/A) are expected to range from 0.1 to 14.0 ppm. These values show that ASANA use should not present a direct toxicity threat to birds, since, the expected residues are 6 to 3 orders of magnitude less than the avian toxicity values. However, because of the toxicity of ASANA to aquatic invertebrates, runoff of this pesticide may affect a significant trophic level that waterfowl are dependent upon, especially during breeding.

101.3 Endangered Species

The following EEC's and Endangered Species Triggers were calculated for ASANA.

<u>Organisms</u>	<u>Toxicity values</u>	<u>Trigger</u>	<u>Max. EEC</u>
Avian	LC ₅₀ =10,000 ppm	1/10 LC ₅₀ = 1000 ppm	14 ppm
Fish	LC ₅₀ =0.31 ug/L	1/20 LC ₅₀ = 0.015 ug/L	0.154 ug/L
Invert.	LC ₅₀ =0.032 ug/L	1/20 LC ₅₀ = 0.0016 ug/L	0.154 ug/L

The above EEC calculations suggest that ASANA exposure to endangered fish, aquatic invertebrates and terrestrial invertebrates exceeds the endangered species trigger of 1/20 LC₅₀ values. However, a full assessment of risk to endangered species can not be concluded at this time because pertinent ecological effects data (mesocosm) have not been submitted.

107.0 Conclusions.

EEB has partially completed an evaluation of the proposed ASANA Section 3 registration on rangelands and pastures. Expected environmental residues were calculated in order to assess the potential hazard of ASANA exposure to avian and aquatic organisms. These calculated residues appear to exceed acute/chronic toxicity values for fish and aquatic invertebrates by one to three orders of magnitude and suggest the potential for risk to aquatic organisms. The calculated residue levels for assessing ASANA exposure and toxicity to avians suggest that the proposed use patterns should not directly affect birds, but could result in a possible indirect effect to certain waterfowl by altering a significant aquatic invertebrate food base. However, at this time EEB can not complete a risk assessment of this chemical because relevant ecological effects data (mesocosm) have not been submitted for review. Until this assessment is completed, any expansion of the ASANA registration could possibly result in an increase risk the wildlife and aquatic organisms.

Michael Rexrode 5/30/90

Michael Rexrode, Fishery Biologist
Ecological Effects Branch
Environmental Fate and Effects Division

Ann Stavola 5/31/90

Ann Stavola, Section Head
Ecological Effects Branch
Environmental Fate and Effects Division

James Akerman 6/5/90

James Akerman, Chief
Ecological Effects Branch
Environmental Fate and Effects Division

Appendix I - EEC Calculations for ASANA Use on Pastures

I. Ground Application

Assumptions:

0.1% runoff
10 acre drainage basin
0.05 lb ai/A of ASANA

(A) Runoff

0.05 lb ai/A x 0.001 x 10 A = 0.0005 lbs ai total runoff
EEC of 1 lb ai, direct application to 1 A pond, 6 ft deep = 61
Therefore, EEC = $\frac{61 \text{ ug/L}}{1 \text{ lb ai}} \times \frac{0.0005 \text{ lb ai}}{1} = 0.03 \text{ ug/L}$

II. Aerial Application

Assumptions

0.1% runoff
60% application efficiency
10 acre drainage basin
5% drift
0.05 lb ai/A of ASANA

(A) Runoff

0.05 lb ai/A x 0.06 x 0.001 x 10 A = 0.00003 lb ai found in total runoff

(B) Drift

0.05 ai/A x 0.05 = 0.0025 lbs ai in total drift

Therefore, EEC = $\frac{61 \text{ ug/L}}{1 \text{ lb ai}} \times \frac{0.0025 \text{ lb ai}}{1} = 0.154 \text{ ug/L}$

Appendix II - EEC Calculations for ASANA Use on Rangelands

I. Ground Application

Assumptions:

0.1% runoff
10 acre drainage basin
0.015 lb ai/A of ASANA

(A) Runoff

$0.015 \text{ lb ai/A} \times 0.001 \times 10 \text{ A} = 0.00015 \text{ lbs ai total runoff}$
EEC of 1 lb ai, direct application to 1 A pond, 6 ft deep = 61
Therefore, $\text{EEC} = \frac{61 \text{ ug/L}}{1 \text{ lb ai}} \times \frac{0.00015 \text{ lb ai}}{1} = 0.009 \text{ ug/L}$

II. Aerial Application

Assumptions

0.1% runoff
60% application efficiency
10 acre drainage basin
5% drift
0.015 lb ai/A of ASANA

(A) Runoff

$0.015 \text{ lb ai/A} \times 0.6 \times 0.001 \times 10 \text{ A} = 0.00009 \text{ lb ai found in total runoff}$

(B) Drift

$0.015 \text{ ai/A} \times 0.05 = 0.0007 \text{ lbs ai in total drift}$

Therefore, $\text{EEC} = \frac{61 \text{ ug/L}}{1 \text{ lb ai}} \times \frac{0.0007 \text{ lb ai}}{1} = 0.046 \text{ ug/L}$

REFERENCES

Lawrence, L.J. and J.E. Casida. 1983. Stereospecific action of pyrethroid insecticides on the Y-aminobutyric acid receptor-ionophore complex. *Science* 221:1399-1401.

Mayer, F.L. and M.R. Ellersieck. 1986. Manual of Acute Toxicity: Interpretation and Data Base for 410 Chemicals and 66 Species of Freshwater Animals. U.S. Dept. of the Interior, Publication 160. p. 234-285.

McKenney, C. L. and D. B. Hamaker. 1984. Effects of fenvalerate on larval development of Palaemonetes pugio (Holthuis) and on larval metabolism during osmotic stress. *Aquat. Tox.* 5:343-355

Miller, T.A. and M.E. Adams 1982. Mode of action of pyrethroids. In Insecticide Mode of Action (J.R. Coats, ed.) pp. 3-24, Academic Press, New York.

Schimmel, S.C.; R.L. Garnas, J.M. Patrick and J.C. Moore. 1983. Acute toxicity, bioconcentration, and persistence of AC 222, 705, bentiocarb, chlorpyrifos, fenvalerate, methyl parathion, and permethrin in the estuarine environment. *J. Agric. Food Chem.* 31(1):104-113.

Smith, A.G., J.J. Stoudt and J.B. Gallop. 1964. Prairie potholes and marshes. In Waterfowl Tomorrow. U.S. Government Printing Office, Washington, D.C. 770 pp.

Swanson, G.A., G.L. Krapu and J.R. Serie. 1979. Foods of laying female dabbling ducks on the breeding grounds. In Waterfowl and Wetlands: Integrated Review. (T. Bookout, ed.) pp. 152.