

US EPA ARCHIVE DOCUMENT



9-22-92 470-A  
CASWELL FILE

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

SEP 22 1992

OFFICE OF  
PESTICIDES AND TOXIC  
SUBSTANCES

MEMORANDUM

**SUBJECT:** RfD for Iprodione

**TO:** George Ghali, Ph.D.  
Manager, RfD/Peer Review Committee  
Science Analysis Branch  
HED (H7509C)

**FROM:** K. Clark Swentzel *K. Clark Swentzel 9/16/92*  
Section Head, Section II  
Toxicology Branch II  
HED (H7509C)

**THROUGH:** Marcia van Gemert, Ph.D. *M van Gemert 9/16/92*  
Branch Chief, Toxicology Branch II  
HED (H7509C)

The RfD for iprodione is based on a 1-year feeding study in dogs (84RH002/179) in which the dietary levels were 0, 100, 600 or 3600 ppm. The registrant has repeated a 1-year study in which the dogs were fed dietary levels of 0, 200, 300, 400 or 600 ppm iprodione (EPA Memorandum, Swentzel to Rose/Lewis, June 24, 1992). The NOEL in the first study, on which the RfD is based, is 100 ppm (approximately 4 mg/kg/day); the NOEL in the current study is 400 ppm (approximately 18 mg/kg/day). The LEL for both studies is 600 ppm (increased number of RBCs with Heinz bodies and decreased prostate weights in the first study and depressed RBC parameters in the second).

