

US EPA ARCHIVE DOCUMENT

6-25-81  
RCB



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

~~GETCK~~  
~~Sanchez~~  
Fate petition

OFFICE OF  
PESTICIDES AND TOXIC SUBSTANCES

JUN 25 1981

JUN 25 1981

MEMORANDUM

DATE:

SUBJECT: Bayleton 25% W.P.; Tolerance of 0.1 ppm on Chick Peas. (Acc. # 099850-51, Caswell 862 AA, Petition # 1E2459).

FROM: George Z. Ghali, Ph.D.  
Toxicology Branch, HED (TS-769)

G. Ghali

TO: Henry Jacoby (PM. 21)  
Registration Division (TS-767)

R. H. 5-28-81  
Refer WZB

Registrant:

Mobay Chemical Corporation  
St. Louis, Missouri

Action Requested:

Establishment of a tolerance of 0.1 ppm of Bayleton residues on chick peas.

Conclusion and Recommendations:

Toxicology Branch recommends for the establishment of the proposed tolerance.

Background Information:

Existing Tolerances:

There are no existing tolerances for this pesticide. However, temporary tolerances were established as follows:

Apples, fresh	0.75 ppm
Pears, fresh	0.75 ppm
Grapes	1.00 ppm
Wheat	0.10 ppm
Barley	0.10 ppm

1/4

Toxicology Data:

(memo by John Doherty, 2/15/78)

A. Bayleton 25% W.P.

Acute oral LD<sub>50</sub>, rat, 2,828 mg/kg (male), 3,668 mg/kg (female)

Acute dermal LD<sub>50</sub>, rabbit, > 5,000 mg/kg

Acute inhalation LC<sub>50</sub>, rat > 20 mg/l

Primary eye irritation; reversible corneal opacity.

Primary skin irritation: not irritating.

B. Bayleton, technical:

(memo by J. Doherty 1/9/80, A. Arce 1/24/80)

1. Acute Studies:

Acute oral <sup>rats</sup> 568 mg/kg (male), 363 mg/kg (female).

Acute I.P. rats, LD<sub>50</sub> 293 and 321 mg/kg for females and males respectively.

Acute dermal, rats LD<sub>50</sub> > 1000 mg/kg.

Acute inhalation; mice, rabbits, hamsters and rats. LC<sub>50</sub> > 174 mg/m<sup>3</sup>.

Primary skin irritation: rabbits negative.

Skin irritation: rabbits negative.

Skin irritation; human. not irritant.

Eye irritation; invalid study, dose was not reported.

2. Mutagenicity:

Dominant lethal test, negative for mutagenicity.

Micronucleus test, negative for mutagenicity.

Ames test, negative at doses from 5 to 1000 ug/ml.

3. Subchronic toxicity:

Twelve-week feeding, rats. NOEL > 2000 ppm.

Thirteen-week feeding, dogs NOEL > 2400 ppm.

4. Subacute toxicity:

Thirty-day oral administration, rats, NOEL 3mg/kg (m), 10 mg/kg (f).

Four-hours inhalation, rats, LC<sub>50</sub> > 453 mg/m<sup>3</sup>.

Six-hours inhalation, rats 15 exposure, NOEL 78.7 mg/m<sup>3</sup>.

Cumulative subacute dermal application for four weeks, rabbits, NOEL 250 mg/kg

5. Embryotoxicity and Teratology:

In an oral administration study in rats, occasionally cleft palates were seen in the groups treated with 75 mg/kg/day and above. These equaled only 4 of the 211 of one experiment and 3 of 183 in another experiment. However, this deformity is seldom seen in this strain. A no-effect level for embryonic and fetal development/teratology was at least 50 mg/kg/day (J. Doherty, 1978).

In a later memo by Roger Gardner dated 4/16/81 it was concluded that the cleft palates observed in this study may not be attributable to Bayleton treatment. However, the memo also indicated that the raw data and background on the historical terata incidence in this strain of rats are needed to further evaluate the significance of this effect.

From the information available until now, the compound is questionably positive with a clear-cut no-effect level for teratogenic effect of 50 mg/kg/day.

Inhalation administration, rats, negative for terata and embryotoxicity of a dose level of 113.6 mg/m<sup>3</sup>.

Oral administration, rabbits, negative up to and including 50 mg/kg (highest dose tested).

6. Chronic Toxicity:

(memo by George Z. Ghali, 3/81)

Two-year feeding (oncogenicity) in rats; not onocgenic, NOEL 50 ppm.

Two-year feeding study in dogs, not oncogenic, NOEL 100 ppm.

Multigeneration reproduction study, rats NOEL 50 ppm.

Toxicology Data Gap:

An adequate metabolism study in an appropriate animal species.

Unverified printout

file last update: 3/24/81

not recorded  
3/24/81

ACCEPTABLE DAILY INTAKE DATA

WAT, Older	NOEL	S.F.	ADI	RPI
mg/kg	ppm		mg/kg/day	mg/day (60kg)
2.500	50.00	100	0.250	1.5000

Unpublished, tox Approved PP# 0G2300, 1G2432

CROP	Tolerance	Food Factor	mg/day (1.5kg)
Apples ( 2)	0.750	2.53	0.02846
Pears (116)	1.000	0.26	0.00383
Barley ( 8)	0.100	0.03	0.00005
wheat (170)	0.100	10.36	0.01554

RPI 1.5000 mg/day (60kg) THRC 0.0479 mg/day (1.5kg) % ADI 3.19

\*\*\*\*\*

Current Action PP# 0E2393, 1E2459, 0F2349

CROP	Tolerance	Food Factor	mg/day (1.5kg)
Cucumbers, not pickl ( 47)	0.100	0.34	0.00051
Tomatoes (163)	0.200	2.87	0.00862
Grapes, not raisins ( 67)	2.000	0.45	0.01349
Melons ( 92)	0.200	2.00	0.00601
<u>chick Peas</u> (214)	0.100	0.03	0.00005

RPI 1.5000 mg/day (60kg) THRC 0.0766 mg/day (1.5kg) % ADI 5.10

\*\*\*\*\*

Proposed based on 2-yr...