

US EPA ARCHIVE DOCUMENT

*Head - Sub File*

**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**

WASHINGTON, D.C. 20460

June 16, 1994

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

**MEMORANDUM**

**SUBJECT:** Metalaxyl (113501) - Case No. 0081.  
Product and Residue Chemistry Chapters for the  
Reregistration Eligibility Document (RED).  
CB No. 12906. DP Barcode: D197037.

**FROM:** Susan V. Hummel, Chemist *Susan V. Hummel*  
Special Review Section II  
Chemistry Branch II - Reregistration Support  
Health Effects Division (H7509C)

**THROUGH:** Francis B. Suhre, Section Head *Francis B. Suhre*  
Special Review Section II  
Chemistry Branch II - Reregistration Support  
Health Effects Division (H7509C)

**TO:** Debra Edwards, Chief  
Chemical Coordination Branch  
Health Effects Division (H7509C)

Attached are the Product and Residue Chemistry chapters for the Metalaxyl RED. These chapters were completed by Dynamac Corporation under the supervision of CBRS, HED. They have undergone secondary review in the branch and have been revised to reflect Agency policies.

All Product Chemistry data requirements for the TGAI are satisfied for the Ciba Geigy 90% T (EPA Reg. No. 100-601). Some additional physical and chemical characteristics are needed for the MP.

The chemical name of the metalaxyl metabolite included in the tolerance expression in 40 CFR 180.408 (a) and (c), 185.4000 (a), and 186.4000 (a) needs to be corrected. The correct chemical name is N-2-hydroxymethyl-6-methylphenyl)-N-methoxyacetyl)-alanine methyl ester. A final decision on the form of the tolerance expression for livestock commodities will be made after additional recovery data are submitted for several livestock metabolites.

According to REFS, a 50% wettable powder (WP) formulation of metalaxyl was recently registered for multiple foliar applications to several crops (EPA Reg. No. 100-735). Residue data are not available to support the use of multiple foliar and other late season postemergence applications (7-30 day PHI) of this type of



**Recycled/Recyclable**  
Printed with Soy/Canola Ink on paper that  
contains at least 50% recycled fiber

formulation. A limited set of side by side field trial data are needed for the WP formulation on asparagus, fruiting vegetables, and strawberries showing comparable residue levels for the WP and EC formulations.

Additional recovery data are needed for several metalaxyl hydroxy metabolites in livestock commodities to permit final calculations of residues in livestock commodities. Upper bound estimates of livestock residues have been made for the RED.

Storage stability data are still needed for processed commodities and livestock commodities. Data are needed for metalaxyl and representative metabolites.

Additional confined rotational crop studies and field rotational crop studies have been submitted and are under review.

For dietary exposure assessment, current and/or recommended tolerance levels should be used, except for livestock commodities. For livestock commodities, the upper bound residue levels calculated should be used. Since tolerance levels and upper bound residue estimates are being used for dietary exposure assessment, the dietary exposure to residues of metalaxyl is likely to be greatly overestimated.

If you need additional input, please advise.

Attachment 1 & 2: Metalaxyl Product and Residue Chemistry RED chapters (respectively)

cc (with Attachments 1 & 2): Circ, Reg.Std.F., SF, S. Hummel, SRRD, Dynamac

cc (without Attachments): RF

RDI:FBS:06/13/94:MM:06/16/94:EZ:06/16/94

H7509C:CBRS:SVH:svh:CM#2:RM804:06/16/94:METALRED.946

**DYNAMAC**  
**CORPORATION**  
*Environmental Services*

---

Final Report

**METALAXYL**  
**Shaughnessy No. 113501**  
**Case No. 0081**  
**(CBRS Nos. 12906 & 12907, DP**  
**Barcodes D197037 & D197066)**

**TASK 2A**  
**Reregistration Eligibility Document:**  
**Product Chemistry Considerations**

May 5, 1994

**Contract No. 68-D2-0053**


**Submitted to:**  
U.S. Environmental Protection Agency  
Arlington, VA 22202

**Submitted by:**  
Dynamac Corporation  
The Dynamac Building  
2275 Research Boulevard  
Rockville, MD 20850-3268

---

**METALAXYL****REREGISTRATION ELIGIBILITY DOCUMENT:****PRODUCT CHEMISTRY CONSIDERATIONS****(Shaughnessy No. 113501; Case No. 0081)****CBRS Nos. 12906 and 12907; DP Barcodes D197037 and D197066****TASK 2A****DESCRIPTION OF CHEMICAL**

Metalaxyl [N-(2,6-dimethylphenyl)-N-(methoxyacetyl)-alanine methyl ester] is a systemic fungicide registered for use on a variety of crops/sites (primarily citrus, cotton, cucurbits, onions, ornamentals, potatoes, soybeans (seed treatment), tobacco, tomatoes, and turf).



Empirical Formula:	$C_{15}H_{21}NO_4$
Molecular Weight:	279.3
CAS Registry No.:	57837-19-1
Shaughnessy No.:	113501

**IDENTIFICATION OF ACTIVE INGREDIENT**

Technical metalaxyl is a white to beige crystalline solid with a melting point of 71-72 C and a vapor pressure of  $2.2 \times 10^{-6}$  mm Hg at 20 C. The solubility of metalaxyl in water at 20 C is 7.1 g/L. Metalaxyl is also readily soluble in most organic solvents (i.e., 65% soluble in methanol and 55% soluble in benzene).

**MANUFACTURING-USE PRODUCTS**

A search of the Reference File System (REFS) conducted 3/22/94 identified a single metalaxyl manufacturing-use product (MP) registered to Ciba-Geigy Chemical Company,

the 90% technical (T; EPA Reg. No. 100-601). This product is the only MP subject to a reregistration eligibility decision.

### REGULATORY BACKGROUND

The 12/81 Metalaxyl Guidance Document was developed in conjunction with the initial registration of the pesticide, which consisted of non-food uses only. Because metalaxyl was subsequently registered for numerous food/feed uses, the data addressed in the original Guidance Document were reevaluated and a second Guidance Document was issued 9/88. Additional data were required for the following GLNs: 61-2, 61-3, 62-1, 62-2, and 62-3. Data submitted in response to the 12/88 Guidance Document were evaluated in the Metalaxyl Reregistration Standard Update dated 3/13/91; additional data were required for GLNs 61-1, 61-2, 61-3, and 62-2. Although physical/chemical data from the 12/81 Guidance Document are considered to be adequate to fulfill data requirements for the TGAI, data pertaining to the following MP GLNs have not been submitted for the 90% T: 63-14, 63-16, 63-17, and 63-20.

The current status of the product chemistry data requirements for the Ciba-Geigy metalaxyl technical is presented in the attached data summary table. Refer to this table for a listing of the outstanding product chemistry data requirements. The nitrosamine content of technical metalaxyl is not expected to be of dietary concern because nitrosamines were detected at <1 ppm.

### CONCLUSIONS

All of the product chemistry data requirements pertaining to the TGAI are satisfied for the Ciba-Geigy 90% T (EPA Reg. No. 100-601); however, some physical/chemical characteristics of the MP remain outstanding. Provided that the registrant either certifies that the suppliers of starting materials and the manufacturing process for the metalaxyl technical product have not changed since the last comprehensive product chemistry review or submits a complete updated product chemistry data package, CBRS has no objections to the reregistration of metalaxyl with respect to product chemistry data requirements.

AGENCY MEMORANDA CITED IN THIS DOCUMENT

CB No(s): 8166  
 DP Barcode(s): D165989  
 Subject: Metalaxyl. Ciba-Geigy Response to the Reregistration Standard: Product Chemistry Data (MRID No. 41912901), Storage Interval Data (MRID No. 41912902), and Corn Fodder Data (MRID No. 41912903).  
 From: J. Abbotts  
 To: L. Rossi  
 Dated: 04/16/92  
 MRID(s): 41912901

CB No(s): 10303 and 9962  
 DP Barcode(s): D181027 and D178783  
 Subject: Metalaxyl. Ciba-Geigy Product Chemistry of Selected Impurities in Tox-Tested Samples vs. "Current" [1982] and Proposed CSFs, and for 61-1, 62-1, 62-2, and 62-3.  
 From: K. Dockter  
 To: L. Rossi and N. Thoa  
 Dated: 9/29/92  
 MRID(s): 42319901, 42409201, and 42409202

CB No(s): 11937  
 DP Barcode(s): D191625  
 Subject: Metalaxyl Reregistration. Ciba's 5/4/93 Response [CSF & Nitrosamine Data for Technical; 100-601] to Our 9/29/92 Review & Related 3/23 & 30/93 Telecons.  
 From: K. Dockter  
 To: P. Perreault  
 Dated: 6/30/93  
 MRID(s): 42762801 and 42762802

## PRODUCT CHEMISTRY CITATIONS

Bibliographic citations include only MRIDs containing data which fulfill data requirements.

### References (cited):

00104483 Ciba-Geigy Corp. (1978) [Chemical Data on the Fungicide Ridomil]. (Compilation; unpublished study received Jul 13, 1978 under 100-EX-63; CDL:234427-A).

00104498 Ciba-Geigy Corp. (1978) [Chemistry of CGA-48988 Technical]. (Compilation; unpublished study received Sep 5, 1978 under 100-601; CDL:235062-A).

40435001 Ciba Geigy Corp. (1982) Product Chemistry for Product Containing Metalaxyl. Unpublished study. 4 p.

41055201 Lail, L. (1989) Technical Metalaxyl: Product Chemistry: Study No. PC-88-005. Unpublished study prepared by Ciba-Geigy Corp. 51 p.

41055202 Lail, L. (1989) Technical Metalaxyl: Product Chemistry: Study No. PC-88-005. Unpublished study prepared by Ciba-Geigy Corp. 120 p.

41912901 Lail, L. (1989) Technical Metalaxyl: A Supplement to the Product Chemistry. Unpublished study prepared by Ciba-Geigy Corp. 31 p.

42319901 Lail, L. (1992) Technical Metalaxyl Supplement to Product Chemistry. Unpublished study prepared by Ciba-Geigy Corp. 18 p.

42409201 Lail, L. (1989) Technical Metalaxyl: Supplement to Product Chemistry. Unpublished study prepared by Ciba-Geigy Corp. 7 p.

42409202 Lail, L. (1989) Technical Metalaxyl: Supplement to Product Chemistry. Unpublished study prepared by Ciba-Geigy Corp. 7 p.

42762801 Lail, L. (1992) Technical Metalaxyl: Supplement to Product Chemistry. Unpublished study prepared by Ciba Plant Protection. 8 p.

42762802 Lail, L. (1992) Technical Metalaxyl: Supplement to Product Chemistry. Unpublished study prepared by Ciba Plant Protection. 28 p.

Case No. 0081  
 Chemical No. 113501

Case Name: Metalaxyl  
 Registrant: Ciba-Geigy Corporation  
 Product(s): 90% T (EPA Reg. No. 100-601)

**PRODUCT CHEMISTRY DATA SUMMARY**

Guideline Number	Requirement	Are Data Requirements Fulfilled? <sup>a</sup>	MRID Number <sup>b</sup>
61-1	Product Identity and Disclosure of Ingredients	Y	00104498 41055201 41912901 42409201 <sup>c</sup> 42762801 <sup>d</sup> 42762802 <sup>d</sup> CSF dated 5/2/93 <sup>d</sup>
61-2	Starting Materials and Manufacturing Process	Y	41055201 41912901
61-3	Discussion of Formation of Impurities	Y	41055201 41055202 41912901
62-1	Preliminary Analysis	Y	41055202 42319901 <sup>e</sup> 42762801 <sup>d</sup> 42762802 <sup>d</sup>
62-2	Certification of Ingredient Limits	Y	41055201 41055202 41912901 42409202 <sup>c</sup>
62-3	Analytical Methods to Verify the Certified Limits	Y	41055202 41912901
63-2	Color	Y	00104483
63-3	Physical State	Y	00104483
63-4	Odor	Y	00104483
63-5	Melting Point	Y	00104483
63-6	Boiling Point	N/A <sup>e</sup>	
63-7	Density, Bulk Density or Specific Gravity	Y	00104483
63-8	Solubility	Y	00104483
63-9	Vapor Pressure	Y	00104483
63-10	Dissociation Constant	Y	00104483
63-11	Octanol/Water Partition Coefficient	Y	40435001
63-12	pH	Y	40435001
63-13	Stability	Y	00104483
63-14	Oxidizing or Reducing Action	N	
63-15	Flammability	N/A <sup>e</sup>	
63-16	Explosibility	N	
63-17	Storage Stability	N	
63-18	Viscosity	N/A <sup>e</sup>	
63-19	Miscibility	N/A <sup>e</sup>	
63-20	Corrosion Characteristics	N	

<sup>a</sup> Y = Yes; N = No; N/A = Not Applicable.

<sup>b</sup> **Bolded** references were reevaluated/reviewed in the Metalaxyl Guidance Document dated 9/88; underlined references were reviewed in the Metalaxyl Reregistration Standard Update dated 3/13/91; *italicized* references were reviewed under CB No. 8166, D165989, dated 4/16/92, by J. Abbotts; remaining references reviewed as noted.

<sup>c</sup> CB Nos. 10303 and 9962, D181027 and D178783, dated 9/29/92, by K. Dockter

<sup>d</sup> CB No. 11937, D191625, dated 6/30/93, by K. Dockter.

<sup>e</sup> Data are not required because the TGAI/MP is a solid at room temperature.

r

**DYNAMAC**  
**CORPORATION**  
*Environmental Services*

---

Final Report

**METALAXYL**  
**Shaughnessy No. 113501**  
**Case No. 0081**  
**(CB No. 12906, DP Barcode D197037)**

**TASK 2B**  
**Reregistration Eligibility Document:**  
**Residue Chemistry Considerations**

June 15, 1994

**Contract No. 68-D2-0053.**

**Submitted to:**  
U.S. Environmental Protection Agency  
Arlington, VA 22202

**Submitted by:**  
Dynamac Corporation  
The Dynamac Building  
2275 Research Boulevard  
Rockville, MD 20850-3288

---

## METALAXYL

REREGISTRATION ELIGIBILITY DOCUMENTRESIDUE CHEMISTRY CONSIDERATIONSShaughnessy No. 113501; Case 0081(CB No. 12906; DP Barcode D197037)

TABLE OF CONTENTS		page
INTRODUCTION . . . . .		1
REGULATORY BACKGROUND . . . . .		1
SUMMARY OF SCIENCE FINDINGS . . . . .		5
GLN 171-3: Directions for Use . . . . .		5
GLN 171-4 (a): Plant Metabolism . . . . .		6
GLN 171-4 (b): Animal Metabolism . . . . .		6
GLN 171-4 (c) and (d): Residue Analytical Methods - Plants and Animals . . . . .		7
GLN 171-4 (e): Storage Stability . . . . .		7
GLN 171-4 (k): Magnitude of the Residue in Plants . . . . .		8
GLN 171-4 (l): Magnitude of the Residue in Processed Food/Feed . . . . .		8
GLN 171-4 (j): Magnitude of the Residue in Meat, Milk, Poultry, and Eggs . . . . .		9
GLN 165-1: Confined Rotational Crops . . . . .		9
GLN 165-2: Field Rotational Crops . . . . .		9
TOLERANCE REASSESSMENT SUMMARY . . . . .		63
Tolerances Listed Under 40 CFR § 180.408 (a) . . . . .		63
Tolerances Listed Under 40 CFR § 180.408(b) . . . . .		64
Tolerances Listed Under 40 CFR § 180.408(c) . . . . .		64
Tolerances Listed Under 40 CFR § 185.4000(a) . . . . .		64
Tolerances Listed Under 40 CFR § 185.4000(b) . . . . .		65
Tolerances Listed Under 40 CFR § 185.4000(d) . . . . .		65
Tolerances Listed Under 40 CFR § 186.4000(a) . . . . .		65
Tolerances Listed Under 40 CFR § 186.4000(b) . . . . .		66
Tolerances Listed Under 40 CFR § 186.4000(d) . . . . .		66
CODEX HARMONIZATION . . . . .		73
DIETARY EXPOSURE ASSESSMENT SUMMARY . . . . .		74
AGENCY MEMORANDA CITED IN THIS DOCUMENT . . . . .		77
MASTER RECORD IDENTIFICATION NUMBERS . . . . .		86

# METALAXYL

## REREGISTRATION ELIGIBILITY DOCUMENT

### RESIDUE CHEMISTRY CONSIDERATIONS

Shaughnessy No. 113501; Case 0081

(CBRS Nos. 12906 and 12907; DP Barcodes D197037 and D197066)

#### TASK 2B

### INTRODUCTION

Metalaxyl [N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester] is a fungicide registered for use as a seed treatment, soil application, and/or foliar application on a variety of food and feed crops, including: alfalfa, almonds, apples, artichokes (Jerusalem), asparagus, avocados, beets (table/garden), blueberries, broccoli, cabbage, carrots, cassava, cauliflower, cereal grains, chicory, Chinese broccoli (gai lon), Chinese cabbage (tight heading varieties), citrus fruits, corn, cotton, cranberries, cucurbit vegetables, dill, eggplants, forage grasses, forage legumes, ginger, ginseng, grapes, hops, horseradish, legume vegetables, lettuce, okra, onions (bulb and green), papaya, parsnips, peanuts, peppers, pineapples, potatoes, radishes, raspberries, rice, rutabaga, salify, soybeans, spinach, stone fruits, strawberries, sugar beets, sweet potato, tomato, turnips, walnuts, and yams. The metalaxyl formulations registered for soil and foliar application include granular (G), wettable powder (WP), and emulsifiable concentrate (EC) formulations. Formulations registered for use as seed treatments include WPs and a flowable concentrate (FIC).

Tolerances for residues of metalaxyl in/on raw and processed plant commodities and animal commodities are currently expressed in terms of the combined residues of metalaxyl [N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester] and its metabolites containing the 2,6-dimethylaniline moiety, and N-(2-hydroxymethyl-6-methyl)-N-(methoxyacetyl) alanine methyl ester or N-(2-hydroxymethyl-6-methylphenyl)-N-(methoxyacetyl) alanine methyl ester [*Source: 40 CFR 180.408(a), (b), and (c), 40 CFR 185.4000, and 40 CFR 186.4000*]. The correct chemical nomenclature for the metalaxyl metabolite included in the tolerance expression is N-(2-hydroxymethyl-6-methylphenyl)-N-(methoxyacetyl) alanine methyl ester. The metalaxyl tolerance expressions in 40 CFR 180.408 (a) and (c), 105.4000 (a), 186.4000 (a) should be corrected.

### REGULATORY BACKGROUND

Metalaxyl was the subject of a Reregistration Standard and Guidance Document 12/81 and a Final Registration Standard and Tolerance Reassessment (FRSTR), Residue Chemistry Chapter dated 6/22/87 and Guidance Document dated 9/88. The Metalaxyl Reregistration

Update is dated 3/13/91. The information contained in this document outlines the Residue Chemistry Science Assessments with respect to the reregistration of metalaxyl.

Metabolites which may not be covered by the current tolerance expression have been identified in livestock metabolism studies. Up to 34% of the residue in ruminants and poultry are compounds containing the 2-hydroxymethyl-6-methyl aniline moiety other than N-(2-hydroxymethyl-6-methyl)-N-methoxyacetyl alanine methyl ester, which is currently regulated. The HED Metabolism Committee (9/10/93) has determined that the residues to be regulated in livestock commodities are metalaxyl, metabolites that can be converted to 2,6-dimethyl aniline (2,6-DMA), and those containing the 2-hydroxymethyl-6-methyl aniline (HMMA) moiety. However, additional data are required to demonstrate the recovery of HMMA-containing metabolites as 2,6-dimethyl aniline by the livestock tolerance enforcement method. The wording of the tolerance expression will depend on the recovery of HMMA containing metabolites using the current enforcement method. These data are outstanding, but considered confirmatory, since sufficient information is available to do a reasonable upper bound dietary exposure assessment. Residues of concern not recovered by enforcement methodology, and therefore not included in the tolerance expression, are included in the anticipated residues provided in this document. The chemical structures of the metabolites of concern are presented in Figure A.

Figure A. The chemical structure of metalaxyl and its metabolites of concern.



Structure Metabolite: Chemical name	Structure Metabolite: Chemical name
 <p><b>metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester</b></p>	 <p><b>N-(2,6-dimethylphenyl)-N-(hydroxyacetyl)alanine methyl ester</b></p>

Figure A (continued).













Structure Metabolite: Chemical name	Structure Metabolite: Chemical name
 <p data-bbox="240 659 581 688">N-(oxalyl)-2,6-dimethylaniline</p>	 <p data-bbox="857 659 1409 716">N-(oxalyl)-N-(2,6-dimethylphenyl)alanine methyl ester</p>
 <p data-bbox="240 1058 732 1087">N-(2,6-dimethylphenyl)-2-hydroxyacetamide</p>	 <p data-bbox="857 1058 1430 1087">N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine</p>
 <p data-bbox="240 1430 808 1486">N-[2-hydroxymethyl]-6-methylphenyl]-N-(methoxyacetyl)alanine methyl ester</p>	 <p data-bbox="857 1430 1409 1459">N-(2,6-dimethylphenyl)-N-(hydroxyacetyl)alanine</p>

Figure A (continued).

Structure Metabolite: Chemical name	Structure Metabolite: Chemical name
 <p>N-(oxalyl)-N-(2,6-dimethylphenyl)alanine</p>	 <p>N-decanoic acid ester of N-(2,6-dimethylphenyl)-N-(hydroxyacetyl)alanine methyl ester</p>
 <p>Sulfuric acid ester of N-[2-(hydroxymethyl)-6-methylphenyl]-N-(methoxyacetyl)alanine methyl ester</p>	 <p>N-[2-(hydroxymethyl)-6-methylphenyl]-N-(hydroxyacetyl)alanine</p>
 <p>N-Fatty acid amide conjugates of N-[2-(hydroxymethyl)-6-methylphenyl]alanine</p>	 <p>N-[2-(hydroxymethyl)-6-methylphenyl]-N-(methoxyacetyl)alanine</p>

SUMMARY OF SCIENCE FINDINGS

GLN 171-3: Directions for Use: A REFs search conducted 3/22/94 revealed that there are 18 end-use products (Eps) of metalaxyl presently registered to Ciba-Geigy Corporation which may be used on food/feed crops. These Eps are presented below.

Registrant EPA Reg. No.	Acceptance Date	Formulation Class	Product Name
<b>Ciba-Geigy Corporation</b>			
100-664	1/25/94	1% G	Ridomil PC® 11G Granular Fungicide
100-713	1/25/94	1% G	Ridomil PC® Granular Fungicide
100-676	1/15/90 <sup>a</sup>	2% G	Subdue® Granular Fungicide
100-628 <sup>b</sup>	12/2/93	5% G	Ridomil® 5G Fungicide
100-646	1/15/90 <sup>a</sup>	5% G	Subdue® 5G Fungicide
100-749	3/4/94	7% WP	Ridomil® MZ Fungicide
100-658	1/31/94	9% WP	Ridomil®/Bravo® 81W Fungicide
100-629 <sup>c</sup>	11/19/92	10% WP	Ridomil® MZ58 Fungicide
100-720	1/31/94	10% WP	Ridomil®/Copper 70W Fungicide
100-639	10/19/92	25% WP	Apron® 25W Fungicide
100-717	1/14/94	25% WP	Subdue® II Fungicide
Not available <sup>d</sup>	--	35% WP	Apron® 70 SD Fungicide
Not available <sup>e</sup>	--	35% WP	Apron® 35 SD Fungicide
100-672	1/15/90 <sup>a</sup>	45% WP	Apron® T69 Fungicide
Not available <sup>f</sup>	--	45% WP	Apron® T69 SD Fungicide
100-735	5/3/94	50% WP	Ridomil® 50W Fungicide
100-738	2/26/93	50% WP	Apron® 50W Fungicide
100-607 <sup>g</sup>	12/7/93	2 lb/gal EC	Ridomil® 2E Fungicide
100-619	12/7/93	2 lb/gal EC	Subdue® 2E Fungicide
100-626	9/18/89	2 lb/gal EC	Apron® 2E Fungicide
100-684	6/8/89	3 lb/gal FIC	Apron® Flowable Fungicide

<sup>a</sup> Copy of label was obtained from a Product Label DCI dated 1/15/90.

<sup>b</sup> Includes SLN Nos. AR90000502 and OK90000500.

<sup>c</sup> Includes SLN Nos. ME92000200, ND92000200, and WA92004200.

<sup>d</sup> Includes SLN Nos. CA85007300, ID83003300, and WA83003500.

<sup>e</sup> Includes SLN No. WA90003300.

<sup>f</sup> Includes SLN Nos. CA87000700, ID86001900, and WA86002800.

<sup>g</sup> Includes SLN Nos. AR90000501, AZ85000800, AZ86001500, CA82002400, CA85006800, CA86001800, CA90004700, CO88000100, FL86000200, FL89002300, ID89000500, LA92000400, OK90000500, OR89000800, TX91001100, WA87000200, and WA89001500.

A comprehensive summary of the registered food/feed use patterns of metalaxyl, based on these product labels, is presented in Table A. The 50% wettable powder (WP) formulation (EPA Reg. No. 100-735) was registered recently without review by the Chemistry Branches. The label permits the first post emergence of a EP formulation on a number of crops. Multiple foliar and other post-emergence applications are permitted throughout the growing season. These uses are not supported by residue data. A tabular summary of the residue chemistry science assessments for reregistration of metalaxyl is presented in Table B. The conclusions listed in Table B regarding the reregistration eligibility of metalaxyl food/feed uses are based on the use patterns registered by the basic producer, Ciba-Geigy Corporation. When end-use product DCIs are developed (e.g., at issuance of the RED), RD should require that all end-use product labels (e.g., MAI labels, SLNs, and products subject to the generic data exemption) be amended such that they are consistent with the basic producer labels.

**GLN 171-4 (a): Plant Metabolism:** The qualitative nature of the residue in plants is adequately understood. Studies with potatoes, lettuce, grapes, and tobacco indicate that metalaxyl is taken up, translocated, and extensively metabolized by plants. Metabolism involves oxidation of the ring-methyl group and hydrolysis of the methyl ester and methyl ether bonds; metabolites can be conjugated to glucose. Studies with [<sup>14</sup>C]metalaxyl-treated seed indicate that no appreciable residue was transferred from treated seed to edible mature plant parts. The residues to be regulated in plant commodities are those in the current tolerance expression, which include metalaxyl, metabolites that can be converted to 2,6-dimethyl aniline (2,6-DMA), and one metabolite containing the 2-hydroxymethyl-6-methyl aniline (HMMA) moiety, N-(2-hydroxymethyl-6-methylphenyl)-N-(methoxyacetyl) alanine methyl ester. The chemical name of the hydroxy metabolite should be corrected in 40 CFR 180.408 (a) and (c), 185.4000 (a), and 186.4000 (a).

**GLN 171-4 (b): Animal Metabolism:** The qualitative nature of the residue in animals is adequately understood. Metabolism in ruminants is via hydrolysis to the ester alcohol and acid alcohol, which may then be de-alkylated. In ruminants, metalaxyl may also be oxidized to benzylic alcohol or phenolic compounds. Some breakdown products may be conjugated with glucuronic acid. A major residue in milk is a mixture of the N-octanoic- and N-decanoic acid ester of metabolite CGA-67869 [(N-(2,6-dimethylphenyl)-N-(hydroxyacetyl)-aniline methyl ester]. 2,6-DMA-based residues accounted for up to ~50% of the residues in ruminant tissues. HMMA containing compounds comprised 34% of the residue in kidney and 12-14% in muscle and fat. The parent compound is rapidly excreted.

In poultry, metalaxyl is hydrolyzed to either the benzyl alcohol or the ester alcohol. The benzylic alcohol is converted to a sulfate and the ester alcohol is conjugated with fatty acid or converted to the acid alcohol, which is subsequently hydrolyzed to the benzylic alcohol. The predominant residues in poultry are the disubstituted free acid forms of the hydroxy metabolite CGA-94689 isomers (~47% in thigh muscle and 24% in egg), the sulfuric acid conjugate of the hydroxy metabolite (30% in thigh muscle), fatty acid conjugates of the disubstituted free acids of CGA-94689 (27% in fat), and CGA-107955, N-(2,6-dimethylphenyl)-N-hydroxyacetyl)alanine, (40% in fat).

The HED Metabolism Committee (9/10/93) has determined that the residues to be regulated are metalaxyl, metabolites that can be converted to 2,6-dimethyl aniline (2,6-DMA), and those containing the 2-hydroxymethyl-6-methyl aniline (HMMA) moiety. Some metabolites containing the HMMA moiety may be convertible to 2,6-DMA and thus measured by the current enforcement methodology. Data showing percent conversion of HMMA metabolites to DMA are outstanding (enforcement method recovery data for HMMA metabolites). The wording of the tolerance expression for livestock commodities will be determined after these data are submitted. Residue levels estimated in this RED chapter for livestock commodities represent an upper bound for HMMA containing metabolites.

GLN 171-4 (c) and (d): Residue Analytical Methods - Plants and Animals: Adequate enforcement methods are available for the determination of residues of metalaxyl and regulated metabolites in plants. Methods I and II in PAM, Vol. II correspond to Methods AG-348 and AG-349. Method AG-395, an improved modification of AG-348, has undergone successful Agency validation with plant matrices.

Metabolites which may not be covered by the current tolerance expression have been identified in livestock commodities. Additional validation data are needed to demonstrate that Method AG-576 adequately recovers metabolites containing the 2,6-DMA moiety and HMMA containing metabolites as 2,6-dimethyl aniline. Method AG-576 is a combination of Method II in PAM, Vol. II and AG-395, both of which have undergone successful Agency validation, and adequately recovers metalaxyl, per se. Method AG-576 may be adequate for enforcement of tolerances in animal commodities, if it is proven to adequately recover all residues of concern. If significantly less than 100% of the total toxic residue is recovered using Method AG-576, then residue levels used in future risk assessments may need to be adjusted to account for this.

Metalaxyl, per se, is completely recovered (> 80%) using FDA Multiresidue Protocol D (PAM, Vol. I Section 232.4), and is not recovered by Protocol E (PAM, Vol. I Sections 211.1/231.1 and 212.1/232.1) [Source: *PESTDATA, PAM, Vol. 1, Appendix, 8/93*]. Multiresidue data for representative metalaxyl metabolites have been forwarded to FDA.

GLN 171-4 (e): Storage Stability: Adequate storage stability data are available to support residue studies on raw agricultural commodities stored less than 24 months at temperatures of -20 C or less, with the exception of oilseed and grain crops. An interim storage stability study on cranberries, potatoes, peppers, and spinach indicate that weathered residues determined as 2,6-DMA are stable in samples stored (-20 C) for 18-24 months. A final report reflecting 38 months storage of these matrices is expected to be confirmatory. Storage stability data for metalaxyl and representative metabolites in a representative oilseed (e.g., soybean or nut) and grain (e.g., wheat) are required.

No storage stability data are available to support processing studies. Storage stability data for metalaxyl and representative metabolites are required from all processed commodities of an oilseed, a grain, and a fruit or fruiting vegetable.

Storage stability data for metalaxyl and representative metabolites in livestock commodities representing one year of frozen storage are required. Considering the demonstrated storage stability on several crops, the outstanding storage stability data are considered confirmatory.

GLN 171-4 (k): Magnitude of the Residue in Plants: All data requirements for magnitude of the residue in plants have been evaluated and deemed adequate, except for the 50% WP formulation. For other formulations, field trials were performed representing the various conditions under which the pesticide can be applied. Geographical representation is adequate and a sufficient number of trials reflecting representative formulation classes were conducted.

To support multiple foliar applications of the 50% WP formulation (EPA Reg. No. 100-735) to strawberries, fruiting vegetables, and asparagus, the registrant must either (1) submit available bridging data for representative commodities which show that residues are similar for late season, foliar applications of WP and EC formulations of metalaxyl at rates similar to those registered for these crops, or (2) perform 3 side by side field trials each for asparagus, strawberries, and tomatoes (9 total trials) in geographically representative growing areas using the 50% WP formulation and a representative EC formulation. If residues resulting from application of the 50% WP are substantially higher than those using the EC formulation, then additional field trial data may be required. These data are considered confirmatory.

CBRS recommends that a tolerance for forage, fodder, and straw for the cereal grains group (excluding wheat, barley, and oats) be established.

GLN 171-4 (l): Magnitude of the Residue in Processed Food/Feed: Processing studies have been conducted on the following RACs: apples, citrus, cereal grains, cottonseeds, grapes, hops, legume vegetables, peanuts, pineapples, potatoes, prunes, sunflowers, soybeans, sugar beets, and tomatoes. All data requirements for magnitude of the residue data in processed food/feed commodities have been deemed adequate to determine the extent to which residues of metalaxyl concentrate upon processing of the raw agricultural commodity.

Residues concentrate in the following food commodities: citrus oil; potato granules/flakes, and chips; prunes; raisins; sugar beet molasses, and tomato puree.

Residues concentrate in the following processed feed items: apple pomace (wet); citrus, molasses and pulp; grape pomace (wet and dried), grape raisin waste; legume vegetable cannery waste; peanut meal; pineapple, process residue, potato peels (dried); soybean hulls and meal; sugar beet molasses, sunflower meal; and tomato pomace (dried).

Food/feed additive tolerances for potato granule/flakes, potato chips, and tomato pomace must be proposed.

Because CBRS now considers cotton gin byproducts and tomato paste processed food/feed commodities, CBRS now requires residue data on these matrices. The required data are considered confirmatory.

GLN 171-4 (j): Magnitude of the Residue in Meat, Milk, Poultry, and Eggs: Tolerances for the combined residues of metalaxyl and its metabolites convertible to 2,6-dimethyl aniline and N-(2-hydroxymethyl-6-methylphenyl)-N-(methoxyacetyl) alanine methyl ester have been established for meat, fat, kidney, liver, and meat byproducts of cattle, goats, horses, sheep and poultry and for milk and eggs. The evaluation of the submitted ruminant feeding studies is deferred until the adequacy of the analytical method used is determined.

The results from the submitted poultry feeding study indicate that residues of metalaxyl will not exceed the established tolerances in poultry tissues and eggs. However, judgement of the adequacy of the available data is reserved until the analytical method used is determined to be adequate and adequate storage stability data are provided.

There are no established or proposed direct animal uses for metalaxyl. The existing ruminant and poultry feeding studies are adequate for risk assessment.

GLN 165-1: Confined Rotational Crops: The registrant has submitted three confined rotational crop studies (MRID Nos. 42196501 through 42196503), which are under review.

GLN 165-2: Field Rotational Crops: Tolerances have been established for the indirect or inadvertent residues of metalaxyl in rotational crops resulting from the application of metalaxyl to the primary crops. A field rotational crop study (MRID No. 41870308) has been submitted and is under review.

Table A. Use patterns subject to reregistration for: Case 0081, Metalaxyl.

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Agricultural Crop Uses</b>						
<b>Alfalfa</b>						
Soil broadcast application At-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	0.5 lb/A	(1)	Not applicable (NA)	12	Application may be made in a minimum of 20 gal/A. A 60-day PFI and PHI have been established. A 12-month plant-back interval has been established for corn and annual crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
<b>Almond</b>						
Soil band or broadcast application After planting (new plantings) or prior to spring plant growth (established plantings) Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	4 lb/A	3	60	12	The grazing of livestock in treated areas and the grazing or feeding of cover crops grown in treated orchards are prohibited.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Apple (bearing and nonbearing)</b>						
Soil broadcast or band application Early spring (prior to tree growth) or fall (postharvest) Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	4 lb/A	(2)	Not specified (NS)	12	Grazing or feeding of cover crops in treated orchards is prohibited.
Soil drench application Early spring (prior to tree growth) or fall (postharvest) Ground equipment		0.5 lb/100 gal				
<b>Asparagus</b>						
Broadcast application Postplant, before first cutting, or preharvest Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	1 lb/A	(2)	NS	12	Application may be made in a minimum of 10 gal/A.
<b>Avocado</b>						
Soil application (under sprinkler or drip irrigation) At the beginning of the growing season or at transplanting Ground equipment	5% G [100-628]	0.075 oz/sq. yd or 1.4 oz/tree ( > 15 ft. diameter tree canopy)	3	90	12	A 28-day PHI has been established. A maximum yearly application rate of 12 lb ai/A is in effect.
Soil application (injected into or sprayed under irrigation water) At the beginning of the growing season or at transplanting Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	1 oz/1,000 gal of irrigation water	3	90	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Avocado (continued)</b>						
Sleeve drench application At transplanting	50% WP [100-735]  2 lb/gal EC [100-607]	0.25 oz/18 gal [1 qt. solution per tree]	NS	NS	12	A 28-day PHI has been established. A maximum yearly application rate of 12 lb ai/A is in effect.
<b>Bean, succulent or dried (see also "Legume vegetables.")</b>						
Soil application At-planting Ground equipment	1% G [100-664 and 100-713]	0.12 oz/1,000 linear ft. of row	NS	NS	12	Feeding of vines or grazing of foliage by livestock is prohibited. A 12-month plant-back interval has been established for Brussels sprouts, corn, garlic, root crops, wheat, and other annual food/feed crops not registered for metalaxyl use.
<b>Blueberry</b>						
Soil band application Spring (prior to growth of established plantings) Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	3.6 lb/A	2	NS	12	
Soil band or broadcast application At-planting or postplant (new plantings) Ground equipment		4 lb/A	3	60	12	A maximum seasonal application rate of 7.2 lb ai/A is in effect for new plantings before bearing harvestable fruit.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Broccoli</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	2 lb/A	(1)	NA	12	Application may be made alone or as a tank mix with other fungicides. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
Soil broadcast application Preplant or at planting Ground equipment	2% G [100-676]  5% G [100-646]	1 lb/A	(1)	NA	NS	A 12-month plant-back interval has been established for other annual food/feed crops not registered for metalaxyl use.
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	
Broadcast foliar application Ground or aerial equipment	9% WP [100-658]	0.18 lb/A	4	14	48	A 7-day PHI has been established. A 12-month plant-back interval has been established for Brussels sprouts, celery, corn, garlic, mint, shallots, and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Broccoli (continued)</b>						
Broadcast foliar application Ground or aerial equipment	2 lb/gal EC [AZ85000800] [CA85006800]	0.25 lb/A	4	14	24 (CA) NS (AZ)	Tank mix use limited to AZ and CA. A 7-day PHI has been established for CA.
<b>Cabbage</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	2 lb/A	(1)	NA	12	See "Broccoli."
Soil broadcast application Preplant or at planting Ground equipment	2% G [100-676]  5% G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	
Broadcast application Foliar Ground or aerial equipment	9% WP [100-658]	0.18 lb/A	4	14	48	See "Broccoli."
	2 lb/gal EC [AZ85000800] [CA85006800]	0.25 lb/A	4	14	24	See "Broccoli."

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Carrot (see also "Root and tuber vegetables.")</b>						
Broadcast foliar application Ground or aerial equipment	10% WP [100-720]	0.2 lb/A	4	14	48	A 7-day PHI has been established. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
<b>Cauliflower</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	2 lb/A	(1)	NA	12	See "Broccoli."
Soil broadcast application Preplant or at planting Ground equipment	2% G [100-676]  5% G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Cauliflower (continued)</b>						
Broadcast foliar application Ground or aerial equipment	9% WP [100-658]	0.18 lb/A	4	14	48	See "Broccoli."
	2 lb/gal EC [AZ85000800] [CA85006800]	0.25 lb/A	4	14	24	See "Broccoli."
<b>Chinese broccoli (gai lan)</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]	2 lb/A	(1)	NA	12	See "Broccoli."
	2 lb/gal EC [100-607]					
Broadcast foliar application Ground or aerial equipment	9% WP [100-658]	0.18 lb/A	4	14	48	See "Broccoli."
	2 lb/gal EC [AZ85000800] [CA85006800]	0.25 lb/A	4	14	24	See "Broccoli."
<b>Chinese cabbage (tight heading varieties only)</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]	2 lb/A	(1)	NA	12	See "Broccoli."
	2 lb/gal EC [100-607]					

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Chinese cabbage (tight heading varieties only) (continued)</b>						
Broadcast foliar application Ground or aerial equipment	9% WP [100-658]	0.18 lb/A	4	14	48	See "Broccoli."
	2 lb/gal EC [AZ85000800] [CA85006800]	0.25 lb/A	4	14	24	See "Broccoli."
<b>Citrus fruit</b>						
Soil band or broadcast application At-planting or transplanting Ground equipment	5% G [100-628]	4 lb/A	3	90	127	Use limited to AZ, CA, FL, and PR. A maximum seasonal application rate of 12 lb ai/A is in effect.
	50% WP [100-735]	4 lb/A (AZ and CA)				
	2 lb/gal EC [100-607]	2 lb/A (FL and PR)				
Trunk spray application Ground equipment	50% WP [100-735]	2 lb/3 gal (AZ, CA, and TX)	3	NS	NS	Use limited to AZ, CA, FL, and TX.
	2 lb/gal EC [100-607]					
	2 lb/gal EC [FL89002300]	2 lb/15 gal (FL)				

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Citrus fruit (nonbearing)</b>						
Soil band or broadcast application At-planting Ground equipment	2% G [100-676]  5% G [100-646]  25% WP [100-717]  50% WP [100-735]  2 lb/gal EC [100-607 and 100-619]	4 lb/A	3	90	NS	
Soil drench At-planting Ground equipment	25% WP [100-717]  50% WP [100-735]  2 lb/gal EC [100-607 and 100-619]	3.75 oz/1,000 linear ft. of row	3	90	NS	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Citrus fruit (nonbearing) (continued)</b>						
Water ring drench	2 lb/gal EC [100-607]  50% WP [100-735]  5% G [100-628]	1.5 oz/100 gal [5 gal/tree] (WP and EC)  0.075 oz/tree (G)	NS	NS	12	
<b>Cotton</b>						
Soil application At-planting Ground equipment	1% G <sup>c</sup> [100-664 <sup>d</sup> and 100-713]	0.1 lb/13,000 linear ft. of row	NS	NS	12	Feeding or grazing of cotton foliage by livestock is prohibited. A 12-month plant-back interval has been established for Brussels sprouts, corn, garlic, root crops, wheat, and other annual food/feed crops not registered for metalaxyl use.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Cotton (continued)</b>						
Soil in-furrow application At-planting Ground equipment	5% G [100-628]  50% WP [100-735]  2 lb/gal EC [100-607]	0.125 lb/13,000 linear ft. of row	NS	NS	12	For the G formulation a 12-month plant-back interval has been established for barley, corn, oats, wheat, and other annual food/feed crops not registered for metalaxyl use. For the EC and WP formulations a 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use and a 14-day plant-back interval has been established for barley, oats, and wheat.
<b>Cranberry</b>						
Soil broadcast application Fall (postharvest), spring, and preharvest Ground or aerial equipment	5% G [100-628]  2 lb/gal EC [100-607]	1.75 lb/A	3	NS	12	A 45-day PHI has been established. A maximum seasonal application rate of 5.25 lb ai/A is in effect.
<b>Cucumber (see also "Cucurbit vegetables.")</b>						
Soil broadcast application Preplant or at planting Ground equipment	2% G [100-676]  5% G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Cucumber (see also "Cucurbit vegetables.") (continued)</b>						
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	
Broadcast foliar application At two-leaf stage of development Ground or aerial equipment	7% WP [100-749]	0.19 lb/A	4	7	24	A 5-day PHI has been established. Applications may be made alone or as a tank mix with other fungicides. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 40-day plant-back interval has been established for barley, oats, and wheat.
Broadcast foliar application Ground or aerial equipment	10% WP [100-629]	0.2 lb/A	4	14	24	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Cucurbit vegetables [including cucumber; gherkin; balsam pear (bitter melon); cantaloupe; casaba; Chinese waxgourd (Chinese preserving melon); citron melon; crenshaw; edible gourds; honeyballs; honeydew melon; mango melon; muskmelon; Persian melon; watermelon; pumpkin; summer squash; winter squash; and cucurbit hybrids only]</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	2 lb/A	(1)	NA	12	A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
Broadcast foliar application Ground and aerial equipment	9% WP [100-658]	0.27 lb/A	4	10	48	A 12-month plant-back interval has been established for Brussels sprouts, celery, corn, garlic, mint, shallots, and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
Broadcast foliar application Ground and aerial equipment	10% WP [100-720]	0.2 lb/A	4	14	48	A 5-day PHI has been established. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Eggplant</b>						
Soil broadcast application Preplant or at planting Ground equipment	5% G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	
Soil band application At-planting Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	2 lb/A	1	NA	12	A 7-day PHI has been established. A maximum seasonal rate of 3 lb ai/A is in effect. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
Directed spray application Postemergence Ground equipment		1 lb/A	2	30	12	
<b>Fruit Trees (nonbearing)</b>						
Soil broadcast application At-planting Ground equipment	2% G [100-676]	8 lb/A	NS	90	NS	Application to plantings that will bear harvestable fruit within 12 months of the last application is prohibited.
	5% G [100-646]					
	25% WP [100-717]	8.2 lb/A	NS	90	12	
	5% G [100-628]	4 lb/A	3	60	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Fruit Trees (nonbearing) (continued)</b>						
Soil band or broadcast application At-planting Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	4 lb/A	NS	90	12	A maximum seasonal rate of 12 lb ai/A is in effect. Application to plantings that will bear harvestable fruit within 12 months of the last application is prohibited.
Soil broadcast application Nurserystock Ground equipment	2 lb/gal EC [100-619]	4 lb/A	NS	90	12	
<b>Ginseng (see also "Root and tuber vegetables.")</b>						
Soil broadcast application Prior to spring growth Ground equipment	5% G [100-628]	0.75 lb/A	5	30	12	A 9-day PHI has been established. A maximum seasonal rate of 3 lb ai/A is in effect.
Soil drench application Prior to spring growth Ground equipment	2 lb/gal EC [100-607]	0.75 lb/A	1	NA	12	
<b>Grape</b>						
Broadcast foliar application Prebloom, early bloom, late bloom, and cluster closing Ground or aerial equipment	10% WP [100-720]	0.2 lb/A	(4)	NS	48	A 66-day PHI has been established. A total of 0.6 lb ai/A may be applied postbloom following a prebloom application.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Hops</b>						
Soil application After pruning, but before training Ground equipment	50% WP [100-735]	0.5 lb/A	1	NA	12	The feeding of hop refuse to livestock is prohibited.
Soil drench application After pruning, but before training Ground equipment	2 lb/gal EC [100-607] [ID89000500] [OR89000800] [WA89001500]	0.5 lb/A	1	NA	12	A 45-day PHI has been established. A maximum of 3 applications (one drench and two foliar) may be made per season. Foliar applications are a tank mix use with copper fungicides.
Broadcast foliar application Ground equipment			2	NS		
<b>Legume vegetables, succulent or dried [including bean: broad (fava), field, French, kidney, lima, mung, navy, pinto, runner, snap, and wax; chickpeas (garbanzo beans); lentil; lupine (sweet, white sweet, white, and grain); pea: field, garden, sugar, southern (black-eyed, catjang, cowpeas, and crowder); and edible soybeans]</b>						
Soil incorporated treatment Preplant Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	1 lb/A	(1)	NA	12	Applications to snap and dry beans may be made alone or as a tank mix. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
Soil band or broadcast application At-planting Ground or aerial equipment						
<b>Lettuce</b>						

Table A (continued).

<b>Site</b> Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
Soil broadcast application Preplant or at planting Ground equipment	2% G [100-676]  5%G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Lettuce (continued)</b>						
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	
<b>Lettuce, head</b>						
Soil band or broadcast application Preplant incorporated, at-planting, or preemergence Ground equipment	5% G [100-628]  50% WP [100-735]  2 lb/gal EC [100-607]	2 lb/A	(1)	NA	12	For the G formulation a 12-month plant-back interval has been established for barley, corn, oats, wheat, and other annual food/feed crops not registered for metalaxyl use. For the EC and WP formulations a 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
Broadcast foliar application Ground or aerial equipment	2 lb/gal EC [AZ86001500] [CA86001800] [CO88000100] [FL86000200] [TX91001100] [WA87000200]	0.25 lb/A	2	NS (AZ, CO, and WA) 14 (CA, FL, and TX)	NS	Use limited to AZ, CA, CO, FL, TX, and WA. Tank mix use. Application may be made in 25-100 gal/A by ground or in 5-10 gal/A by air. A 10-day PHI has been established for CA, CO, and WA and a 7-day PHI has been established for TX.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Melon (including cantaloupe, casaba, crenshaw, honeydew, and watermelon) (see also "Cucurbit vegetables.")</b>						
Soil broadcast application Preplant or at planting Ground equipment	2% G [100-676]  5% G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	
Broadcast foliar application At two-leaf stage of development Ground or aerial equipment	7% WP [100-749]	0.19 lb/A	4	7	24	See "Cucumber."
Broadcast application Foliar Ground or aerial equipment	10% WP [100-629]	0.2 lb/A	4	14	24	See "Cucumber."
<b>Nut Trees (nonbearing)</b>						
Soil broadcast application At-planting Ground equipment	2% G [100-676]	8 lb/A	NS	90	NS	See "Fruit Trees (nonbearing)."
	5% G [100-646]					
	25% WP [100-717]	8.2 lb/A	NS	90	12	
	5% G [100-628]	4 lb/A	3	60	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Nut Trees (nonbearing) (continued)</b>						
Soil band or broadcast application At-planting Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	4 lb/A	NS	90	12	See "Fruit Trees (nonbearing)."
Soil broadcast application Nurserystock Ground equipment	2 lb/gal EC [100-619]	4 lb/A	NS	90	12	See "Fruit Trees (nonbearing)."
<b>Onion, bulb (including onions grown for seed)</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	1 lb/A	(1)	NA	12	A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
Broadcast foliar application Ground or aerial equipment	10% WP [100-629]	0.2 lb/A	4	14	24	A 7-day PHI has been established. Applications may be made alone or as a tank mix with other fungicides. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 40-day plant-back interval has been established for barley, oats, and wheat.
	7% WP [100-749]	0.19 lb/A	4	7	24	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Onion, bulb (including onions grown for seed) (continued)</b>						
Broadcast foliar application Ground or aerial equipment	9% WP [100-658]	0.18 lb/A	4	7	48	A 7-day PHI has been established. A 12-month plant-back interval has been established for Brussels sprouts, celery, corn, garlic, mint, shallots, and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
<b>Onion, green (including green onions, leeks, spring onions, scallions, Japanese bunching onions, green shallots, and green eschalots)</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	1 lb/A	(1)	NA	12	See "Onions, bulb (including onions grown for seed)."
Broadcast foliar application Ground or aerial equipment	9% WP [100-658]	0.18 lb/A	3	7	48	A 21-day PHI has been established. A 12-month plant-back interval has been established for Brussels sprouts, celery, corn, garlic, mint, shallots, and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Papaya</b>						
Soil drench application Seedling (nurserystock) Ground equipment	50% WP [100-735]	3.4 lb/A	2	30	12	Use limited to HI. A 90-day PHI has been established.
Soil broadcast application At transplanting Ground or aerial equipment	2 lb/gal EC [100-607]	3.5 lb/A	2	14	12	
<b>Peanut</b>						
Soil band application At-planting Ground equipment	1% G [100-664 and 100-713]	0.25 lb/14,520 linear ft. of row	NS	NS	12	The feeding or grazing of peanut foliage or harvest residues by livestock is prohibited. A maximum seasonal rate of 1 lb ai/14,520 linear ft. of row is in effect. A 12-month plant-back interval has been established for Brussels sprouts, corn, garlic, root crops, wheat, and other annual food/feed crops not registered for metalaxyl use.
Soil band application At early pegging Ground equipment		1 lb/14,520 linear ft. of row	NS	NS	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Peanut (continued)</b>						
Soil band or in-furrow application At-planting Ground equipment	. 5% G [100-628]	0.25 lb/A (based on 42-inch rows)	NS	NS	12	A 12-month plant-back interval has been established for barley, corn, oats, wheat, and other annual food/feed crops not registered for metalaxyl use.
Soil band application At early pegging Ground equipment		0.5 lb/12,400 linear ft. of row	NS	NS	12	
Soil band or in-furrow application At-planting Ground equipment	50% WP [100-735]	0.25 lb/A	NS	NS	12	A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14- day plant-back interval has been established for barley, oats, and wheat.
Soil application At early pod set or pegging Irrigation equipment	2 lb/gal EC [100-607]	1 lb/A	NS	NS	12	
<b>Pepper</b>						
Soil broadcast application Preplant or at planting Ground equipment	5% G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Pepper (continued)</b>						
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	
Soil band application At-planting Ground equipment	50% WP [100-735]	2 lb/A	1	NA	12	See "Eggplant."
Directed spray application Postemergence Ground equipment	2 lb/gal EC [100-607]	1 lb/A	2	30	12	
Broadcast foliar application Ground or aerial equipment	10% WP [100-720]	0.25 lb/A	4	10	48	A 7-day PHI has been established. A maximum seasonal rate of 3 lb ai/A is in effect. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
<b>Pineapple</b>						
Seedpiece dip (crown dip) treatment Preplant	50% WP [100-735]  2 lb/gal EC [100-607]	1 lb/100 gal [100 gal of dip solution per planted acre]	NA	NA	NA	If crop failure occurs within one year of planting treated crowns, the use of crop or crop residue for animal feed is prohibited.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Potato</b>						
Broadcast foliar application Ground or aerial equipment	10% WP [100-720]	0.25 lb/A	4	14	48	A 7-day PHI has been established. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
	7% WP [100-749]	0.16 lb/A	4	14	24	A 3-day PHI has been established for CT, DE, FL, ME, MA, MI, NH, NY, OH, PA, RI, VT, and WI. A 14-day PHI has been established for all other states. Applications may be made alone or as a tank mix with other fungicides. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 40-day plant-back interval has been established for barley, oats, and wheat.
	10% WP [100-629]	0.2 lb/A				

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Potato (continued)</b>						
Broadcast foliar application Ground or aerial equipment	10% WP [ME92000200] [ND92000200]	0.2 lb/A	4	14	NS	Use limited to ME and ND. A 7-day PHI has been established. Only sweet corn, corn grown for grain or feed where the corn and cob are ground together may be rotated the spring following the last application of metalaxyl to potatoes. Grazing or feeding forage or fodder from rotational corn to livestock is prohibited.
	9% WP [100-658]	0.18 lb/A	4	14	48	A 7-day PHI has been established. A 12-month plant-back interval has been established for Brussels sprouts, celery, corn, garlic, mint, shallots, and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
<b>Radish (see also "Root and tuber vegetables.")</b>						
Broadcast foliar application Ground or aerial equipment	10% WP [100-720]	0.2 lb/A	4	14	48	See "Carrot."

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Radish (grown for seed)</b>						
Broadcast foliar application After bolting Ground or aerial equipment	10% WP [WA92004200]	0.2 lb/A	2	14	NS	Use limited to WA. Applications may be made in a minimum of 30 gal/A using ground equipment and 10 gal/A using aerial equipment. The feeding or grazing of radish forage or fodder and the cutting of radish tops for hay or forage are prohibited. The use of any portion of the treated field, including seed, seed screenings, hay, forage, or stubble for human or animal consumption is prohibited.
<b>Raspberry</b>						
Soil band application Spring and fall (postharvest) Ground equipment	5% G [100-628]  50% WP [100-735]  2 lb/gal EC [100-607]	0.25 lb/1,000 linear ft. of row [3.6 lb/A on broadcast basis of G formulation]	2	NS	12	A 45-day PHI has been established.
<b>Rice</b>						
Soil broadcast application Preplant/preflooding Ground or aerial equipment	2 lb/gal EC [LA92000400] <sup>c</sup>	0.5 lb/A	(1)	NA	NS	Use limited to LA. Applications may be made in a minimum of 20 gal/A using ground equipment and 5-10 gal/A using aerial equipment.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Root and tuber vegetables [including artichoke (Jerusalem); beet (sugar and table); carrot; cassava; chicory; ginger; ginseng; horseradish; parsnip; radish; rutabaga; salsify; sweet potato; turnip; and yam]</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	2 lb/A	(1)	NA	12	Application to potato is prohibited.
<b>Soybean</b>						
Soil band or in-furrow application At-planting Ground equipment	5% G [100-628]  50% WP [100-735]  2 lb/gal EC [100-607]	0.3 oz/1,000 linear ft. of row	(1)	NA	12	See "Cotton."
Soil band or broadcast application At-planting Ground or aerial equipment	50% WP [100-735]  2 lb/gal EC [100-607]	1.25 lb/A	(1)	NA	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Spinach</b>						
Soil broadcast application Preplant or at planting Ground equipment	2% G [100-676]  5% G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."
Soil incorporated treatment Preplant Ground equipment	5% G [100-628]  50% WP [100-735]  2 lb/gal EC [100-607]	2 lb/A	(1)	NA	12	A 21-day PHI has been established. Two additional sidedress applications may be made at 0.25 lb ai/A/application. For the G formulation a 12-month plant-back interval has been established for barley, corn, oats, wheat, and other annual food/feed crops not registered for metalaxyl use. For the EC and WP formulations a 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use and a 14-day plant-back interval has been established for barley, oats, and wheat.
Soil band or broadcast application At-planting/preemergence Ground or aerial equipment						
Broadcast application Preplant or at planting Ground or aerial equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Spinach (continued)</b>						
Soil band or broadcast application At-planting (fall) and breaking dormancy (late winter) Ground equipment	5% G [AR90000502] [OK90000500]  2 lb/gal EC [AR90000501] [OK90000500]	0.5 lb/A	2	NS	NS	Use limited to AR and OK. A maximum seasonal rate of 1 lb ai/A is in effect. A 21-day PHI has been established.
Broadcast foliar application Ground or aerial equipment	10% WP [100-720]	0.25 lb/A	2	14	48	A 21-day PHI has been established. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
<b>Squash (see also "Cucurbit vegetables.")</b>						
Soil broadcast application Preplant or at planting Ground equipment	2% G [100-676]  5% G [100-646]	1 lb/A	(1)	NA	NS	See "Broccoli."
Soil broadcast application Preplant or at planting Ground equipment	25% WP [100-717]	0.68 lb/A	NS	NS	12	
Broadcast foliar application At two-leaf stage of development Ground or aerial equipment	7% WP [100-749]	0.19 lb/A	4	7	24	See "Cucumber."

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Squash (see also "Cucurbit vegetables.") (continued)</b>						
Broadcast foliar application Ground or aerial equipment	10% WP [100-629]	0.2 lb/A	4	14	24	See "Cucumber."
<b>Stone fruits [including apricots; cherries (sweet and sour); nectarines; peaches; plums (Chickasaw, Damson, Japanese); and prunes]</b>						
Soil band or broadcast application After planting (new plantings) or prior to spring plant growth (established plantings) Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	4 lb/A	3	60	12  7	See "Almond."
<b>Strawberry</b>						
Soil band or broadcast application Post-transplant, prebloom, preharvest, or postharvest Ground or irrigation equipment	50% WP [100-735]  2 lb/gal EC [100-607] [CA90004700]	1 lb/A	3	NS	12	A maximum seasonal rate of 3 lb ai/A is in effect.
Soil broadcast application Postplant (nurserystock)	2 lb/gal EC [CA82002400]	1 lb/A	2	30	NS	A maximum seasonal rate of 2 lb ai/A is in effect.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Sugar beet (see also "Root and tuber vegetables.")</b>						
Soil band or broadcast application Preplant incorporated or at-planting Ground equipment	5% G [100-628]	2 lb/A	(1)	NA	12	
Broadcast foliar application Ground or aerial equipment	10% WP [100-629]	0.2 lb/A	4	14	24	A 14-day PHI has been established. The feeding of treated tops to livestock is prohibited.
<b>Tobacco</b>						
Soil broadcast application Preplant or at-planting (plant bed) Ground equipment	50% WP [100-735]	0.5 lb/A	(1)	NA	12	Application may be made in 50 gal/A. In PA, 1 lb ai/A may be used.
Soil incorporated treatment Pretransplant Ground equipment	2 lb/gal EC [100-607]	3 lb/A	(1)	NA	12	In PA, the application rate of 3 lb ai/A is prohibited.
<b>Tomato</b>						
Soil broadcast application Preplant or at-planting Ground equipment	2% G [100-676]	1 lb/A	(1)	NA	NS	See "Broccoli."
	5% G [100-646]					
	25% WP [100-717]	0.68 lb/A	NS	NS	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Tomato (continued)</b>						
Soil band application At-planting Ground equipment	5% G [100-628]	1 lb/A	NS	NS	12	A 7-day PHI has been established. Two additional applications may be made at 0.5 lb ai/A. A maximum seasonal rate of 2 lb ai/A is in effect. A 12-month plant-back interval has been established for barley, corn, oats, wheat, and other annual food/feed crops not registered for metalaxyl use.
Soil band application At-planting Ground equipment	50% WP [100-735] 2 lb/gal EC [100-607]	2 lb/A	NS	NS	12	A 7-day PHI has been established for the WP formulation. A maximum seasonal rate of 3 lb ai/A is in effect. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.
Directed soil or soil injection application Postplant Ground equipment		1 lb/A	2	NS	12	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Tomato (continued)</b>						
Broadcast foliar application Ground or aerial equipment	7% WP [100-749]	0.19 lb/A (east of the Mississippi)  0.16 lb/A (west of the Mississippi)	4	7	24	A 5-day PHI has been established. Applications may be made alone or as a tank mix with other fungicides. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 40- day plant-back interval has been established for barley, oats, and wheat.
	10% WP [100-629]	0.2 lb/A	4	14	24	
Broadcast foliar application Ground or aerial equipment	9% WP [100-658]	0.27 lb/A	4	10	48	A 7-day PHI has been established. A 12-month plant-back interval has been established for Brussels sprouts, celery, corn, garlic, mint, shallots, and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
Broadcast foliar application Ground or aerial equipment	10% WP [100-720]	0.25 lb/A	NS	5	48	A 7-day PHI has been established. A maximum seasonal rate of 3 lb ai/A is in effect. A 12-month plant-back interval has been established for corn and other annual food/feed crops not registered for metalaxyl use. A 14-day plant-back interval has been established for barley, oats, and wheat.

Table A (continued).

<b>Site</b> Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Walnut</b>						
Soil band or broadcast application After planting (new plantings) or prior to spring plant growth (established plantings) Ground equipment	50% WP [100-735]  2 lb/gal EC [100-607]	4 lb/A	3	60	12	See "Almond."

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Seed Treatments</b>						
<b>Beet (table/garden)</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	
<b>Cotton (acid delinted and machine delinted)</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	
<b>Corn (sweet)</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	1 oz/cwt	NA	NA	None	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Corn [field and pop (including forage and fodder)]</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]	0.5 oz/cwt	NA	NA	None	
	50% WP [100-738]					
	3 lb/gal FIC [100-684]					
<b>Corn</b>						
Seed treatment (export use) Slurry	25% WP [100-639]	3.5 oz/cwt (WP) 3.75 oz/cwt (FIC)	NA	NA	None	Use of seed for feed, food, or oil purposes is prohibited.
	50% WP [100-738]					
	3 lb/gal FIC [100-684]					
<b>Dill</b>						
Seed treatment (domestic use) Slurry	3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Forage grasses (hay and silage)</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	
<b>Forage legumes [alfalfa, bean forage, clover, cowpea forage and hay, lespedeza, pea forage, pea vine hay, soybean forage and hay, trefoil, velvet bean forage, and vetch]</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	
<b>Grain crops [barley, buckwheat, corn: field, sweet, and pop, millet, oats, rice, rye, triticale, and wheat (including forage and fodder)]</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Legume vegetables [beans (field, green, kidney, lima, navy, pole, snap, string, and wax); chick peas (garbanzo beans); cowpeas; lentils; lupines; peas (black-eyed, field, and garden); and edible soybeans]</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	
<b>Okra</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	
<b>Millet</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. *	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Millet (continued)</b>						
Seed treatment (export use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	3.5 oz/cwt (WP) 3.75 oz/cwt (FIC)	NA	NA	None	Use of seed for feed, food, or oil purposes is prohibited.
<b>Pea</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	1.25 oz/cwt (WP) 1.4 oz/cwt (FIC)	NA	NA	None	
Seed treatment (export use) Slurry	35% WP [CA85007300] [ID83003300] [WA83003500]	1.12 oz/cwt	NA	NA	None	Use limited to CA, ID, and WA for export only. Use of seed for feed, food, or oil purposes is prohibited.
	35% WP [WA90003300]	1.1375 oz/cwt				
	45% WP [CA87000700] [ID86001900] [WA86002800]	1.125 oz/cwt	NA	NA	None	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Pea (continued)</b>						
Seed treatment (export use) Slurry	25% WP [100-639]	0.5 oz/cwt	NA	NA	None	Use of seed for feed, food, or oil purposes is prohibited.
	50% WP [100-738]					
<b>Sorghum</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]	1 oz/cwt	NA	NA	None	
	50% WP [100-738]					
	3 lb/gal FIC [100-684]					
Seed treatment (domestic use) Slurry	2 lb/gal EC [100-626]	0.5 oz/cwt	NA	NA	None	Use of seed for feed, food, or oil purposes is prohibited.
Seed treatment (export use) Slurry	25% WP [100-639]	3.5 oz/cwt (WP) 3.75 oz/cwt (FIC)	NA	NA	None	Use of seed for feed, food, or oil purposes is prohibited.
	50% WP [100-738]					
	3 lb/gal FIC [100-684]					

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Soybean</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  2 lb/gal EC [100-626]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	EC formulation: Use of seed for feed, food, or oil purposes is prohibited.
<b>Sugar beet</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	0.5 oz/cwt	NA	NA	None	

Table A (continued).

Site Application Type Application Timing Application Equipment	Formulation [EPA Reg. No.]	Max. Single Application Rate (ai)	Max. # Apps. <sup>a</sup>	Min. Retreatment Interval (Days)	Restricted Entry Interval (Hours)	Use Limitations <sup>b</sup>
<b>Sunflower</b>						
Seed treatment (domestic use) Slurry	25% WP [100-639]  50% WP [100-738]  2 lb/gal EC [100-626]  3 lb/gal FIC [100-684]	1 oz/cwt	NA	NA	None	EC formulation: Use of seed for feed, food, or oil purposes is prohibited.
Seed treatment (export use) Slurry	25% WP [100-639]  50% WP [100-738]  3 lb/gal FIC [100-684]	3.5 oz/cwt (WP) 3.75 oz/cwt (FIC)	NA	NA	None	Use of seed for feed, food, or oil purposes is prohibited.

- <sup>a</sup> Label instructions imply a maximum number of applications but it is not strictly stated; the implied number of applications is indicated in parentheses.
- <sup>b</sup> PHI = preharvest interval and PFI = prefeeding interval.
- <sup>c</sup> CBRS considers the feeding restrictions for cotton forage to livestock impractical. This feeding restriction must be removed from the use limitations for cotton on the EPA Reg. Nos. 100-664 and 100-713 labels. The registrant must amend the labels for the 1% G formulations (EPA Reg. Nos. 100-664 and 100-713).
- <sup>d</sup> The 1% G formulation [EPA Reg. No. 100-664] product label should be amended to list lb of product/13,000 linear ft. of row instead of lb of product/linear ft. of row.
- <sup>e</sup> Revised label for SLN No. LA92000400 must be submitted with the following restrictions: When using this product as directed in making a direct soil application at-planting to water seeded rice, do not use Apron<sup>®</sup>-treated rice seed or any other rice seed that has been treated with metalaxyl (CBTS Nos. 11708 and 11709, DP Barcodes D189772 and D189764, 5/5/93, F. Griffith).

Table B. Residue Chemistry Science Assessments for Reregistration of Metalaxyl.

GLN: Data Requirements	Current Tolerances, ppm [40 CFR]	Must Additional Data Be Submitted?	References <sup>1</sup>
171-3: Directions for Use	N/A	No <sup>2</sup>	
171-4 (a): Plant Metabolism	N/A	No	00071601 00071602 00071603 00071604 00071605 00071606 00071607 00071608 00071609 00071610 00114379 00128102
171-4 (b): Animal Metabolism	N/A	Yes <sup>3</sup>	00071611 00071612 41664503 41664504 41664505 41664506 42115801 <sup>4</sup> 42115802 <sup>4</sup> 42115803 <sup>4</sup> 42115804 <sup>4</sup> 42115805 <sup>4</sup> 42115806 <sup>4</sup>
171-4 (c/d): Residue Analytical Methods	N/A	No	00071622 00071623 00071676 00104378 00104656 00148440 00157480 40503101 41055203 41689701 41689702 42115809 <sup>4</sup> 42115810 <sup>4</sup> 42115807 <sup>4</sup> 42115808 <sup>4</sup>
171-4 (e): Storage Stability	N/A	Yes <sup>5</sup>	00148440 40106601 <sup>6</sup> 41449001 <sup>7</sup> 42021101 <sup>8</sup> 42115809 <sup>4</sup> 42350101 <sup>9</sup> 42919401 <sup>5</sup> 41912902 <sup>10</sup>
171-4 (k): Magnitude of the Residue in Plants			
<u>Root and Tuber Vegetables Group</u>	0.5 [§180.408(a)]	No <sup>11</sup>	40838301 <sup>12</sup> 40838302 <sup>12</sup> 40838303 <sup>12</sup>
- Beets	0.1 [§180.408(a)]	No <sup>13</sup>	00128102
- Ginseng	3.0 [§180.408(a)]	No	41688900 <sup>14</sup> 41688901 <sup>14</sup>
- Potatoes	0.5 [§180.408(a)]	No <sup>11</sup>	00104654 00071616 00098428 <sup>15</sup>
- Sugar beets	0.1 [§180.408(a)] 0.5 (roots) [§180.408(a)] <sup>11</sup>	No <sup>13</sup> No	00128102 40569301 <sup>16</sup>
<u>Leaves of Root and Tuber Vegetables Group</u>	15.0 [§180.408(a)]	No	40838301 <sup>12</sup> 40838302 <sup>12</sup> 40838303 <sup>12</sup>
- Beet, tops	0.1 [§180.408(a)]	No <sup>13</sup>	00128102
- Sugar beet, tops	10.0 [§180.408(a)]	No <sup>13</sup>	40569301 <sup>16</sup>

Table B (continued).

GLN: Data Requirements	Current Tolerances, ppm [40 CFR]	Must Additional Data Be Submitted?	References <sup>1</sup>
<b><u>Bulb Vegetables Group</u></b>			
- Onions, dry bulb	3.0 [§180.408(a)]	No	00071615 00098428 00130694 00148103
- Onions, green	10.0 [§180.408(a)]	No	00071615 00098428 00130694
<b><u>Leafy Vegetables (Except Brassica Vegetables) Group</u></b>			
	0.1 [§180.408(a)]	No <sup>17</sup>	41587801 <sup>18</sup> 42021101 <sup>17</sup>
- Lettuce, head	5.0 [§180.408(a)]	No <sup>17</sup>	00071615 00097511 00114377 00103695 40775801 <sup>19</sup>
- Spinach	10.0 [§180.408(a)]	No	00071672 00114378 00130695 40790201 <sup>20</sup> 41636201 <sup>21</sup>
<b><u>Brassica (Cole) Leafy Vegetables Group</u></b>			
- Brassica (cole) leafy vegetable group [except broccoli, cabbage, and cauliflower]	0.1 [§180.408(a)]	No <sup>22</sup>	
- Broccoli	2.0 [§180.408(a)]	No	00071615 00130773
- Cabbage	2.0 [§180.408(a)]	No <sup>22</sup>	00071615 00130773
- Cauliflower	2.0 [§180.408(a)]	No <sup>22</sup>	00071615 00130773
<b><u>Legume Vegetables Group</u></b>			
- Legume vegetables group	0.2 [§180.408(a)]	No <sup>23</sup>	00129003 <b>40569303</b>
- Soybean, grain	1.0 [§180.408(a)]	No	00071672 00104390 00148440
<b><u>Foliage of Legume Vegetables Group</u></b>			
	8.0 [§180.408(a)]	No	00129003 00071672 00104390 00148440 <b>40569303</b>
<b><u>Fruiting Vegetables (Except Cucurbits) Group</u></b>			
	1.0 [§180.408(a)]	Yes <sup>24</sup>	00148103 00148440 00157480
<b><u>Cucurbit Vegetables Group</u></b>			
	1.0 [§180.408(a)]	No	00071615 00098428 00130693 00148103
<b><u>Citrus Fruits Group</u></b>			
	1.0 [§180.408(a)]	No	00117969 00133020 00148440
<b><u>Pome Fruits Group</u></b>			
- Apples	0.2 [§180.408(a)]	No	00126315 00141519
<b><u>Stone Fruits Group</u></b>			
	1.0 [§180.408(a)]	No	00164650 <sup>25</sup>
<b><u>Small Fruits and Berries Group</u></b>			
- Blueberries	2.0 [§180.408(a)]	No	00164649 <sup>25</sup>

Table B (continued).

GLN: Data Requirements	Current Tolerances, ppm [40 CFR]	Must Additional Data Be Submitted?	References <sup>1</sup>
- Cranberries	4.0 [§180.408(a)]	No	41996401 <sup>26</sup>
- Grapes	2.0 [§180.408(a)]	No	00138818 <sup>27</sup> 00155845 <sup>27</sup> 41150101 <sup>28</sup>
- Raspberries	0.5 [§180.408(a)]	No	00127769
- Strawberries	10.0 [§180.408(a)]	Yes <sup>29</sup>	00155237 <sup>30</sup> 40880401 <sup>31</sup>
<b><u>Tree Nuts Group</u></b>			
- Almonds	0.5 (nutmeats) 10.0 (hulls) [§180.408(a)]	No	00164651 <sup>25</sup>
- Walnuts	0.5 [§180.408(a)]	No	00164651 <sup>25</sup>
<b><u>Cereal Grains Group</u></b>			
- Grain, crops	0.1 [§180.408(a)]	No <sup>32</sup>	00071672 00104387 00128102 <b>41689701</b> <b>41689702</b> 41870301 <sup>33</sup>
<b><u>Forage, Fodder and Straw of Cereal Grains Group</u></b> <b><u>(except wheat, barley, and oats)</u></b>	None	No <sup>32</sup>	00128102 <b>41689701</b> <b>41689702</b> 41870301 <sup>33</sup> 41870305 <sup>33</sup> 41912903 <sup>34</sup>
<b><u>Grass Forage, Fodder, and Hay Group</u></b>			
- Grasses, forage	0.1 [§180.408(a)]	No <sup>35</sup>	
<b><u>Non-grass Animal Feeds Group</u></b>			
- Alfalfa	6.0 (forage) 20.0 (hay) [§180.408(a)]	No No	40832901 <sup>37</sup>
<b><u>Miscellaneous Commodities</u></b>			
- Asparagus	7.0 [§180.408(a)]	Yes <sup>38</sup>	00154446 <sup>38</sup>
- Avocados	4.0 [§180.408(a)]	No	00074488
- Cottonseed	0.1 [§180.408(a)]	Yes <sup>39</sup>	00109402 41870305 <sup>33</sup>
- Hops, green	2.0 [§180.408(a)]	No <sup>40</sup>	00079433 40746901 <sup>41</sup> 40909401 <sup>42</sup>
- Papaya	0.1 [§180.408(c)]	No	40490501 <sup>43</sup>
- Peanut	0.2 (nutmeats) 20.0 (vines) 20.0 (hay) 2.0 (hulls) [§180.408(a)]	No No <sup>44</sup> No No	00128738 41870306 <sup>45</sup> 42498701 <sup>46</sup>

Table B (continued).

GLN: Data Requirements	Current Tolerances, ppm [40 CFR]	Must Additional Data Be Submitted?	References <sup>1</sup>
- Pineapples	0.1 (fruit) 0.1 (fodder) 0.1 (forage) [§180.408(a)]	No No <sup>47</sup> No <sup>47</sup>	00109472
- Sunflowers	0.1 (seeds) 0.1 (forage) [§180.408(a)]	No No	00128102 4191290210
- Tobacco	N/A	No	00100467 00100476 00100477 00100478 00104485 00140371 00148440 42196503
171-4(l): Magnitude of the Residue in Processed Food/Feed			
- Apples	2.0 (pomace, dry) 0.4 (pomace, wet) [§186.4000 (a)]	No <sup>48</sup> No	00126315
- Citrus	7.0 (oil) [§185.4000 (a)] 7.0 (molasses) 7.0 (pulp) [§186.4000 (a)]	No No No	00117969
- Cereal grains group (except wheat, barley, and oats)	None	No	41870301 <sup>33</sup> 41870302 <sup>33</sup> 41870303 <sup>33</sup> 42498701 <sup>49</sup> 42259805 <sup>50</sup>
- Cottonseed	None (cotton gin bypds)	Yes <sup>39</sup>	
- Grapes	6.0 (raisins) [§185.4000 (a)] 10.0 (raisin waste) 10.0 (pomace, dry) 10.0 (pomace, wet) [§186.4000 (a)]	No	00138818 <sup>27</sup> 00155845 <sup>27</sup> 41150101 <sup>28</sup>
- Hops	20.0 (dried) [§185.4000 (d)] 2.0 (dry) 20.0 (spent) [§186.4000 (d)]	No <sup>40</sup> No <sup>40</sup> No <sup>40</sup>	00079433 40746901 40909401
- Peanut	1.0 (meal) 2.0 (soapstock) [§186.4000 (a)]	No No <sup>51</sup>	00128738
- Pineapples	None	No <sup>52</sup>	42233501 <sup>52</sup>

Table B (continued).

GLN: Data Requirements	Current Tolerances, ppm [40 CFR]	Must Additional Data Be Submitted?	References <sup>1</sup>
- Potatoes	4.0 (processed, including potato chips) [§185.4000 (a)] 4.0 (waste, dried, processed) [§186.4000 (a)]	No <sup>53</sup>	41870307 <sup>54</sup>
- Prunes (dried)	4.0 [§185.4000 (a)]	No	00164650 <sup>25</sup>
- Soybean	2.0 (hulls) 2.0 (meal) 2.0 (soapstock) [§186.4000 (a)]	No No No <sup>55</sup>	00071672
- Sugar beet	1.0 (molasses) [§186.4000 (a)]	No <sup>56</sup>	401066016 40569301 <sup>57</sup>
- Sunflower	None	No <sup>58</sup>	41870304 <sup>59</sup> 42498701 <sup>60</sup>
- Tomatoes	20.0 (pomace, dry and wet) [§186.4000 (a)] 3.0 (processed) [§185.4000 (a)]	Yes <sup>61</sup>	00148440 00157480
171-4 (j): Magnitude of the Residue in Meat, Milk, Poultry, and Eggs			
- Fat, kidney, and liver of cattle, goats, hogs, horses, poultry, and sheep	0.4 [§180.408 (a)]	Reserved	00100753 42115809 <sup>4</sup> 00071673 00071674
- Meat and meat byproducts (except kidney and liver) of cattle, goats, hogs, horses, poultry, and sheep	0.05 [§180.408 (a)]	Reserved	00100753 42115809 <sup>4</sup> 00071673 00071674
- Eggs	0.05 [§180.408 (a)]	Reserved	00071673 42115809 <sup>4</sup>
- Milk	0.02 [§180.408 (a)]	Reserved	00071674
165-1: Rotational Crops (Confined)		Reserved	42196501 <sup>62</sup> 42196502 <sup>62</sup> 42196503 <sup>62</sup>
165-2: Rotational Crops (Field)		Reserved	41870308 <sup>63</sup>
- Barley, oat, and wheat	2.0 (forage, fodder, and straw) [§180.408 (b)] 0.2 (grain) [§180.408 (b)] 1.0 (milling fractions) [§185.4000 (b) and §186.4000 (b)]		00114376 00071672 00104387 00114376 00071672 00104387 00114376

Table B (continued).

1. **Bolded** references were reviewed in the Update of 3/13/91. Unbolded references were reviewed in the Residue Chemistry Science Chapter (6/22/87) of the Metalaxyl Final Reregistration Standard and Tolerance Reassessment (FRSTR) (Guidance Document date 9/88). *Italicized* references were reviewed in the 12/81 Registration Standard. Otherwise, references were reviewed as noted.
2. A revised label for SLN No. LA920004 must be submitted with the following restriction: When using this product as directed in making a direct soil application at planting to water seeded rice, do not use Apron®-treated rice seed or any other rice seed that has been treated with metalaxyl (CBTS Nos. 11708 and 11709, DP Barcodes D189772 and D189764, 5/5/93, F. Griffith).  
  
In addition, The registrant must amend the labels for the 1% G formulations (EPA Reg. Nos. 100-664 and 100-713) to impose a 75-day PHI for peanuts and remove the feeding restriction for cotton forage, which CBRS considers to be impractical.
3. CB No. 9102, D172350, S. Hummel, 9/16/93. Additional method validation data are needed, including recoveries of representative metalaxyl metabolites including at least one containing the 2,6-dimethyl aniline moiety, CGA-94689, P1, and P2.
4. CB No. 9102, DP Barcode D172350, 9/15/93, S. Hummel.
5. A storage stability data requirement is outstanding for a representative oilseed (e.g., soybeans or a nut) and a grain crop (e.g., wheat). Storage stability data requirements are also outstanding for all processed plant commodities from an oilseed, a grain, and a fruit or fruiting vegetable. In addition, storage stability data requirements are outstanding for livestock commodities (CBRS No. 12888, DP Barcode D197066, 2/17/94, S. Hummel). Storage stability data are needed for metalaxyl and representative metabolites including CGA-94689.
6. CB No. 1996 and 2013, 3/13/87, F. Griffith.
7. CBRS No. 7193, 1/15/91, R. Perfetti.
8. CBTS No. 8898, DP Barcode D171050, 12/18/91, R. Lascola.
9. CBTS No. 10062, DP Barcode No. D179628, 7/3/92, M. Peters.
10. CBRS No. 8166, DP Barcode D165608, 4/16/92, J. Abbotts.
11. The group tolerance for root and tuber vegetables should be renamed "root and tuber vegetables group (except ginseng)," concomitant with the revocation of the established tolerances for beets, potatoes, and sugar beets.
12. PP#9F3698, CB Nos. 4788 to 4791, 8/3/89, J. Garbus; CB No. 5836, 10/23/89, J. Garbus; CB No. 6176, 2/14/90, J. Garbus; CB No. 8625, 12/12/91, K. Dockter.
13. The 0.1 ppm tolerances on beets, beet tops, and sugar beets are based on residue data generated from field trials on treated seeds. As group tolerances for root and tuber vegetables (except ginseng) and for leaves of root and tuber vegetables will cover these commodities, and as the preplant and foliar uses have superseded the seed treatment uses, these tolerances should be revoked.
14. PP#1E3926, CB No. 7340, 2/22/91, G. Otakie; CB No. 7787, 4/8/91, G. Otakie; CBTS Nos. 9054 and 9058, 3/6/92, W. Chin.

Table B (continued).

15. CB No. 60, 2/14/86, W. Anthony.
16. PP#3F3617/FAP#8D5554, CB Nos. 3908 through 3912, 11/28/88, F. Griffith; and CB Nos. 5934, 5935, and 6545, 4/30/90, F. Griffith.
17. CBTS recommends in favor of establishing a crop group tolerance of 5 ppm for leafy vegetables, (except Brassica), excluding spinach. Once the 5 ppm group tolerance has been established, the 5 ppm tolerance for lettuce should be revoked (PP#0F3893; CBTS No. 8898, DP Barcode 171050, 12/18/91, R. Lascola; and CBTS No. 9710, DP Barcode 176810, 5/12/92, R. Lascola).
18. CBTS No. 7171, DP Barcode D156811, 6/21/91, S. Bacchus.
19. CB No. 4808, 2/9/89, L. Propst.
20. CB No. 4809, 2/14/89, L. Propst; and CB No. 7418, 1/24/91, M. Metzger.
21. CBRS No. 7165, DP Barcode D156892, 2/19/91, K. Dockter.
22. The established 0.1 ppm crop group tolerance for Brassica (cole) leafy vegetables group (except broccoli, cabbage, and cauliflower) was established to cover seed treatments. However, as there is no registered seed treatment on any member of this group, this tolerance should be revoked. A petition proposing a crop group tolerance of 5 ppm on brassica leafy vegetables to cover soil and foliar uses is in reject status (PP#2F04072, DP Barcode D174355, 1/15/93, M. Peters). Data submitted with this petition (MRID 42159801) indicate that the established 2 ppm tolerances for cabbage and cauliflower can be reduced to 1 ppm. Data submitted with the petition indicate that a crop group tolerance is inappropriate because residue levels differ by more than a factor of 5.
23. The group tolerance for the legume vegetable group should be changed to "legume vegetables group excluding soybeans."
24. Post emergence use of the 50% WP formulation on fruiting vegetables is not supported. Supporting data must be provided or the use removed from the 50% WP label.
25. PP#7F3470/FAP#7H5520, CB Nos. 1676, 1677, and 1678, 3/6/87, M. Nelson; CB Nos. 2604, 2699, and 2700, 8/26/87, M. Nelson; and CB Nos. 3788 and 3789, 5/13/88, M. Nelson.
26. PP#1E4024, CB No. 8582, DP Barcode D168812, 10/31/91, J. Morales; and CBTS No. 9537, DP Barcode D175506, 7/17/92, J. Morales.
27. PP#4G3031/FAP#4H5424, 5/30/84, L. Cheng; and CB No. 960, 6/3/86, L. Cheng.
28. PP#6F3362/FAP#6H5493, CB Nos. 5557 and 5558, 12/11/89, G. Otakie; and CB Nos. 7646 and 7647, DP Barcodes D160990 and D161029, 9/9/91, G. Otakie.
29. Post emergence use of the 50% WP formulation on strawberries is not supported. Supporting data must be provided or the use removed from the 50% WP label.
30. PP#6F3337, CB No. 360, 2/21/86, M. Firestone; CB No. 988, 9/8/86, M. Firestone; and CB No. 2301, 9/3/87, M. Kovacs.
31. CB Nos. 4814, 4815, and 4816, 5/10/89, M. Kovacs; and CB No. 5895, 11/2/89, D. Edwards.

Table B (continued).

32. CBRS recommends that the proposed tolerances of 0.1 ppm in/on cereal grains group (excluding wheat, barley, and oats) and 1 ppm in/on forage, fodder, and straw of cereal grains group (excluding wheat, barley, and oats) be established concomitant with the revocation of the group tolerance for grain crops (PP#1F3993; CBRS No. 10833, DP Barcode D183914, 9/30/93, D. Miller).
33. CBRS No. 8043, DP Barcode D164655, 6/1/92, L. Cheng.
34. CBRS No. 8166, DP Barcode D165608, 4/16/92, J. Abbotts.
35. The tolerance of 0.1 ppm for grasses, forage was established to cover seed treatment. Ciba-Geigy has petitioned for crop group tolerances for grasses, forage at 10 ppm and grasses, hay at 20 ppm to cover proposed soil and foliar uses. This petition is currently in reject status (PP#2F04063; CBTS No. 9338, DP Barcode D173024, 9/22/93, M. Peters).
36. A crop group tolerance for the Non-Grass Animal Feeds Group is not appropriate because of the more than 5 fold difference between residues in alfalfa and clover hays. Individual tolerances must be established. (PP#2F4105, CBTS No. 9625, DP Barcode D175974, M. Rodriguez, 8/17/93).
37. PP#9F3675, CB Nos. 4792, 4793 and 4794, 8/3/89, J. Garbus.
38. PP#6F3330, CB No. 312, 2/7/86, M. Firestone. Postemergence use of the 50% WP formulation is not supported. Supporting data must be provided or the use removed from the 50% WP label.
39. It is unclear from the submitted residue data for cotton if the de-linted or undelinted seed was analyzed. Residue data for the undelinted cottonseed are required. In addition, CBRS now requires residue data for cotton gin byproducts (commonly called gin trash) which includes burrs, leaves, stems, lint, immature seeds, sand, and dirt. As these data requirements are based on the Livestock Feeds Table (Table II) for Subdivision O (Residue Chemistry) of the Pesticide Assessment Guidelines (June 1994), they are considered confirmatory data.
40. In accordance with PR Notice 93-12 (12/23/93), the regulated raw agricultural commodity for hops is now considered to be dried hops. The 20 ppm tolerance for hops, dried listed under 40 CFR §185.4000(d) should be moved to §180.408(a). The tolerance for the commodities hops, green (0.5 ppm under §180.408[a]) and the feed additive tolerance for hops, dry (2 ppm under 40 CFR § 186.4000[a]) should be revoked. In addition, the 20 ppm tolerance for hops, spent listed under 40 CFR § 186.4000(d) should be revoked as metalaxyl residues in spent hops will now be covered by the tolerance on the RAC.
41. FAP# 7H5532, CB Nos. 4078 and 4190, 9/28/88, S. Willett; and CB No. 4630, 11/17/88, S. Willett.
42. PP#9F3712, CB Nos. 4811, 4812 and 4813, 4/11/89, S. Willett; CB No. 5579, 8/18/89, F. Toghrol; and CBTS No. 9011, DP Barcode D169779, 6/19/92, J. Morales.
43. PP#8E3605, CB No. 3360, 5/5/88, F. Griffith.
44. The tolerance for peanut, vines should be revoked, as this commodity is no longer regulated because it is not a major livestock feed item.
45. CBRS No. 8043, DP Barcode D164655, 6/1/92, L. Cheng; and CBRS No. 10743, DP Barcode D183659, 11/24/92, L. Cheng.
46. CBRS No. 11238, DP Barcode D187081, 7/22/93, J. Abbotts.
47. The tolerances for pineapple fodder and forage should be revoked as these commodities are not major livestock feed items.

Table B (continued).

- 
48. The feed additive tolerance for apple, pomace, dry should be revoked, as the regulated feed item is apple pomace, wet.
  49. CBRS No. 11238, DP Barcode No. D187081, 7/22/93, J. Abbotts.
  50. CBTS No. 11708 and 11709, DP Barcode D189772 and 189764, 5/5/93, F. Griffith.
  51. The tolerance for peanut, soapstock should be revoked as this commodity is not a major livestock feed item.
  52. A feed additive tolerance of 0.5 ppm for pineapple, process residue is appropriate and should be proposed and established (CBRS No. 9596, DP Barcode D175852, 6/22/92, R. Perfetti; and CBRS No. 11238, DP Barcode D187081, 7/22/93, J. Abbotts).
  53. Metalaxyl residues were found to concentrate up to 15.3x, 2.4x, and 1.4x in dried peel, potato flakes, and potato chips, respectively. Residues did not concentrate in the wet peel. Based on a 0.5 ppm tolerance for potatoes, food additive tolerances for the processed commodities potato granules/flakes and chips be proposed at 2 ppm, concomitant with the revocation of the established 4.0 ppm tolerance for potato, processed (including potato chips). In addition, the feed additive tolerance for potato waste, dried, processed should be increased to 10 ppm from 4 ppm, and renamed as processed potato waste (CBRS No. 8043, DP Barcode D164655, 6/1/92, L. Cheng).
  54. CBRS No. 8043, DP Barcode D164655, 6/1/92, L. Cheng.
  55. The tolerance for soybean, soapstock should be revoked, as this commodity is not a major livestock feed item.
  56. Residues in molasses were found to concentrate up to 10x. Based on the 0.5 ppm tolerance for sugar beets, the feed additive tolerance for molasses listed under 40 CFR §186.4000(a) must be increased from 1.0 to 5 ppm (CB Nos. 3908 through 3912, 11/28/88, F. Griffith; and CB Nos. 5934, 5935, and 6545, 4/30/90, F. Griffith). The feed additive tolerance of 5.0 ppm listed under 40 CFR §186.4000(b) (inadvertent tolerance) should be revoked.
  57. CB Nos. 3908 through 3912, 11/28/88, F. Griffith; and CB Nos. 5934, 5935, and 6545, 4/30/90, F. Griffith.
  58. The available data indicate that residues concentrated by up to 1.5x in sunflower, meal; a 0.2 ppm feed additive tolerance for sunflower, meal is appropriate (CBRS No. 11238, DP Barcode D187081, 7/22/93, J. Abbotts).
  59. CBRS No. 8043, DP Barcode D164655, 6/1/92, L. Cheng.
  60. CBRS No. 11238, DP Barcode D187081, 7/22/93, J. Abbotts.
  61. Data are required to determine if residues concentrate in tomato paste (Table II, June, 1994)
  62. Confined Rotational Crop studies have been submitted and are currently under review.
  63. Partially reviewed in CB No. 12312, DP Barcode No. D193692, 8/16/93, G. Herndon.

## TOLERANCE REASSESSMENT SUMMARY

### Tolerances Listed Under 40 CFR § 180.408 (a):

The tolerances listed in 40 CFR § 180.408(a) are for the combined residues of metalaxyl and its metabolites containing the 2,6-dimethylaniline moiety, and *N*-(2-hydroxy methyl-6-methylphenyl)-*N*-(methoxyacetyl)-alanine methylester, each expressed as metalaxyl.

Sufficient data are available to ascertain the adequacy of the established tolerances listed in 40 CFR § 180.408(a) for the following commodities: alfalfa, almonds, apples, asparagus, avocados, blueberries, broccoli, cabbage, cauliflower, cereal grains (excluding wheat, barley, and oats) group, citrus fruits group, cranberries, cucurbit vegetables group, fruiting vegetables (except cucurbits) group, ginseng, grapes, leafy vegetables (excluding brassica) group, leaves of root and tuber vegetables group, legume vegetables (succulent or dried group, excluding soybeans), foliage of legume vegetables group, onions (bulb and green), peanuts, pineapples, potatoes, raspberries, root and tuber vegetables (excluding ginseng) group, soybeans, spinach, stone fruits group, strawberries, sunflowers, and walnuts.

The established tolerances for green hops (0.2 ppm), peanut vines (20 ppm), and pineapple forage and fodder (0.1 ppm) should be revoked as these are no longer regulated RACs.

The established tolerances for beets (0.1 ppm), potatoes (0.5 ppm), and sugar beets (0.1 ppm and 0.5 ppm) should be revoked as these commodities are covered by the established root and tuber vegetable group tolerance (0.5 ppm). The established root and tuber group should exclude ginseng which has an established 3.0 ppm tolerance. In addition, the established tolerances for beet tops (0.1 ppm) and sugar beet tops (10.0 ppm) should be revoked as they are covered under the established crop group tolerance for leaves of root and tuber vegetables (15.0 ppm).

The established tolerance of 0.1 ppm for the Brassica (cole) leafy vegetable group (excluding broccoli, cabbage, and cauliflower) should be revoked as this tolerance was established to cover residues resulting from seed treatments which are no longer registered uses. Separate tolerances will be required for each member of the Brassica leafy vegetable group, because of the greater than five fold difference in residue levels. The available residue data indicate that the established tolerances for cabbage and cauliflower (2 ppm) should be lowered to 1 ppm.

A petition (PP#0F3893) proposing a crop group tolerance for leafy vegetables (except Brassica, excluding spinach) has been submitted. Sufficient data have been submitted to ascertain the adequacy of the pending tolerances. The established tolerance for the leafy vegetables group (excluding Brassica) should be increased from 0.1 ppm to 5 ppm and the group should be renamed to also exclude spinach. Once the increased tolerance is established, the established tolerance for head lettuce (5 ppm) should be revoked.

A petition (PP#1F3993) for crop group tolerances for the cereal grains group (excluding wheat, barley, and oat) and the forage, fodder, and straw of cereal grain group (excluding wheat, barley, and oat) has been submitted. Sufficient data have been submitted to ascertain the adequacy of the pending tolerances. A 0.1 ppm tolerance must be established for the cereal grains group (excluding wheat, barley, and oats), and a tolerance of 1.0 ppm must be established for the forage, fodder, and straw of cereal grains group (excluding wheat, barley, and oats).

Additional data are required to support the 0.1 ppm tolerances for cottonseed.

The adequacy of the established tolerances for meat, fat, and meat byproducts of cattle, goats, horses, sheep and poultry and for milk and eggs will be determined once questions pertaining to the analytical method are resolved and the submitted ruminant feeding studies are re-evaluated.

Tolerances Listed Under 40 CFR § 180.408(b):

The tolerances listed in 40 CFR § 180.408(b) are for the indirect or inadvertent residues of metalaxyl present as a result of the application of metalaxyl to the growing of the primary crops listed in 40 CFR § 180.408(a) and tobacco.

Sufficient data are available to ascertain the adequacy of the established indirect or inadvertent tolerances listed in 40 CFR § 180.408(b). Additional field rotational crop studies have been submitted and are currently under review.

Tolerances Listed Under 40 CFR § 180.408(c):

The tolerance listed in 40 CFR § 180.408(c) are with regional registration for the combined residues of metalaxyl and its metabolites containing the 2,6-dimethylaniline moiety, and *N*-(2-hydroxy methyl-6-methylphenyl)-*N*-(methoxyacetyl)-alanine methylester, each expressed as metalaxyl in/on papaya in HI.

Sufficient data are available to ascertain the adequacy of the established tolerances listed in 40 CFR § 180.408(c) for papaya.

Tolerances Listed Under 40 CFR § 185.4000(a):

The food additive tolerances listed in 40 CFR § 185.4000(a) are for the combined residues of metalaxyl and its metabolites containing the 2,6-dimethylaniline moiety, and *N*-(2-hydroxy methyl-6-methylphenyl)-*N*-(methoxyacetyl)-alanine methylester, each expressed as metalaxyl.

Sufficient data are available to ascertain the adequacy of the established food additive tolerances listed in 40 CFR § 185.4000(a) for the following food commodities: citrus oil, dried prunes, raisins, potatoes processed (including potato chips), and tomato puree.

The established tolerance for dried apricots (4 ppm) should be revoked as it is not a regulated commodity.

A 2.0 ppm food additive tolerance for potato granules/flakes and potato chips should be proposed. Once these tolerances are established, the food additive tolerance of 4.0 ppm for potatoes, processed (including potato chips) should be revoked.

Tolerances Listed Under 40 CFR § 185.4000(b):

The food additive tolerances listed in 40 CFR § 185.4000(b) are for the indirect or inadvertent residues of metalaxyl present as a result of the application of metalaxyl to the growing of the primary crops listed in 40 CFR § 180.408(a) and tobacco.

Sufficient data are available to ascertain the adequacy of the established inadvertent or indirect tolerances listed in 40 CFR § 185.4000(b). Additional field rotational crop studies have been submitted and are currently under review.

Tolerances Listed Under 40 CFR § 185.4000(d):

The food additive tolerance listed in 40 CFR § 185.4000(d) is for the combined residues of metalaxyl and its metabolites containing the 2,6-dimethylaniline moiety, and *N*-(2-hydroxymethyl-6-methylphenyl)-*N*-(methoxyacetyl) alanine methylester, each expressed as metalaxyl.

Sufficient data are available to ascertain the adequacy of the established food additive tolerance listed in 40 CFR § 185.4000(d) for dried hops. However, as the regulated RAC for hops is now considered to be dried hops [*Source: PR Notice 93-12 (12/23/93)*], the 20 ppm tolerance for hops, dried should be moved to 40 CFR §180.408(a).

Tolerances Listed Under 40 CFR § 186.4000(a):

The feed additive tolerances listed in 40 CFR § 186.4000(a) are for the combined residues of metalaxyl and its metabolites containing the 2,6-dimethylaniline moiety, and *N*-(2-hydroxy methyl-6-methylphenyl)-*N*-(methoxyacetyl)-alanine methylester, each expressed as metalaxyl.

Sufficient data are available to ascertain the adequacy of the established feed additive tolerances listed in 40 CFR § 186.4000(a) for the following feed commodities: apple pomace (wet), citrus molasses and dried pulp, grape pomace (wet and dried), peanut meal, processed potato waste, raisin waste, soybean hulls and meal, sugar beet molasses, and tomato pomace (dried).

The tolerance for potato waste, dried, processed should be increased from 4.0 ppm to 10 ppm and renamed potatoes, waste from processing. The tolerance for sugar beet molasses should be increased from 1.0 ppm to 5.0 ppm concomitant with the revocation of the 5.0

ppm for indirect or inadvertent residues in sugar beet molasses listed under 40 CFR 186.4000 (b).

The established 2.0 ppm tolerances for apple pomace (dry), hops (dry), peanut soapstock, and soybean soapstock and the 5.0 ppm tolerance for legume vegetable cannery waste should be revoked as these commodities are no longer being regulated because they are not considered major livestock feed items.

Feed additive tolerances must be proposed and established for pineapple process residue (0.5 ppm) and sunflower meal (0.2 ppm).

Tolerances Listed Under 40 CFR § 186.4000(b):

The feed additive tolerances listed in 40 CFR § 186.4000(b) are for the indirect or inadvertent residues of metalaxyl present as a result of the application of metalaxyl to the growing of the primary crops listed in 40 CFR § 180.408(a) and tobacco.

Sufficient data are available to ascertain the adequacy of the established indirect or inadvertent tolerances listed in 40 CFR § 186.4000(b) for the following feed commodities: barley, oats, and wheat, milling fractions.

The indirect or inadvertent tolerance for sugar beet molasses of 5 ppm should be revoked as the need for this tolerance is superseded by the 5 ppm tolerance for this commodity in 40 CFR § 186.4000(a).

Tolerances Listed Under 40 CFR § 186.4000(d):

The tolerances listed in 40 CFR § 186.4000(d) are for the combined residues of metalaxyl and its metabolites containing the 2,6-dimethylaniline moiety, and *N*-(2-hydroxymethyl-6-methylphenyl)-*N*-(methoxyacetyl) alanine methylester, each expressed as metalaxyl.

The 20 ppm tolerance for spent hops should be revoked as residues in spent hops will be covered by the 20 ppm tolerance on the RAC (hops, dried).

Table C. Tolerance Reassessment Summary for Metalaxyl

Commodity	Current Tolerance (ppm)	Tolerance Reassessment (ppm)	Comment/Correct Commodity Definition
<b>Tolerances listed under 40 CFR §180.408(a)</b>			
Alfalfa, forage	6.0	6.0	
Alfalfa, hay	20.0	20.0	
Almonds	0.5	0.5	
Almonds, hulls	10.0	10.0	
Apples	0.2	0.2	
Asparagus	7.0	7.0	
Avocados	4.0	4.0	
Beets	0.1	Revoke	Established 0.1 ppm tolerance was based on seed use which has been superseded by preplant and foliar uses. In addition, tolerance is covered by 0.5 ppm crop group tolerance for <i>root and tuber vegetables (exc. ginseng)</i> .
Beet, top	0.1	Revoke	Tolerance covered by 15.0 ppm crop group tolerance for <i>leaves of root and tuber vegetables</i> .
Blueberries	2.0	2.0	
Brassica (cole) leafy vegetables group [excluding broccoli, cabbage, and cauliflower]	0.1	Revoke	Separate tolerances are required for each member of the Brassica vegetable crop group for which metalaxyl use is registered.
Broccoli	2.0	2.0	
Cabbage	2.0	1.0	
Cauliflower	2.0	1.0	
Cattle, fat	0.4	To be determined	Additional data may be required.
Cattle, kidney	0.4		
Cattle, liver	0.4		
Cattle, meat	0.05		
Cattle, mbyp (excluding kidney and liver)	0.05		
Citrus fruit	1.0	1.0	<i>Citrus fruits group</i>
Cottonseed	0.1	To be determined	<i>Cotton, undelinted seed</i> Additional residue data are required.
Cranberry	4.0	4.0	<i>Cranberries</i>

Table C (continued).

Commodity	Current Tolerance (ppm)	Tolerance Reassessment (ppm)	Comment/Correct Commodity Definition
Cucurbit vegetables group	1.0	1.0	
Eggs	0.05	To be determined	Additional data may be required.
Fruiting vegetables (excluding cucurbits) group	1.0	1.0	
Ginseng	3.0	3.0	
Goats, fat	0.4	To be determined	Additional data may be required.
Goats, kidney	0.4		
Goats, liver	0.4		
Goats, meat	0.05		
Goats, mbyop (excluding kidney and liver)	0.05	To be determined	Additional data may be required.
Grain, crops	0.1	0.1	<i>Cereal grains (exc. wheat, barley, and oats).</i>
Grapes	2.0	2.0	
Grasses, forage	0.1	0.1	
Hogs, fat	0.4	To be determined	Additional data may be required.
Hogs, kidney	0.4		
Hogs, liver	0.4		
Hogs, meat	0.05		
Hogs, mbyop (excluding kidney and liver)	0.05		
Hops, green	2.0	Revoke	Hops, green is no longer a regulated RAC.
Horses, fat	0.4	To be determined	Additional data may be required.
Horses, kidney	0.4		
Horses, liver	0.4		
Horses, meat	0.05		
Horses, mbyop (excluding kidney and liver)	0.05		
Leafy vegetables (excluding brassica) group	0.1	5.0	<i>Leafy vegetables (exc. Brassica, exc. spinach)</i>

Table C (continued).

Commodity	Current Tolerance (ppm)	Tolerance Reassessment (ppm)	Comment/Correct Commodity Definition
Lettuce, head	5.0	Revoke	Tolerance will be covered by 5.0 ppm crop group tolerance for leafy vegetables (exc. Brassica, exc. spinach).
Leaves of root and tuber vegetables (human food or animal feed) group	15.0	15.0	
Legume vegetable foliage	8.0	8.0	<i>Legume vegetables group, Foliage of</i>
Legume vegetable group (dry or succulent)	0.2	0.2	<i>Legume vegetables (succulent or dried) group, exc. soybeans</i>
Milk	0.02	To be determined	Additional data may be required.
Onions, dry bulb	3.0	3.0	
Onions, green	10.0	10.0	
Peanut, hay	20.0	20.0	
Peanut, vines	20.0	Revoke	Peanut vines are no longer a regulated RAC.
Peanut, Nuts	0.2	0.2	<i>Peanuts, nutmeat</i>
Peanut, shells	2.0	2.0	<i>Peanuts, hulls (shells)</i>
Pineapples	0.1	0.1	
Pineapple fodder	0.1	Revoke	Pineapple fodder and forage are no longer regulated RACs.
Pineapple forage	0.1	Revoke	
Potatoes	0.5	Revoke	Tolerance covered by 0.5 ppm crop group tolerance for <i>root and tuber vegetables (exc. ginseng)</i> .
Poultry, fat	0.4	To be determined	Additional data may be required.
Poultry, kidney	0.4		
Poultry, liver	0.4		
Poultry, meat	0.05		
Poultry, mbyp (excluding kidney and liver)	0.05		
Raspberries	0.5	0.5	
Root and tuber vegetables group	0.5	0.5	<i>Root and tuber vegetables (exc. ginseng)</i> .
Soybean, grain	1.0	1.0	<i>Soybeans</i>
Spinach	10.0	10.0	

Table C (continued).

Commodity	Current Tolerance (ppm)	Tolerance Reassessment (ppm)	Comment/Correct Commodity Definition
Sugar beets	0.1	Revoke	Tolerance covered by 0.5 ppm crop group tolerance for <i>root and tuber vegetables (exc. ginseng)</i> .
Sugar beet (tops)	10.0	Revoke	
Sugar beet (roots)	0.5	Revoke	
Sheep, fat	0.4	To be determined	Additional data may be required.
Sheep, kidney	0.4		
Sheep, liver	0.4		
Sheep, meat	0.05		
Sheep, mby (excluding kidney and liver)	0.05		
Stone fruit group	1.0	1.0	<i>Stone fruits group</i>
Strawberries	10.0	10.0	
Sunflowers	0.1	0.1	<i>Sunflower, seed</i>
Sunflower, forage	0.1	0.1	
Walnuts	0.5	0.5	
<b>Tolerances listed under 40 CFR §180.408(b)</b>			
Barley, grain	0.2	0.2	
Barley, fodder	2.0	Revoke	No longer regulated
Barley, forage	2.0	2.0	
Barley, straw	2.0	2.0	
Oat, fodder	2.0	Revoke	No longer regulated
Oat, forage	2.0	2.0	<i>Oats, forage</i>
Oat, grain	0.2	0.2	<i>Oats, grain</i>
Oat, straw	2.0	2.0	<i>Oats, straw</i>
Wheat, fodder	2.0	Revoke	No longer regulated
Wheat, forage	2.0	2.0	
Wheat, grain	0.2	0.2	
Wheat, straw	2.0	2.0	
<b>Tolerances listed under 40 CFR §180.408(c)</b>			
Papaya	0.1	0.1	

Table C (continued).

Commodity	Current Tolerance (ppm)	Tolerance Reassessment (ppm)	Comment/Correct Commodity Definition
<b>Food Additive Tolerances listed under 40 CFR §185.4000(a)</b>			
Apricots (dried)	4.0	Revoke	Dried apricots are not a regulated commodity.
Citrus, oil	7.0	7.0	
Potatoes, processed (including potato chips)	4.0	Revoke	See <i>potato granules/flakes</i> and <i>potato chips</i> below.
Prunes (dried)	4.0	4.0	
Raisins	6.0	6.0	<i>Grapes, raisins</i>
Tomatoes, processed	3.0	3.0	<i>Tomato, puree</i>
<b>Food Additive Tolerances listed under 40 CFR §185.4000(b)</b>			
Barley, milling fractions	1.0	1.0	
Oat milling fractions	1.0	1.0	<i>Oats, milling fractions</i>
Wheat, milling fractions	1.0	1.0	
<b>Food Additive Tolerances listed under 40 CFR §185.4000(d)</b>			
Hops, dried	20.0	Move to 40 CFR §180.408(a)	Hops, dried are now regulated as a RAC
<b>Feed Additive Tolerances listed under 40 CFR §186.4000(a)</b>			
Apple, pomace (dry)	2.0	Revoke	Apple, pomace (dry) is no longer produced as a significant livestock feed.
Apple, pomace (wet)	0.4	0.4	<i>Apple, pomace</i>
Citrus, molasses	7.0	7.0	
Citrus, pulp	7.0	7.0	<i>Citrus, pulp, dried</i>
Grape pomace (dry)	10.0	10.0	<i>Grapes, pomace, dried</i>
Grape pomace (wet)	10.0	Revoke	Covered by 10 ppm tolerance on <i>Grapes, pomace, dried</i>
Hops, dry	2.0	Revoke	Hops, dry is no longer a regulated commodity.
Legume vegetable, cannery waste	5.0	Revoke	Legume vegetable cannery waste is no longer regulated.
Peanut, meal	1.0	1.0	<i>Peanuts, meal</i>
Peanut, soapstock	2.0	Revoke	Peanut soapstock is no longer a regulated commodity.
Potato waste, dried, processed	4.0	10.0	<i>Potatoes, waste from processing</i>
Raisin waste	10.0	10.0	<i>Grapes, raisin waste</i>
Soybean, hulls	2.0	2.0	<i>Soybeans, hulls</i>

Table C (continued).

Commodity	Current Tolerance (ppm)	Tolerance Reassessment (ppm)	Comment/Correct Commodity Definition
Soybean, meal	2.0	2.0	<i>Soybeans, meal</i>
Soybean, soapstock	2.0	Revoke	Soybean soapstock is no longer a regulated commodity.
Sugar beet molasses	1.0	5.0	<i>Sugar beets, molasses</i>
Tomato pomace (dry and wet)	20.0	20.0	<i>Tomato, pomace, dried</i>
<b>Feed Additive Tolerances listed under 40 CFR §186.4000(b)</b>			
Barley, milling fractions	1.0	1.0	
Oat milling fractions	1.0	1.0	<i>Oats, milling fractions</i>
Sugar beet molasses	5.0	Revoke	Inadvertent tolerance covered by primary use tolerance listed under 40 CFR §186.4000(a).
Wheat, milling fractions	1.0	1.0	
<b>Feed Additive Tolerances listed under 40 CFR §186.4000(d)</b>			
Hops, spent	20	Revoke	Tolerance will be covered by the 20.0 ppm tolerance on the RAC, <i>Hops, dried</i> .
<b>Tolerances required under 40 CFR §180.408(a)</b>			
Forage, fodder, and straw of cereal grains group (excluding wheat, barley, and oats)	None	1.0	Must be established.
<b>Food Additive Tolerances required under 40 CFR §185.4000(a)</b>			
Potatoes, granules/flakes	None	2.0	Must be proposed.
Potatoes, chips	None	2.0	Must be proposed.
<b>Feed Additive Tolerances required under 40 CFR §186.4000(a)</b>			
Pineapples, process residue	None	0.5	Must be established.
Sunflower seeds, meal	None	0.2	Must be established.

**CODEX HARMONIZATION**

Numerous maximum residue limits (MRLs) for metalaxyl residues in plant commodities have been established by Codex. Codex currently regulates metalaxyl *per se* in plant commodities; however, CBTS has concluded that the U.S tolerance definition for plant commodities should include metalaxyl and its metabolites that can be converted to 2,6-dimethylaniline and N-(2-hydroxymethyl-6-methylphenyl)-N-(methoxyacetyl) alanine methyl ester. The Codex MRLs and the applicable U.S. tolerances are presented in Table D.

Table D. Codex MRLs and Applicable U.S. Tolerances.

Commodity	MRL (mg/kg) <sup>a</sup>	U.S. Tolerance (ppm) <sup>b</sup>	Recommendation/ Comments <sup>c</sup>
Apple	0.05 <sup>d</sup>	0.2	--
Asparagus	0.05 <sup>d</sup>	7.0	--
Avocado	0.2	4.0	--
Broccoli	0.5	2.0	--
Brussels sprouts	0.2	0.1 (Brassica leafy vegetables)	--
Cabbages, head	0.5	2.0	--
Cacao beans	0.2	None	--
Carrot	0.05 <sup>d</sup>	0.5 (Root and tuber vegetables)	--
Cauliflower	0.5	2.0	--
Cereal grains	0.05 <sup>d</sup>	0.1 (grain, crops)	--
Citrus fruits	5	1	--
Cotton seed	0.05 <sup>d</sup>	0.1	--
Cucumber	0.5	1.0 (Cucurbit vegetables)	--
Gherkin	0.5	1.0 (Cucurbit vegetables)	--
Grapes	1	None	--
Hops, dry	10	20.0 (Hops, dried)	--
Lettuce, head	2 <sup>e</sup>	5.0	--
Melons, except watermelon	0.2	1.0 (Cucurbit vegetables)	--
Onion, bulb	0.2 <sup>e</sup>	0.5	--
Peanut	0.1	0.01	--
Peas, shelled	0.05 <sup>d</sup>	0.1 (Legume vegetables)	--
Peppers	1	1.0 (Fruiting vegetables)	--
Pome fruits	1	None	--

Commodity	MRL (mg/kg) <sup>a</sup>	U.S. Tolerance (ppm) <sup>b</sup>	Recommendation/ Comments <sup>c</sup>
Potato	0.05 <sup>d</sup>	0.5	--
Raspberries, Red, Black	0.2	0.5	--
Soya bean (dry)	0.05 <sup>d</sup>	1.0	--
Spinach	2 <sup>e</sup>	10.0	--
Squash, Summer	0.2	1.0 (Cucurbit vegetables)	--
Strawberry	0.2	10.0	--
Sugar beet	0.05 <sup>d</sup>	0.5 (Sugar beet, roots)	--
Sunflower seed	0.05 <sup>d</sup>	0.1	--
Tomato	0.5	1.0 (Fruiting vegetables)	--
Watermelon	0.2	1.0 (Cucurbit vegetables)	--
Winter squash	0.2	1.0 (Cucurbit vegetables)	--

- <sup>a</sup> All metalaxyl MRLs are final (CXL), except MRLs for lettuce, onions, and spinach which are at step 8 and the MRL for strawberries which is at step 7B.
- <sup>b</sup> Based on the combined residues of metalaxyl and its metabolites that can be converted to 2,6-DMA and N-(2-hydroxymethyl-6-methylphenyl)-N-(methoxyacetyl) alanine methyl ester.
- <sup>c</sup> As the tolerance definitions for Codex MRLs and U.S. tolerances are different, harmonization of Codex MRLs and U.S. tolerances is not currently possible.
- <sup>d</sup> At or about the limit of detection.
- <sup>e</sup> MRL is temporary.

The following conclusion can be made regarding efforts to harmonize the U.S. tolerances with the Codex MRLs:

- Harmonization of Codex MRLs and U.S. tolerances for metalaxyl is not possible at the present time as the Codex and U.S. tolerance definitions are incompatible.

#### DIETARY EXPOSURE ASSESSMENT SUMMARY

Current and/or recommended tolerance levels should be used for exposure assessment. For livestock commodities, we have calculated upper bound residue estimates, correcting for the recovery by the enforcement analytical method, as measured in the metalaxyl livestock metabolism studies.

Upper Bound Residues in Meat, Milk, Poultry, and Eggs are estimated from the total radioactive residue (TRR) from existing feeding studies discussed in the Metalaxyl FRSTR

Residue Chemistry Chapter of 6/22/87. The residues found in feeding studies are corrected for enforcement method recoveries from the metalaxyl metabolism studies (S. Hummel memo of 9/15/93, CB No. 9102, DP Barcode D172350). Calculations of the dietary burden for beef cattle, dairy cattle, and poultry are presented below, followed by the calculation of the upper bound residue estimates for each tissue, milk, and eggs. The upper bound residue estimates calculated here should be used for metalaxyl dietary exposure assessment instead of the established tolerances.

Dietary Intake Analysis for Beef Cattle

=====Feeds Items=====		Feed	Feed	Feed	Dietary
Crop	Feed	%DM	Consump. Percent	Tolerance (ppm)	Burden (ppm)
Alfalfa	Hay	89%	0.0%	20.000	0.000
Almond	Hulls	90%	25%	10.000	2.778
Bean*	Forage	35%	30%	8.000	6.857
Soybean	Seed	89%	15%	1.000	0.169
Wheat	Grain	89%	0%	0.200	0.000
Wheat	Milled bypts	88%	10%	1.000	0.114
Grape	Pomace, Wet	15%	20%	10.000	13.333
Total:			100.0%	Total:	23.250

Dietary Intake Analysis For Dairy Cattle

=====Feeds Items=====		Feed	Feed	Feed	Dietary
Crop	Feed	%DM	Consump. Percent	Tolerance (ppm)	Burden (ppm)
Bean*	Forage	35%	60%	8.000	13.714
Almond	Hulls	90%	15%	10.000	1.667
Soybean	Seed	89%	20%	1.000	0.225
Wheat	Milled bypts	88%	5%	0.200	0.011
Grape	Pomace, Wet	15%	0%	10.000	0.000
Total:			100.0%	Total:	15.617

Dietary Intake Analysis for Poultry

=====Feeds Items=====		Feed	Feed	Dietary
Crop	Feed	Consump. Percent	Tolerance (ppm)	Burden (ppm)
Alfalfa	Meal	10.0%	20.000	2.000
Soybean	Seed	20.0%	1.000	0.200
Wheat	Milled bypts	45.0%	1.000	0.450
Peanut	Meal	25.0%	1.000	0.250
Total:		100.0%	Total:	2.900

Calculation of Upper Bound Residues of Metalaxyl in Meat, Milk, Poultry, and Eggs

Tissue	Feed Lvl (ppm)	Metalaxyl TRR (ppm)	% Recovery by enforcement method <sup>a</sup>	Corrected Metalaxyl Residue (ppm)	Dietary Burden (ppm)	Estimated Upper Bound Residue (ppm)
<b>Dairy Cattle</b>						
muscle	75	0.14	58	0.24	23	0.074
fat	75	<0.05	89	0.06	23	0.017
liver <sup>b</sup>	75	0.14	30	0.47	23	0.143
kidney <sup>b</sup>	75	0.13	30	0.43	23	0.133
milk	75	0.02	49	0.04	16	0.009
<b>Poultry</b>						
Eggs	5	<0.05	22	0.23	2.9	0.132
muscle	5	<0.05	26	0.19	2.9	0.112
fat	5	<0.05	18	0.28	2.9	0.161
liver	5	<0.05	19	0.26	2.9	0.153
skin	5	<0.05	18	0.28	2.9	0.161
<sup>a</sup> Recovery of Total Radioactive Residue (TRR) using the Enforcement Analytical Method  <sup>b</sup> Cattle liver and kidney samples were from 14-28 feeding periods; higher levels were reported at shorter feeding times, but cattle were slaughtered several hours after the final dose. At the 14 and 28 day dosing period, cattle were slaughtered 24 hours after the final dose.						

AGENCY MEMORANDA CITED IN THIS DOCUMENT

**CB No.:** 312  
**Subject:** PP#6F3330. Metalaxyl on Asparagus. Evaluation of Analytical Methods and Residue Data.  
**From:** M. Firestone  
**To:** H. Jacoby and Toxicology Branch (HED)  
**Dated:** 02/07/86  
**MRID(s):** 00154446

**CB No.:** 60  
**Subject:** EPA Reg. #100-629 [Accession #259156]. Metalaxyl (Ridomil®-MZ58): Amended Registration  
**From:** W. Anthony, HED  
**To:** H. Jacoby, RD  
**Dated:** 02/14/86  
**MRID(s):** 00098428

**CB No.:** 360  
**Subject:** PP#6F3337 - Metalaxyl on Strawberries--Evaluation of Analytical Methods and Residue Data (Accession Numbers 260658 and 260659).  
**From:** M. Firestone, HED  
**To:** H. Jacoby, RD, and Toxicology Branch, HED  
**Dated:** 02/21/86  
**MRID(s):** 00155237

**CB No.:** 960  
**Subject:** FAP#4H5425. Metalaxyl on Grapes. Amendment of May 19, 1986. No Accession Number.  
**From:** L. Cheng, HED  
**To:** H. Jacoby/B. Cool, RD  
**Dated:** 06/03/86  
**MRID(s):** 00138818

**CB No.:** 986  
**Subject:** PP#6F3337. Metalaxyl on Strawberries - Amendment Dated April 11, 1986  
**From:** M. Firestone  
**To:** H. Jacoby and Toxicology Branch (HED)  
**Dated:** 09/08/86  
**MRID(s):** 00155237

**CBRS No(s).:** 1676, 1677, 1678  
**Subject:** PP#7F3470. Metalaxyl in or on Blueberries, Walnuts, Almonds, Almond Hulls, Stone Fruits, Dried Apricots, and Prunes.  
**From:** M. Nelson  
**To:** L. Rossi and Toxicology Branch (HED)  
**Dated:** 03/06/87  
**MRID(s):** 00164649, 00164650, 00164651

CB No(s): 1996 and 2013  
 Subject: PP#6F3387. Metalaxyl on Fruiting Vegetables (except Curcubits), Sugar Beets and Sugar Beet Tops. Evaluation of February 26, 1987, Amendment.  
 From: F. Griffith  
 To: L. Rossi and Toxicology Branch (HED)  
 Dated: 03/13/87  
 MRID(s): 40106601

CB No.: 2604, 2699, 2700  
 Subject: PP#7F3470/FAP#7H5520, EPA Reg. No. 100-601. Metalaxyl in or on Blueberries, Walnuts, Almonds, Almond Hulls, Stone Fruits, Dried Apricots and Prunes.  
 From: M. Nelson  
 To: L. Rossi and Toxicology Branch, HED  
 Dated: 08/26/87  
 MRID(s): None

CB No.: 2301  
 Subject: PP#6F3337. Metalaxyl on Strawberries - Amendment Dated March 4, 1987.  
 From: M. Kovacs  
 To: L. Rossi and Toxicology Branch (HED)  
 Dated: 09/03/87  
 MRID: None

CB No.: 3360  
 Subject: PP#8E3605. Petition Review for Establishment of Tolerance(s). Metalaxyl on Papaya  
 From: F. Griffith  
 To: H. Jamerson and Toxicology Branch (HED)  
 Dated: 05/05/88  
 MRID: 40490500 and -01

CBRS No(s): 3788 and 3789  
 Subject: PP#7F3470/FAP#7H5520. Metalaxyl in or on Blueberries, Walnuts, Almonds, Almond Hulls, Stone Fruits, Dried Apricots, and Prunes.  
 From: M. Nelson, HED  
 To: L. Rossi, RD, and Toxicology Branch, HED  
 Dated: 05/13/88  
 MRID(s): 40503100 and -01

CB No(s): 4078 and 4190  
 Subject: FAP#7H5532. Metalaxyl (Ridomil) on Hops. Amendments of July 8 and 19, 1988.  
 From: S. Willett, HED  
 To: L. Rossi, RD, and Toxicology Branch, H-FS, HED  
 Dated: 09/28/88  
 MRID(s): 40746900 and -01

CB No.: 4630  
 Subject: FAP#7H5532. Metalaxyl (Ridomil) on Hops. Revised Sections B and F.  
 From: S. Willett, HED  
 To: L. Rossi/M. Fiol, RD, and Toxicology Branch, HED  
 Dated: 11/17/88  
 MRID(s): None

CB No.: 3908 through 3912  
 Subject: PP#8F3617/FAP#8H5554. Metalaxyl on Sugar Beet Tops and Roots; Legume Vegetables (Dry and Succulent) Crop Groups; Legume Vegetables Foliage Crop Group; Grass Forage, Fodder, and Hay Crop Group; and Nongrass Animal Feeds Crop Group. Evaluation of Analytical Methods and Residue Data  
 From: F. Griffith  
 To: L. Rossi and Toxicology Branch (HED)  
 Dated: 11/28/88  
 MRID: 40534800 through -03 and 40569301 through -03

CB No.: 4808  
 Subject: EPA Reg. No. 100-628. Ridomil 5G (Metalaxyl). Amended Registration Request to Add Use on Head Lettuce.  
 From: L. Propst, CBRS, HED  
 To: L. Rossi, RD  
 Dated: 02/09/89  
 MRID: 40775801

CB No.: 4809  
 Subject: EPA Reg. No. 100-628. Ridomil 5G (Metalaxyl). Amended Registration Request to Add Use on Spinach.  
 From: L. Propst, CBRS, HED  
 To: L. Rossi, RD  
 Dated: 02/14/89  
 MRID: 40790201

CB No(s): 4811, 4812, 4813  
 Subject: PP#9F3712. Metalaxyl on Green Hops. Petition to Revise Tolerance.  
 From: S. Willett  
 To: L. Rossi and Toxicology Branch (HED)  
 Dated: 04/11/89  
 MRID(s): 40909401

CB No(s): 4814, 4815, 4816  
 Subject: PP#6F3337. Metalaxyl on Strawberries - Amendment Dated November 4, 1988.  
 From: M. Kovacs  
 To: S. Lewis and Toxicology Branch (HED)  
 Dated: 05/10/89  
 MRID(s): 40880400 and 40880401

CB No(s): 4792, 4793, and 4794  
 Subject: PP#8F3695/FAP#8H5569: Metalaxyl: Tolerances for Alfalfa and for Barley as Rotational Crop. Evaluation of Residue Data and Analytical Methodology.  
 From: J. Garbus, CBRS, HED  
 To: L. Rossi, RD  
 Dated: 08/03/89  
 MRID(s): 40832901

**CB No(s):** 4788 to 4791  
**Subject:** PP#8F3698: Metalaxyl: Tolerances for Metalaxyl for the Crop Grouping Root and Tuber Vegetables. Evaluation of Residue Data and Analytical Methodology.  
**From:** J. Garbus, CBRS, HED  
**To:** L. Rossi, RD  
**Dated:** 08/03/89  
**MRID(s):** 40838301 to 40838303

**CB No.:** 5579  
**Subject:** OR-890008; 24(c) Registration for the Use of Metalaxyl (Ridomil 2E Fungicide) on Hops. EPA Reg. No. 100-607.  
**From:** F. Toghrol, CBRS, HED  
**To:** S. Lewis, RD  
**Dated:** 08/18/89  
**MRID(s):** None

**CB No.:** 5836  
**Subject:** PP#9F3698: [Formerly PP#8F3698]: Metalaxyl: Tolerances for the Crop Grouping: Root and Tuber Vegetables. Amendment of September 25, 1989. Revised Sections B and F for PP#8F3698.  
**From:** J. Garbus, CBRS, HED  
**To:** S. Lewis/B. Chambliss, RD  
**Dated:** 10/23/89  
**MRID(s):** None

**CBRS No.:** 5895  
**Subject:** PP#6F3337. Metalaxyl on Strawberries. Amendment of 09/27/89.  
**From:** D. Edwards  
**To:** S. Lewis  
**Dated:** 11/02/89  
**MRID(s):** None

**CBRS No.:** 5557 and 5558  
**Subject:** PP#6F3362/FAP#6H5493. Metalaxyl and Mancozeb on Grapes. Evaluation of Amendment Dated June 23, 1989.  
**From:** G. Otakie  
**To:** S. Lewis  
**Dated:** 12/11/89  
**MRID:** 41150101

**CBRS No.:** 6176  
**Subject:** PP#8F3698. Metalaxyl: Tolerances for Crop Grouping: Root and Tuber Vegetables. Amendment of December 7, 1989. Revised Section B.  
**From:** J. Garbus  
**To:** S. Lewis and B. Chambliss  
**Dated:** 02/14/90  
**MRID(s):** None

**CBRS No.:** 5934, 5935, 6545  
**Subject:** PP#8F3617/FAP#8H5554. Metalaxyl on Sugar Beet Tops and Roots; Legume Vegetables (Dry and Succulent) Crop Group; Legume Vegetables Foliage Crop Group; Grass Forage, Fodder, and Hay Crop Group; and Non-grass Animal Feeds Crop Group.  
**From:** F. Griffith  
**To:** S. Lewis and Toxicology Branch (HED)  
**Dated:** 04/30/90  
**MRID:** 41250101 and 41055203

**CBRS No.:** 7193  
**Subject:** Ciba-Geigy Corp.: Response to the Metalaxyl Reregistration Standard: Storage Stability and Sample Histories and Multiresidue Protocol Data.  
**From:** R. Perfetti, CBRS, HED  
**To:** R. Engler, HED and L. Rossi, SRRD  
**Dated:** 01/15/91  
**MRID(s):** 41449001, 41055203, and 40534803

**CBRS No.:** 7418  
**Subject:** OK-900005. Metalaxyl on Spinach. 24(c) Special Local Needs Registration for (1) Decreased PHI, (2) Additional Applications, and (3) Use of the Granular Formulation  
**From:** M. Metzger, CBRS, HED  
**To:** S. Lewis/J. Stone, RD  
**Dated:** 01/24/91  
**MRID:** 40790201

**CBRS No.:** 7165  
**Subject:** EPA Reg. No. 100-628. Ridomil 5G (Metalaxyl) Add Use on Spinach. Additional Residue Studies Submitted 9/19/90 in Response to DEB 2/14/89 Review.  
**From:** K. Dockter, CBRS, HED  
**To:** S. Lewis/B. Chambliss, RD  
**Dated:** 02/19/91  
**MRID:** 41636201

**CBTS No.:** 7340  
**Subject:** PP#1E3926. Evaluation of Analytical Method(s) and Residue Data. Metalaxyl on Ginseng.  
**From:** G. Otakie  
**To:** H. Jamerson and Toxicology Branch (HED)  
**Dated:** 02/22/91  
**MRID(s):** 41688900 and 41688901

**CBRS No(s).:** 7787  
**Subject:** PP#1E3926. Metalaxyl on Ginseng - Amendment of March 21, 1991  
**From:** G. Otakie  
**To:** H. Jamerson and Toxicology Branch (HED)  
**Dated:** 04/08/91  
**MRID(s):** None

**CBTS No.:** 9011  
**DP Barcode:** D169779  
**Subject:** PP#1F3993. Metalaxyl. Tolerance Petition for Residues in Cereal Grains, Soybeans, Potatoes, Peanuts, Cottonseed, Sunflower Seed, and Hops. Evaluation of Residue Data and Analytical Methodology. HED # 2-0630.  
**From:** J. Morales, CBTS, HED  
**To:** S. Lewis, RD, and Toxicology Branch, HED  
**Dated:** 06/19/91  
**MRID(s):** 41870101, -02, -03, -04, -05, -06, -07, and -08

**CBRS No(s):** 7171  
**Subject:** PP#0F3893. Metalaxyl Technical: (Ridomil®2E). Tolerance in/on Leafy Vegetables (Excluding Brassica Vegetables, Excluding Spinach). Analysis of Residue Data and Analytical Method. f`  
**From:** S. Bacchus  
**To:** B. Chambliss and S. Lewis  
**Dated:** 06/21/92  
**MRID(s):** 41587800 and 41587801

**CBRS No(s):** 7646 and 7647  
**Subject:** PP#8F3362/FAP#6H5493-Metalaxyl on Grapes--Evaluation of Amendment Dated January 25, 1991  
**From:** G. Otakie, CBTS, HED  
**To:** S. Lewis, RD, and Toxicology Branch, HED  
**Dated:** 09/09/91  
**MRID(s):** 41664503 through 41664506

**CBTS No.:** 8582  
**DP Barcode:** D168812  
**Subject:** PP#1E4024. Metalaxyl on Cranberries. Evaluation of Residue Data and Analytical Methodology.  
**From:** J. Morales, CBTS, HED  
**To:** H. Jamerson, RD, and Toxicology Branch, HED  
**Dated:** 10/31/91  
**MRID:** 41996401

**CBRS No.:** 8625  
**DP Barcode:** D169183  
**Subject:** ID#100-TEN. Ridomil®/Copper 70W Fungicide (Metalaxyl/Copper Hydroxide) "Me Too." For Foliar Use on Carrots, Cucurbits, Peppers, Potatoes, Radishes, Spinach, and Tomatoes. Accession Nos. 251021 and 262112.  
**From:** K. Dockter, CBRS, HED  
**To:** C. Giles-Parker, RD  
**Dated:** 12/12/91  
**MRID:** 41718301

CBTS No.: 8898  
DP Barcode: D171050  
Subject: PP#0F3893: Metalaxyl (Ridomil 2E, 5G) In/On Leafy Vegetables (Excluding Spinach). Amendment of 9/9/91.  
From: R. Lascola, CBTS, HED  
To: B. Chambliss/S. Lewis, RD  
Dated: 12/18/91  
MRID(s): 42021101

CBRS No.: 9054 and 9058  
Subject: EPA Reg. Nos. 100-628, 100-607. Amended Registration of Metalaxyl in/on Ginseng.  
From: W. Chin  
To: S. Lewis  
Dated: 03/06/92  
MRID(s): 41688900 and 41688901

CBRS No.: 8166  
DP Barcode: D165608  
Subject: Metalaxyl. Ciba-Geigy Response to the Reregistration Standard: Product Chemistry Data, Storage Interval Data, and Corn Fodder Data.  
From: J. Abbotts, CBRS, HED  
To: L. Rossi, SRRD  
Dated: 04/16/92  
MRID(s): 41912901, -02, and 03

CBTS No.: 9710  
DP Barcode: D176810  
Subject: PP#0F3893. Metalaxyl (Ridomil 2E, 5G) In/On Leafy Vegetables (Excluding Brassica, Excluding Spinach). Amendment of 3/17/92.  
From: R. Lascola, CBTS, HED  
To: B. Chambliss/S. Lewis, RD  
Dated: 05/12/92  
MRID(s): None

CBRS No.: 8043  
DP Barcode: D164655  
Subject: Metalaxyl. Ciba-Geigy Response to Guidance Document (FRSTR) Dated 9/88.  
From: L. Cheng, CBRS, HED  
To: L. Rossi, SRRD  
Dated: 06/01/92  
MRID(s): 41870301 through -07

CBRS No.: 9596  
DP Barcode: D175852  
Subject: Ciba-Geigy Corp.: Response to the Metalaxyl Reregistration Standard: Residue and Processing Data  
From: R. Perfetti, CBRS, HED  
To: W. Burnam, HED/L. Rossi, SRRD  
Dated: 06/22/92  
MRID: 42233501

CBTS No.: 9537  
 DP Barcode: D175506  
 Subject: PP#1E4024. Metalaxyl on Cranberries. Amendment to Review of 10/31/91.  
 From: J. Morales, CBTS, HED  
 To: H. Jamerson, RD, and Toxicology Branch, HED  
 Dated: 07/17/92  
 MRID(s): None

CBRS No(s): 12312  
 DP Barcode(s): D193692  
 Subject: Registration for Application of Ridomil MZ58® (Metalaxyl + Mancozeb, EPA Reg #100-629) to Potatoes with a Rotation to Corn. Request to Change the Rotational Crop Restriction on Corn from 12 to 9 Months. Submission of the Rotational Crop Data Requested in the CBTS Memo of 05/05/93.  
 From: G. Herndon  
 To: S. Lewis and D. Greenway  
 Dated: 08/16/93  
 MRID(s): 41870308

CBRS No(s): 10743 and 10744  
 DP Barcode(s): D183657 and D183659  
 Subject: Metalaxyl. Label Amendments for Peanuts. Ridomil PC 11G (EPA Reg. No. 100-664), Ridomil PC (EPA Reg. No. 100-713)  
 From: L. Cheng, CBRS, HED  
 To: B. Chambliss, RD  
 Dated: 11/24/92  
 MRID(s): None

CBRS No.: 9342  
 Subject: PP#2F04072: 1. Metalaxyl (Ridomil® 2E Fungicide, EPA Reg. No. 100-607) In or On Members of the Brassica (Cole) Leafy Vegetables Crop Grouping; Evaluation of Request for Amended Registration, Analytical Method, and Residue Data of Mustard Greens.  
 From: M. Peters, CBTS, HED  
 To: B. Chambliss/S. Lewis, RD  
 Dated: 01/15/93  
 MRID(s): None

CBTS No(s): 11708 and 11709  
 DP Barcode(s): D189772 and D189764  
 Subject: LA-920004 Metalaxyl (Ridomil®) - Special Local Need [24(c)] Registration and Amended Use Registration for EPA Registration Number 100-607 (Ridomil® 2E Fungicide) on Water Seeded Rice. Review of Amendment with New Magnitude of the Residue Data - Crop Field Trials.  
 From: F. Griffith, CBTS, HED  
 To: J. Fairfax/S. Lewis, RD  
 Dated: 05/05/93  
 MRID(s): 42259805

**CBRS No.:** 11238  
**DP Barcode:** D187081  
**Subject:** Metalaxyl Reregistration. Registrant Ciba-Geigy Submission on Magnitude of the Residue in Peanuts and Processed Commodities of Rice, Sunflower, and Pineapple.  
**From:** J. Abbotts, CBRS, HED  
**To:** L. Rossi, SRRD  
**Dated:** 07/22/93  
**MRID(s):** 42498701 and -02

**CBRS No.:** None  
**Subject:** Metalaxyl (113501) Ruminant and Poultry Metabolism: HED Metabolism Committee Meeting Held September 8, 1993.  
**From:** S. Hummel, CBRS, HED  
**To:** HED Metabolism Committee  
**Dated:** 09/10/93  
**MRID(s):** None

**CBRS No.:** 9102  
**DP Barcode:** D172350  
**Subject:** Metalaxyl (113501) Additional Ruminant and Poultry Metabolism Data Poultry Feeding Study.  
**From:** S. Hummel, CBRS, HED  
**To:** C. Peterson/W. Waldrop, SRRD  
**Dated:** 09/15/93  
**MRID(s):** 42115801 to -10

**CBTS No.:** 9338  
**DP Barcode:** D173024  
**Subject:** PP#2F04063, Metalaxyl In or On the Grass Forage, Fodder, and Hay Crop Grouping. Evaluation of Analytical Method and Residue Data (Ridomil® 2E Fungicide, EPA Reg. No. 100-607).  
**From:** M. Peters, CBTS, HED  
**To:** B. Chambliss, RD  
**Dated:** 09/22/93  
**MRID(s):** 42134500 and -01

**CBRS No.:** 10833  
**DP Barcode:** D183914  
**Subject:** PP#1F3993. Proposed Metalaxyl Tolerances  
**From:** D. Miller, SA, HSO, USPHS  
**To:** B. Chambliss, RD  
**Dated:** 9/30/93  
**MRID(s):** None

**CBRS No.:** 12888  
**DP Barcode:** D197066  
**Subject:** Metalaxyl (113501) Storage Stability Data Interim.  
**From:** S. Hummel, CBRS, HED  
**To:** P. Perreault/L. Propst, SRRD  
**Dated:** 02/17/94  
**MRID(s):** 42919401

MASTER RECORD IDENTIFICATION NUMBERS

- 00071601 Fischer, W.C.; Cassidy, J.E. (1978) Balance and Metabolism of [Phenyl]-14C-CGA-48988 in Potatoes: M6-69-1P, 1S; Report No. ABR-78001. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-A)
- 00071602 Foster, R.A.; Fischer, W.C.; Cassidy, J.E. (1978) Uptake of [Phenyl]-14C-CGA-48988 in Potatoes Grown in a Field Plot - Preparation of Rotational Plots: M6-69-2P, 2S; Report No. ABR-78013. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-B)
- 00071603 Honeycutt, R.C.; Fischer, W.C.; Madrid, S.; et al. (1980) Uptake, Balance and Metabolism of [Phenyl]-14C-CGA-48988 in Field Grown Potatoes: M11-69-14P, 14S; Report No. ABR-80042. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-C)
- 00071604 Honeycutt, R.C.; Cassidy, J.E. (1980) Translocation of [Phenyl]-14C-CGA-48988 and Metabolites from Leaves into Potato Tubers: M11-69-12P, 12S; Report No. ABR-80048. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-D)
- 00071605 Gross, D. (1977) Metabolism of CGA 48 988 in Field Grown Potato Plants: Project Report 30/77. (Unpublished study received Apr 15, 1981 under 100-607; prepared by Ciba-Geigy Ltd., Switzerland, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-F)
- 00071606 Gross, D. (1978) Metabolism of CGA 48 988 in Grapevine: Project Report 11/78. (Unpublished study received Apr 15, 1981 under 100-607; prepared by Ciba-Geigy Ltd., Switzerland, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-F)
- 00071607 Gross, D. (1979) Identification of Metabolites of CGA 48 988 (Ridomil® in Grapevine: Project Report 06/79. (Unpublished study received Apr 15, 1981 under 100-607; prepared by Ciba-Geigy Ltd., Switzerland, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-G)
- 00071608 Gross, D. (1979) Fate of CGA 48 988 in Lettuce: Project Report 38/79. (Unpublished study received Apr 15, 1981 under 100-607; prepared by Ciba-Geigy Ltd., Switzerland, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-H)
- 00071609 Gross, D. (1979) Identification of Metabolites of CGA 48 988 (Ridomil®) in Field Grown Potato Plants: Project Report 39/79. (Unpublished study received Apr 15, 1981 under 100-607; prepared by Ciba-Geigy Ltd., Switzerland, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-I)
- 00071610 Gross, D. (1980) Identification of Degradation Products of CGA 48988 (Ridomil®) in Lettuce: Project Report 38/80. (Unpublished study received Apr 15, 1981 under 100-607; prepared by Ciba-Geigy Ltd., Switzerland, submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-J)
- 00071611 Seim, V.W. (1978) Biological Report for the Metabolism of [Phenyl]-14C-CGA-48988 in a Lactating Goat: Report No. BIOL-78002. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-K)

- 00071612 Fischer, W.C.; Foster, R.A.; Cassidy, J.E. (1978) Balance and Metabolism of [Phenyl]-<sup>14</sup>C-CGA-48988 in a Lactating Goat: M6-69-3A: Report No. ABR-78046. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070018-L)
- 00071615 Ciba-Geigy Corporation (1980) [Study of Metalaxyl (Ridomil 2E) Residue Tolerances in Cottonseed, Soybeans, Wheat, and Certain Vegetable Crops]: AG-A 5457 I. (Compilation; unpublished study, including AG-A 5725, 5456 I, 5545 I, II..., received Apr 15, 1981 under 100-607; CDL:070019-A)
- 00071616 Ciba-Geigy Corporation (1981) [Study of Various Compounds for Residue Tolerances in Potatoes]: AG-A 4601. (Compilation; unpublished study, including AG-A 4614, 4615, 4903..., received Apr 15, 1981 under 100-607; CDL:070020-A)
- 00071622 Balasubramanian, K. (1980) Analytical method for the Determination of Total Residues of Metalaxyl in Animal Tissues, Milk and Eggs as 2,6-Dimethylaniline. Method no. AG-349 dated Nov 25, 1980. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070022-L)
- 00071623 Balasubramanian, K. (1980) Analytical Method for the Determination of Total Residues of Metalaxyl in Oil Seed Fractions as 2,6-Dimethylaniline. Method no. AG-350 dated Dec 18, 1980. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070022-M)
- 00071672 Ciba-Geigy Corporation (1981) [Series of Tests Performed with Metalaxyl on Various Crops]: AG-A 5638. (Compilation; unpublished study, including AG-A nos. 5639 I,II, 5754 I,II, 5805 I,II, ..., received Apr 15, 1981 under 100-607; CDL:070021-A)
- 00071673 Seim, V.W.; Thomas, W.A. (1980) Biological Report For CGA-48988: Residue Test in Laying Hens: Report No. Biol-80009. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070021-B)
- 00071674 Seim, V.; Thomas, W.; Brown, G. (1980) Biological Report for CGA-48988: Residue Test in Lactating Cows: Report No. Biol-79009. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070021-C)
- 00071676 Balasubramanian, K.; Ross, J.A. (1981) Comparison of Analyses of Potato Tubers Treated with Metalaxyl Using the Parent Method AG-325 and the Total Method AG-348: Report No. ABR-80056. (Unpublished study received Apr 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070021-E)
- 00074488 Ciba-Geigy Corporation (1981) Residues of Metalaxyl in Avocados Resulting from Soil Surface Granular, Soil Drench, and Drip Irrigation Applications: Report No. ABR-81016. (Compilation; unpublished study received Jun 9, 1981 under 100-607; CDL:070151-A)
- 00079433 Kahrs, R.A. (1981) Residues of Metalaxyl in Hops Resulting from Soil Drench Applications: Report No. ABR-81024. (Unpublished study received Jul 15, 1981 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070188-A)
- 00097511 Ciba-Geigy Corporation (1981) [Residues of Ridomil® in Lettuce]. (Compilation; unpublished study received Apr 1, 1982 under 100-EX-71; CDL:070732-A)
- 00098428 Kahrs, R.A. (1982) Metalaxyl - Melons, Cucumbers, Squash, Green Onions, Bulb Onions, Potatoes - Supportive Data Requested by EPA: Report No. ABR-82014. (Unpublished study received Apr 7, 1982 under 1F2500; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:070765-A)

- 00100467 Honeycutt, R.; Cassidy, J. (1978) Uptake and Balance of 0-14C-CGA-48988 and Its Metabolites in Greenhouse Grown Bright and Burley Tobacco: Report no. ABR-78036. (Unpublished study received Apr 26, 1979 under 100-600; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:238232-D)
- 00100476 Cassidy, J.; Ross, J. (1979) Metalaxyl (CGA-48988) - Tobacco: Results of Tests on the Amounts and Nature of Metalaxyl Residues in Tobacco and Its Smoke: Report No. ABR-79103. Summary of studies 241245-A, 241245-B, 241246-A. (Unpublished study received Oct 31, 1979 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:241244-A)
- 00100477 Honeycutt, R.; Szolics, I.; Simoneaux, B.; et al. (1979) Identification of the Major Aglycones of [Phenyl]-14C-CGA-48988 Conjugated Metabolites in Cured Greenhouse Grown Bright Tobacco, M11-69-2Y: Report No. ABR-79008. (Unpublished study received Oct 31, 1979 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:241245-A)
- 00100478 Honeycutt, R.; Fischer, W.; Cassidy, J. (1979) Uptake and Balance of [phi]-14C-CGA-48988 and Its Metabolites in Field Grown Bright Tobacco, M11-69-11P, 11S: Report No. ABR-79100. (Unpublished study received Oct 31, 1979 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:241245-B)
- 00100753 Ciba-Geigy Corp. (1982) [Residues of Metalaxyl in Cattle and Goats]. (Compilation; unpublished study received May 11, 1982 under 1F2500; CDL:070836-A)
- 00104378 Balasubramanian, K.; Nixon, W. (1978) Gas Chromatographic Residue Determination of CGA-48988 in Crop Samples: Method No. AG-325. (Unpublished study received Oct 31, 1979 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:241243-D)
- 00104387 Ciba-Geigy Corp. (1979) [Residues of CGA-48988 on Wheat, Rye, Forage and Other Crops]. (Compilation; unpublished study received Oct 31, 1979 under 100-607; CDL:241243-N)
- 00104390 Ciba-Geigy Corp. (1979) [Residue Studies of CGA-48988 on Soybeans and Other Crops]. (Compilation; unpublished study received Oct 31, 1979 under 100-607; CDL:241243-Q)
- 00104485 Honeycutt, R.; Simoneaux, B.; Szolics, I.; et al. (1978) Characterization of Polar Metabolites of [Phenyl]-14C-CGA-48988 in Greenhouse Grown Bright Tobacco: Report No. ABR-78044. (Unpublished study received Jul 13, 1978 under 100-EX-62; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:234431-C)
- 00104654 Ross, J.; Marco, G. (1978) CGA-48988 - Potatoes Residues Observed and Metabolism Data Including the Analytical Methods Used: Report No. ABR-78055. (Unpublished study received Sep 5, 1978 under 100-EX-1; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:097387-A)
- 00104656 Ciba-Geigy Corp. (1978) [Ridomil 2E Fungicide Residue Study - Potatoes]. (Compilation; unpublished study received Sep 5, 1978 under 100-EX-1; CDL:097389-A)
- 00109402 Ciba-Geigy Corp. (1981) Metalaxyl - Cotton In-furrow Applications of Ridomil 2E and 5G Including Mixtures with PCNB: Report No. ABR-81046. (Compilation; unpublished study received Jul 28, 1982 under 100-607; CDL:247934-A)
- 00109472 Ciba-Geigy Corp. (1982) Metalaxyl Residues in Pineapple Resulting from a Preplant Dip Application of Ridomil 2E. (Compilation; unpublished study received Aug 12, 1982 under 100-607; CDL: 071034-A)

- 00114376 Ciba-Geigy Corp. (1982) Metalaxyl - Rotational Wheat: Report # ABR-82061. (Compilation; unpublished study received Sep 23, 1982 under 100-607; CDL:071104-A)
- 00114377 Ciba-Geigy Corp. (1982) Metalaxyl - Lettuce: Report #ABR-82060. (Compilation; unpublished study received Sep 23, 1982 under 100-607; CDL:071106-A)
- 00114378 Ciba-Geigy Corp. (1982) Metalaxyl - Spinach: Report #ABR-82050. (Compilation; unpublished study received Sep 23, 1982 under 100-607; CDL:071107-A)
- 00114379 Collins, P.; Williams, S.; Cassidy, J. (1982) Metabolism of [Phenyl]-14C-CGA-48988 in Greenhouse Head Lettuce: Report No. ABR-82054. (Unpublished study received Sep 23, 1982 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:071108-A)
- 00117969 Cheung, M.; Manuli, P. (1982) Metalaxyl - Citrus: Residues Resulting from Soil or Trunk-spray Applications of Ridomil 2E: Report No. ABR-82069. (Unpublished study received Nov 16, 1982 under 100-607; submitted by Ciba-Geigy Corp., Greensboro, NC; CDL:071238-A)
- 00126315 Ciba-Geigy Corp. (1983) Summary of Section D: [Metalaxyl - Apples]. (Compilation; unpublished study received Mar 7, 1983 under 100-607; CDL:071476-A)
- 00127769 Ciba-Geigy Corp. (1983) Metalaxyl - Red Raspberries: Residues Resulting from Soil Application of Ridomil 2E or 5G. (Compilation; unpublished study received Mar 7, 1983 under 100-607; CDL:071478-A)
- 00128102 Ciba-Geigy Corp. (1982) Metalaxyl (Apron 2E) Seed Treatments. (Compilation; unpublished study received Feb 3, 1983 under 100-626; CDL:071388-A)
- 00128738 Ciba-Geigy Corp. (1983) Summary of Section D: Metalaxyl - Peanuts: [Residue Reports]. (Compilation; unpublished study received Jun 29, 1983 under 100-607; CDL:071718-A)
- 00129003 Ciba-Geigy Corp. (1983) Summary of Section D: Metalaxyl - Pod Vegetables. (Compilation; unpublished study received Jun 23, 1983 under 3F2918; CDL:071711-A)
- 00130693 Ciba-Geigy Corp. (1983) Residue Summary: Metalaxyl - Cucumbers, Melons, and Squash. (Compilation; unpublished study received Aug 19, 1983 under 100-607; CDL:251019-A)
- 00130694 Ciba-Geigy Corp. (1983) Residue Summary: Metalaxyl - Green and Dry Bulb Onions. (Compilation; unpublished study received Aug 19, 1983 under 100-607; CDL:251020-A)
- 00130695 Ciba-Geigy Corp. (1983) Residue Summary: Metalaxyl - Lettuce and Spinach. (Compilation; unpublished study received Aug 19, 1983 under 100-607; CDL:251021-A)
- 00130773 Ciba-Geigy Corp. (1983) Residue Summary: Metalaxyl - Broccoli, Cabbage and Cauliflower. (Compilation; unpublished study received Aug 26, 1983 under 100-607; CDL:071918-A)
- 00133020 Ciba-Geigy Corp. (1983) Metalaxyl - Citrus Residues Resulting from Soil Applications of Ridomil 5G. (Compilation; unpublished study received Dec 8, 1983 under 100-628; CDL:251988-A)
- 00138818 Ciba-Geigy Corp. (1984) Residue Summary: Metalaxyl - Field Grapes. (Compilation; unpublished study received Jan 30, 1984 under 100-EX-81; CDL:072298-A)

- 00140371 Cassidy, J.E.; Ross, J.A. (1978) CGA-48988--Tobacco: Results of Tests on the Amounts and Nature of Residues in Tobacco and Its Smoke: Report No. ABR-78045. (Unpublished study received Jul 13, 1978 under 100-EX-62; submitted by Ciba-Geigy Corp., Greensboro, N.C.; CDL:234430-A)
- 00141519 Ciba-Geigy Corp. (1984) Metalaxyl - Apples: Residue Chemistry Data . Unpublished compilation. 77 p. (CDL:072991A, 072747A)
- 00148103 Ciba-Geigy Corp. (1985) Metalaxyl (Ridomil 2E and Ridomil MZ58) in All Crops: Comparison Studies on Ground Vs. Aerial Treatments: Residue Data. Unpublished compilation. 88 p. (CDL:259156A)
- 00148440 Ciba-Geigy Corp. (1985) Metalaxyl (Ridomil 5G) in All Crops: Cross-over Comparison Studies of Ridomil 2E and 5G. Unpublished compilation. 577 p. (CDL:257618A)
- 00154446 Ciba-Geigy Corp. (1985) Analytical Methodology and Metalaxyl Residues in Asparagus. Unpublished compilation. 85 p. (CDL:260563A)
- 00155237 Ciba-Geigy Corp. (1985) Metalaxyl - Strawberries Residue and Analytical Methodology. Unpublished compilation. 102 p. (CDL:260659A)
- 00155845 Ciba-Geigy Corp. (19??) Metalaxyl - Field Grapes Residue Analysis. Unpublished compilation. 161 p. (CDL:261148A)
- 00157480 Ciba-Geigy Corp. (1984) Residue Chemistry: Methods and Summary of Results: Metalaxyl. Unpublished compilation. 280 p. (CDL:262112B)
- 00164649 Marco, G.; Gold, B. (1986) Summary of Section D: Metalaxyl - Blueberries Residue Data from Field Trials in New York. Unpublished compilation. 103 p. (CDL:265762XA)
- 00164650 Nixon, W. (1985) Summary of Section D: Metalaxyl - Stone Fruits Residue Data. Unpublished compilation. 129 p. (CDL:265764A)
- 00164651 Nixon, W. (1985) Summary of Section D: Metalaxyl - Walnuts and Almonds Residue Data. Unpublished compilation. 111 p. (CDL:265765A)
- 40106601 Gold, B. (1987) Total Metalaxyl Residues in Sugar Beet Fractions following Applications of Ridomil 2E and Ridomil MZ58: (Magnitude of Residues): Including Residue Analytical Method and Storage Stability: Study No. ABR-87009. Unpublished compilation prepared by Ciba-Geigy Corp. 56 p.
- 40490501 Biehn, W. (1987) Metalaxyl - Magnitude of Residue on Papayas, including a Description of the Analytical Method Used: IR-4 PR #2532. Unpublished study prepared by University of Hawaii. 104 p.
- 40503101 Williams, B.; Condra, C. (1987) Multiresidue Method Testing of Metalaxyl and Metabolites in Crops and Animal Tissues: Final Report No. 36320. Unpublished study prepared by Analytical Bio-Chemistry Laboratories. 188 p.
- 40569301 Gold, B.; Cheung, M. (1987) Metalaxyl - Sugar Beets: [Residue Chemistry Data]: Project ID: ABR-87077. Unpublished study prepared by Ciba-Geigy Corp., in cooperation with Enviro-Bio-Tech, Ltd. 130 p.
- 40746901 Kuehne, R. (1988) Additional Analyses of Metalaxyl on German-treated Hops: Laboratory Study No.: 2366/67/87. Unpublished study prepared by Ciba-Geigy Ltd. 10 p.

- 40775801 Cheung, M. (1988) Metalaxyl (Ridomil 5G) - Head Lettuce: Laboratory Study No. ABR-88079. Unpublished study prepared by Ciba-Geigy Corp. 86 p.
- 40838301 Cheung, M. (1988) Metalaxyl (Ridomil 2E and Ridomil MZ58)--Root and Tuber Vegetable Crop Grouping: Laboratory Project ID: ABR-88112. Unpublished study prepared by Ciba-Geigy Corp. 265 p.
- 40838302 Singh, H. (1987) Standard Operating Procedure for Determination of Dithiocarbamate Residues by Carbon Disulfide Evolution Method: Laboratory Project ID: SOP EBT-201.00. Unpublished study prepared by Enviro-Bio-Tech, Ltd. 9 p.
- 40838303 Eckert, J. (1988) Standard Operating Procedure for Determination of Ethylene Thiourea (ETU) in Crops and Feed: Laboratory Project ID: SOP EBT-200.01. Unpublished study prepared by Enviro-Bio-Tech, Ltd. 14 p.
- 40832901 Cheung, M. (1988) Metalaxyl Residues in Alfalfa Resulting from the Application of Ridomil 2E to Alfalfa at Seeding: Laboratory Project ID: ABR-88114. Unpublished study prepared by Ciba-Geigy Corp. 84 p.
- 40880401 Cheung, M. (1988) Response to EPA Review of Metalaxyl (Ridomil 2E) on Strawberries Including Results of Additional Field Trials: Laboratory Project ID ABR-88131. Unpublished study prepared by Ciba-Geigy Corp. 152 p.
- 40909401 Cheung, M. (1988) Summary of Residue in Hops Following Applications of Ridomil 2E and Metalaxyl 48.8 WP: Lab. Proj. ID ABR-88152. Unpublished study prepared by Ciba-Geigy Corp. 47 p.
- 41055203 Hubbard, H. (1989) Determination of the Metalaxyl Metabolites CGA-100255 and CGA-94689 (A&B Isomers) by U.S. Food and Drug Administration Multiresidue Procedures: Proj. ID ABR-88156. Unpublished study prepared by Ciba-Geigy Corp. 78 p.
- 41150101 Cheung, M. (1989) Metalaxyl Residues in Grapes and Grape Fractions Resulting from Applications of Ridomil MZ58: Project ID ABR-89016. Unpublished study prepared by Ciba-Geigy Corp. 225 p.
- 41250101 Cheung, M. (1989) Metalaxyl: Response to EPA Review of Metalaxyl on Sugar Beets: Project ID ABR-89075. Unpublished study prepared by Ciba-Geigy Corp. 97 p.
- 41449001 Ross, J. (1990) Metalaxyl Sample Storage Interval Summary: Lab Project Nos. ABR-90016; 409925. Unpublished study prepared by Ciba-Geigy Corp., Residue Chemistry Dept. 265 p.
- 41587801 Cheung, M. (1990) Metalaxyl: Leafy Vegetables Crop Grouping (Excluding Spinach) Residue Summary: Lab Project Number: ABR-89113; 409187. Unpublished study prepared by Ciba-Geigy Corp. 574 p.
- 41636201 Cheung, M. (1990) Metalaxyl: Response to EPA Review of Metalaxyl (Ridomil 5G) on Spinach Including Results of Additional Field Trials: Lab Project Number: ABR-90042. Unpublished study prepared by Ciba-Geigy Corp. 127 p.
- 41664503 Emrani, J. (1990) Metalaxyl: Metabolism of [carbon 14]-Metalaxyl in Goats: Lab Project Number: ABR/90078. Unpublished study prepared by CIBA-GEIGY Corp., and Others. 205 p.
- 41664504 Kennedy, E. (1990) Metalaxyl: Metabolism of [phi-carbon 14]-Metalaxyl in Hens: Lab Project Number: ABR/90077. Unpublished study prepared by CIBA-GEIGY Corp., and Others. 129 p.

- 41664505 Ross, J. (1990) Preparation of Diazomethane: Lab Project Number: AG/345. Unpublished study prepared by CIBA-GEIGY Corp. 9 p.
- 41664506 McDonald, J. (1990) Procedure for Incubating Animal Tissue Samples with the Enzyme Collagenase: Lab Project Number: SOP/NUMBER/71-22. Unpublished study prepared by CIBA-GEIGY Corp. 7 p.
- 41688900 Interregional Research Project No. 4 (1990) Submission of Data in Support of Proposed Tolerance for Metalaxyl: Residue Chemistry.
- 41688901 Eudy, L. (1990) Metalaxyl-Ginseng: Residue Summary: Lab Project Number: ABR-90033. Unpublished study prepared by Ciba-Geigy Corp. 178 p.
- 41689701 Ross, J. (1990) Metalaxyl-Magnitude of Residues in Field Corn Forage Silage-stage Forage, Fodder, and Grain Following Seed Treatment with Aproff 25W: Lab Project Number: ABR 90075. Unpublished study prepared by Residue Chemical Department, Agriculture Div. Ciba-Geigy Corp. 187 p.
- 41689702 Ross, J. (1990) Metalaxyl Magnitude of Residues in Field Corn Forage Silage-stage forage, Fodder, and Grain following Seed Treatment with Apron 25W: Lab Project No. ABR-90075. Unpublished study prepared by Residue Chemistry Department, Agricultural Division, Ciba-Geigy Corp. 107 p.
- 41870301 Ross, J. (1990) Amendment 1 to ABR-90075: Metalaxyl Magnitude of the Residues in Field Corn Forage, Silage-stage Forage, Fodder, and Grain Following Seed Treatment with Apron 25W. (Related to MRIDs 41689701 and 41689702): Lab Project Number: ABR-90075. Unpublished study prepared by Ciba-Geigy Corp. 222 p.
- 41870302 Eudy, L. (1991) Metalaxyl: Magnitude of the Residues in Processed Food/Feed Commodities of Grain Sorghum Following Seed Treatment with Apron 25W: Lab Project Number: ABR-90103. Unpublished study prepared by Ciba-Geigy Corp. 147 p.
- 41870303 Smith, J. (1991) Metalaxyl: Magnitude of the Residues in Processed Food/Feed Commodities of Rice Following Seed Treatment with Apron 25W plus Preemergence Application of Ridomil 2E: Lab Project Number: ABR-90102. Unpublished study prepared by Ciba-Geigy Corp. 135 p.
- 41870304 Smith, J. (1991) Metalaxyl: Magnitude of the Residues in Processed Food/Feed Commodities of Sunflowers Following Seed Treatment with Apron 25W: Lab Project Number: ABR-90105. Unpublished study prepared by Ciba-Geigy Corp. 138 p.
- 41870305 Smith, J. (1991) Metalaxyl: Magnitude of the Residues in Processed Food/Feed Commodities of Cottonseed Following Seed Treatment with Apron 25W plus In-Furrow Application of Ridomil 5G: Lab Project Number: ABR-90104. Unpublished study prepared by Ciba-Geigy Corp. 120 p.
- 41870306 Eudy, L. (1991) Metalaxyl: Magnitude of the Residues in Peanut Vines, Hay, Nutmeats and Shells Following In-Furrow plus Banded Applications of Ridomil 5G: Lab Project Number: ABR-90107. Unpublished study prepared by Ciba-Geigy Corp. 172 p.
- 41870307 Eudy, L. (1991) Metalaxyl: Magnitude of the Residues in Processed Food/Feed Commodities of Potatoes Following Application of Ridomil 2E plus Ridomil MZ58: Lab Project Number: ABR-90106. Unpublished study prepared by Ciba-Geigy Corp. 119 p.
- 41870308 Dickson, G. (1983) Metalaxyl: Rotational Crop Studies: Lab Project Number: EIR-82011. Unpublished study prepared by Ciba-Geigy Corp. 491 p.

- 41912902 Ross, J. (1990) Metalaxyl: Sample Storage Interval Summary Amendment 1: Lab Project Number: ABR-90016. Unpublished study prepared by Ciba-Geigy Corporation. 29 p.
- 41912903 Ross, J. (1991) Metalaxyl: Response to EPA Review of Metalaxyl on Corn: Lab Project Number: ABR-91035. Unpublished study prepared by Ciba-Geigy Corporation. 13 p.
- 41996401 Smith, J. (1991) Metalaxyl - Cranberries/Residue Summary: Lab Project Number: ABR-90046. Unpublished study prepared by Ciba-Geigy Corp., Ag. Div. 161 p.
- 42021101 Cheung, M. (1991) Metalaxyl: Response to EPA Review of Metalaxyl on Leafy Vegetables Crop Grouping (Excluding Spinach): Lab Project Number: ABR-91058. Unpublished study prepared by Ciba-Geigy Corp. 20 p.
- 42115801 Ermani, J. (1990) Amendment to: Metabolism of [phi-carbon 14] Metalaxyl in Goats. Unpublished study prepared by Ciba-Geigy Corp. 4 p.
- 42115802 Emrani, J. (1990) Supplemental to: Metabolism of [phi-carbon 14] Metalaxyl in Goats. Unpublished study prepared by Ciba-Geigy Corp. 74 p.
- 42115803 Kennedy, E. (1990) Amendment to Metabolism of [phi-carbon 14] Metalaxyl in Hens. Unpublished study prepared by Ciba-Geigy Corp. 4 p.
- 42115804 Kennedy, E. (1990) Supplemental to Metabolism of [phi-carbon 14] Metalaxyl in Hens. Unpublished study prepared by Ciba-Geigy Corp. 176 p.
- 42115805 Kim Ha, J.; Lindsay, R. (1990) Method for the quantitative analysis of volatile free and total branched-chain fatty acids in cheese and milk fat. *J. of Dairy Sci.* 73(8): 1988-1999.
- 42115806 Massart-Leen, A.; De Pooter, H.; DeCloedt, M.; et al. (1981) Composition and variability of the branched-chain fatty acid fraction in the milk of goats and cows. *Lipids* 16(5): 286-292.
- 42115807 Eudy, L. (1991) Improved Analytical Method for the Determination of Total Terminal Residues of Metalaxyl in Poultry Tissues and Eggs as 2,6-Dimethyl Aniline: Lab Project Number: AG-576. Unpublished study prepared by Ciba-Geigy Corp. 45 p.
- 42115808 Yokley, R. (1991) Validation of Analytical Method AG-576 for the Determination of Total Residues of Metalaxyl in Goat Tissues, Milk, Poultry Tissues and Eggs: Lab Project Number: ABR-91008. Unpublished study prepared by Ciba-Geigy Corp. 125 p.
- 42115809 Eudy, L. (1991) Metalaxyl: Three Level/28-Day Poultry Study: Lab Project Number: ABR-91047. Unpublished study prepared by Ciba-Geigy Corp. 23 p.
- 42115810 Cheung, M. (1982) Metalaxyl: Gas Chromatographic Determination of Metalaxyl (Apron) Coatings on Seed: Lab Project Number: AG-397. Unpublished study prepared by Ciba-Geigy Corp. 11 p.
- 42159801 Smith, J. (1991) Metalaxyl: Magnitude of Residue in Brassica Vegetables, Cole Crop Grouping Following a Preplant, Broadcast, Incorporated Application of Ridomil 2E and Four Post Foliar Spray Applications of Ridomil Bravo 81W: Lab Project Number: ABR-91034. Unpublished study prepared by Ciba-Geigy Corp. 127 p.

42196501 McFarland, J. (1992) Uptake and Metabolism of Metalaxyl in Greenhouse Rotational Crops Following Target Tobacco Grown in Soil Treated with [Phenyl-carbon 14]-Metalaxyl: Lab Project Number: ABR-91084. Unpublished study prepared by Ciba-Geigy Corp. 272 p.

42196502 Madrid, S. (1981) Uptake, Balance and Metabolism of [carbon 14]-CGA-48988 in Field Grown Potatoes: Lab Project Number: ABR-81037. Unpublished study prepared by Ciba-Geigy Corp. 32 p.

42196503 Honeycutt, R. (1980) Identification of Four Sugar Conjugate Metabolites of [carbon 14]-CGA-48988 in Tobacco: Lab Project Number: ABR-80035. Unpublished study prepared by Ciba-Geigy Corp. 25 p.

42233501 Eudy, L. (1992) Metalaxyl: Magnitude of the Residue in Processed Food/Feed Commodities of Pineapples Following Seed Piece Dip with Ridomil 2E: Lab Project Number: ABR-91086. Unpublished study prepared by Ciba-Geigy Corp. 110 p.

42259805 Eudy, L. (1992) Metalaxyl: Magnitude of Residues in Water-Seeded Rice Following Application of Ridomil 2E: Lab Project Number: ABR-91018. Unpublished study prepared by Ciba-Geigy Corp. 200 p.

42498700 Ciba-Geigy Corp. (1992) Submittal of residue data and response to EPA review in support of Metalaxyl registration standard. Transmittal of 2 studies.

42498701 Ross, J. (1992) Response to EPA Review of Metalaxyl on Cereal Grains, Soybeans, Potatoes, Peanuts, Cottonseed, Sunflower Seed, and Pineapples: Lab Project Number: ABR-92054. Unpublished study prepared by Ciba-Geigy Corp. 81 p.

42919401 Eudy, L. (1993) Storage Stability of Total Residues of Metalaxyl in Weathered Crops Under Freezer Storage Conditions: Lab Project Number: ABR-93009. Unpublished study prepared by Ciba-Geigy Corporation. 127 p.