US ERA ARCHIVE DOCUMENT

Data Evaluation Report on the acute toxicity of AE C421200 (Metabolite of Thidiazuron) to Rainbow Trout (Oncorhynchus mykiss)

PMRA Submission Number {......}

EPA MRID Number 46203511

Data Requirement:

PMRA DATA CODE

EPA DP Barcode OECD Data Point

EPA MRID **EPA** Guideline D294536

46203511 §72-1c

Test material:

AE C421200

Metabolite of thidiazuron

Common name: Chemical name:

IUPAC: 1-Cyano-3-phenylurea

CAS name: Not reported CAS No.: Not reported Synonyms: None reported **Purity:** 98.4% (w:w)

Primary Reviewer: Greg Hess

Staff Scientist, Dynamac Corporation

Signature:

Date: 4/21/04

OC Reviewer: Christie E. Padova Staff Scientist, Dynamac Corporation Signature: C.E. Pader Date: 4/26/04

Primary Reviewer: Bill Evans, Biologist

OPP/EFED/ERB - I

Date:

Secondary Reviewer(s):

{EPA/OECD/PMRA}

Date:

Reference/Submission No.:

Company Code:

Active Code:

EPA PC Code: 120301

Date Evaluation Completed:

CITATION: Palmer, S.J., et al. 2003. AE C421200: A 96-Hour Static Acute Toxicity Test with the Rainbow Torut (Oncorhynchus mykiss). Unpublished study performed by Wildlife International, Ltd., Easton, MD. Laboratory Project No. 149A-159. Study sponsored by Bayer CropScience, Frankfurt am Main, Germany. Study initiated April 8, 2003 and completed June 30, 2003.

EXECUTIVE SUMMARY:

In a 96-hour acute toxicity study, Rainbow Trout (*Oncorhynchus mykiss*) were exposed under static conditions to AE C421200 (a metabolite of thidiazuron) at nominal concentrations of 0 (negative control), 6.3, 13, 25, 50, or 100 ppm. Mean-measured concentrations were <3.00 (<LOQ, control), 6.3, 13, 25, 51, and 103 ppm a.i.

No mortality or sub-lethal effects were observed in any test group following 96 hours of exposure. The 96-hour LC_{50} was >103 ppm a.i., which categorizes AE C421200 (a metabolite of thidiazuron) as practically non-toxic to Rainbow trout (*Oncorhynchus mykiss*) on an acute toxicity basis. The NOEC and LOEC, based on both mortality and sub-lethal effects, were 103 and >103 ppm a.i., respectively.

This study is scientifically sound and satisfies the guideline requirements for an acute toxicity study with Rainbow Trout (§72-1c). This study is classified as CORE.

Results Synopsis

Test Organism Size/Age (mean Weight or Length):

11 weeks old (reviewer-calculated); mean of 1.1 g (wet) and 5.3 cm (mean of 10 negative control fish at test

termination)

Test Type (Flow-through, Static, Static Renewal):

Static

96-Hour

LC₅₀: >103 ppm a.i. NOEC: 103 ppm a.i. LOEC: >103 ppm a.i. Endpoints affected: None

I. MATERIALS AND METHODS

GUIDELINE FOLLOWED:

The study protocol was based on procedures outlined in the OECD Guideline No. 203 (1993); the U.S. EPA OPPTS No. 850.1075 (Draft, 1996); and ASTM Standard E729-88a (1994). Deviations from U.S. EPA §72-1 included:

- Aeration of the test vessels was not addressed.
- The water hardness (132 mg/L as CaCO₃) was three times higher than recommended (40-48 mg/L as CaCO₃).
- 3. The pH range (8.1-8.6) was greater than recommended (7.2-7.6).
- 4. The total organic carbon (TOC), particulate matter, and residual chlorine concentrations in the dilution water were not reported.

These deviations did not affect the validity or acceptability of the study.

COMPLIANCE:

Signed and dated GLP, Confidentiality, and Quality Assurance statements were provided. This study was conducted in accordance with GLP standards of the U.S. EPA (40 CFR Part 160), OECD, and Japan MAFF (p. 3).

A. MATERIALS:

1. Test Material

AE C421200 (a metabolite of thidiazuron)

Description:

White powder

Lot No./Batch No.:

DSC 1441 (Product code: AE C421200 00 1B98 0001)

Purity:

98.4% (w:w) a.i.

Stability of Compound

Under Test Conditions:

The stability of the test substance in the dilution water during the course of the study was verified by analytical determination at 0, 48, and 96 hours (Table 1, p. 17). Recoveries were 97.8-103% of nominal concentrations at 0 hours, 97.3-103% of nominal at 48 hours, and 97.5-

104% of nominal at 96 hours.

Storage conditions of

test chemicals:

Stored frozen.

OECD requires water solubility, stability in water and light, pK_{a} , P_{ow} , and vapor pressure of the test compound. OECD requirements were not reported.

2. Test organism:

Species:

Rainbow Trout (Oncorhynchus mykiss)

Age at test initiation:

11 weeks old (reviewer-calculated, hatched on February 24, 2003).

Weight at study initiation:

Not provided; the blotted wet weight of 10 negative control fish measured at test termination averaged 1.1 g (range of 0.87-1.8 g).

Length at study initiation:

Not provided; the length of 10 negative control fish measured at test

termination averaged 5.3 cm (range of 4.8-6.1 cm).

Source:

Thomas Fish Company, Anderson, CA.

B. STUDY DESIGN:

1. Experimental Conditions

a. Range-finding Study: The definitive nominal test concentrations were selected in consultation with the sponsor, and were based upon the results of an exploratory range-finding toxicity test. The results of the range-finding study were not reported (p. 9).

b. Definitive Study

Table 1. Experimental Parameters

Parameter	Details	Remarks		
		Criteria		
Acclimation period:	At least 14 days prior to testing			
Conditions: (same as test or not)	Same as test			
Feeding:	Fed commercially-prepared diet supplied by Zeigler Brothers Inc., Gardners, PA. Fish were not fed two days prior to and during testing.	EPA requires: minimum 14 days; no feeding during test OECD requires minimum of 12 days.		
Health: (any mortality observed)	During the 48 hours prior to testing, fish showed no signs of disease, stress or mortality.			
Duration of the test	96-hour			
		EPA/OECD requires: 96 hour		
Test condition				
static/flow through	Static			
Type of dilution system- for flow through method.	N/A	EPA: Must provide reproducible supply of toxicant, with a consistent		
Renewal rate for static renewal	N/A	flow rate of 5-10 vol/24 hours, and meter systems calibrated before study and checked twice daily during test period		
Aeration, if any	Not reported			
a a		EPA requires: no aeration; OECD permits aeration		
Test vessel				
Material: (glass/stainless steel)	Stainless steal aquaria with stainless steal cover			
Size: Fill volume:	38 L 25 L (19.5-cm depth)	EPA requires: Size 19 L (5 gal) or 30 x 60 x 30 cm Fill volume: 15-30 L of solution		
Source of dilution water	The dilution water was freshwater obtained from an on-site laboratory well (40-m deep). The well water was sand-filtered, aerated, then filtered (0.45 µm) again prior to use.	EPA 1975; Soft reconstituted water or water from a natural source, not dechlorinated tap water; OECD permits dechlorinated tap water.		

Data Evaluation Report on the acute toxicity of AE C421200 (Metabolite of Thidiazuron) to Rainbow Trout (Oncorhynchus mykiss) PMRA Submission Number {........}

EPA MRID Number 46203511

Parameter	Details	Remarks
12		Criteria
Water parameters: Hardness pH Dissolved oxygen Total Organic Carbon Particulate Matter	132 mg CaCO ₃ /L 8.1-8.6 7.4-9.4 mg/L (≥68% saturation) Not reported Not reported	The hardness and pH were higher than recommended. Total alkalinity was 190 mg/L as CaCO ₃ . Results of the analysis of the well water on July 31, 2002 for pesticides, organics, and metals are provided in Appendix 3, pp. 26-27.
Metals Pesticides	See Appendix 3, p. 27. <lod< td=""><td>provided in Appendix 3, pp. 20-27.</td></lod<>	provided in Appendix 3, pp. 20-27.
Chlorine Temperature Intervals of water quality measurement	Not reported 11.5 to 13.3°C The DO, pH and temperature were measured in both replicate aquaria at 0-, 24-, 48-, 72- and 96-hours. Temperature was also measured continuously in one negative control aquaria.	Hardness and pH EPA requires hardness of 40-48 mg/L as CaCO₃ and pH of 7.2-7.6. OECD allows hardness of 10-250 mg/L as CaCO₃ and pH between 6 and 8.5. Dissolved Oxygen Renewal: ≥60% during 1 st 48 hrs and ≥ 40% during 2 nd 48 hrs Flow-through: ≥60% through out test. OECD requires at least 80% saturation value. Temperature EPA requires 12°C for coldwater species and 17-22°C for warmwater species. OECD requires range of 21- 25°C for bluegill and 13-17°C for rainbow trout. EPA water quality measured at beginning of test and every 48 hours

Parameter	Details	Remarks
		Criteria
Concentration of test material: nominal:	0 (negative control), 6.3, 13, 25, 50, and 100 ppm	Mean-measured concentrations are provided in Table 1, p. 17.
measured:	<3.00 (<loq, 103="" 13,="" 25,="" 51,="" 6.3,="" a.i.<="" and="" control),="" ppm="" td=""><td>Concentrations were stable during the 96-hour study.</td></loq,>	Concentrations were stable during the 96-hour study.
		Stock solutions were adjusted for purity of the test material (p. 11).
		EPA/OECD requires: Control and five treatment levels. Each conc. should be 60% of the next highest conc., and should be in a geometric series
Solvent (type, percentage, if used)	N/A	
	a e	EPA requires: Not to exceed 0.5 mL/L for static tests or 0.1 mL/L for flow-through tests; OECD requires solvent, exceed 100 mg/L.
Number of fish/replicates: negative control:	20 fish, divided into two replicates containing 10 fish each	# I
solvent control:	N/A	EPA: ≥ 10/concentration;
treated:	20 fish, divided into two replicates containing 10 fish each	OECD requires at least 7 fish/concentration
Biomass loading rate	0.46 g fish/L	
C.		Static: ≤ 0.8 g/L at ≤ 17°C, ≤ 0.5 g/L at > 17°C; flow-through: ≤ 1 g/L/day; OECD requires maximum of 1 g fish/L for static and semi-static with higher rates accepted for flow-through
Lighting	16-hours light/8-hours dark, with a 30-minute transition period.	Light intensity of 272 lux at the water surface during daylight hours.
		EPA requires: 16 hours light/8 hours dark); OECD requires 12 -16 hours photoperiod.

PMRA Submission Number {......}

EPA MRID Number 46203511

Parameter	Details	Remarks		
		Criteria		
Feeding	Animals were not fed during testing.	EPA/OECD requires: No feeding during the study		
Recovery of chemical Level of Quantitation Level of Detection	99.5-102% of nominal 3.00 ppm a.i. Not reported	Based on quality control matrix spikes fortified at 6.00, 20.0, or 100 ppm and analyzed concurrently with the samples (Appendix 4.5, p. 33).		
Positive control {if used, indicate the chemical and concentrations}	N/A			
Other parameters, if any	· N/A			

2. Observations:

Table 2: Observations

Criteria	Details	Remarks/Criteria	
Parameters measured including the sub-lethal effects/toxicity symptoms	Mortality and sub-lethal effects		
Observation intervals	at 5 hours and every 24 hours thereafter	(EPA/OECD requires: minimally every 24 hours)	
Were raw data included?	Yes, sufficient		
Other observations, if any	N/A		

II. RESULTS AND DISCUSSION:

A. MORTALITY:

No mortality occurred in any group during the 96-hour study (Table 4, p. 20). The 96-hour LC_{50} was >103 ppm a.i., the highest concentration tested (Table 5, p. 21). The NOEC based on mortality data was 103 ppm a.i.

PMRA Submission Number {......}

EPA MRID Number 46203511

Table 3: Effect of AE C421200 on mortality of Rainbow Trout (Oncorhynchus mykiss).

Treatment, ppm a.i. Measured and (nominal) concn.	No of	Observation Period					
	No. of fish at start of study	0-48 Hours		72 Hours		96 Hours	
		No Dead	% mortality	No Dead	% mortality	No Dead	% mortality
Negative control	20	0	0	0	0	0	0
6.3 (6.3)	20	0	0	0	0	0	0
13 (13)	20	0	0	0	0	0	0
25 (25)	20	0	0	0	0	0	0
51 (50)	20	0	0	0	0	0	0
103 (100)	20	0	0	0	0	0	0
NOEC (mortality)	103 ppm a.i.						
LC ₅₀ (95% C.I.)	>103 ppm a.i.						
Positive control, if used mortality: LC ₅₀ :	N/A	N/A	N/A	N/A	N/A	N/A	N/A

B. NON-LETHAL TOXICITY ENDPOINTS:

No sub-lethal effects were observed in any group during the 96-hour study (Table 4, p. 20). The NOEC based on sub-lethal effects was 103 ppm a.i.

PMRA Submission Number {......

EPA MRID Number 46203511

Table 4. Sub-lethal effects of AE C421200 on Rainbow Trout (Oncorhynchus mykiss).

Treatment, ppm a.i. Measured and (nominal) concn.	Observation Period				
	endpoint at 5-24 Hours	endpoint at 48 Hours	endpoint at 72 Hours	endpoint at 96 Hours	
	% affected	% affected	% affected	% affected	
Negative control	AN	AN	AN	AN	
6.3 (6.3)	AN	AN	AN	AN	
13 (13)	AN	AN	AN	AN	
25 (25)	AN	AN	AN	AN	
51 (50)	AN	AN	AN	AN	
103 (100)	AN	AN	AN	AN	
NOEC (sub-lethal)	103 ppm a.i.				
LOEC (sub-lethal)	>103 ppm a.i.				
EC ₅₀	Not determined				
Positive control, if used % sub-lethal effect: EC ₅₀ :	N/A	N/A	N/A	N/A	

AN - All surviving fish appeared normal.

C. REPORTED STATISTICS:

Due to a lack of mortality or sub-lethal effects at any treatment level by 96 hours, the 96-hour LC₅₀, NOEC, and LOEC values were determined by visual observation, and were based on mean-measured treatment concentrations (p. 14).

96-Hour

LC₅₀: >103 ppm a.i. NOEC: 103 ppm a.i. LOEC: >103 ppm a.i. Endpoints affected: None

D. VERIFICATION OF STATISTICAL RESULTS:

The 96-hour LC₅₀ was determined visually due to a lack of mortality at any treatment level by 96 hours. The NOEC was visually determined as the highest concentration which exhibited no significant mortality or sublethal effects. All toxicity values were determined in terms of the reported mean-measured treatment concentrations.

Data Evaluation Report on the acute toxicity of AE C421200 (Metabolite of Thidiazuron) to Rainbow Trout (Oncorhynchus mykiss)

PMRA Submission Number {......}

EPA MRID Number 46203511

96-Hour

LC₅₀: >103 ppm a.i. NOEC: 103 ppm a.i. LOEC: >103 ppm a.i. Endpoints affected: None

E. STUDY DEFICIENCIES:

There were no significant deviations from U.S. EPA guideline §72-1c that affected the acceptability or validity of this study.

F. REVIEWER'S COMMENTS:

Results of the reviewer's statistical verification were identical to those of the study author.

The test solutions appeared clear and colorless at test initiation and termination (p. 12).

G. CONCLUSIONS:

This study is scientifically sound and satisfies the guideline requirements for an acute toxicity study with freshwater fish, cold water species (§72-1c). This study is classified as CORE. The 96-hour LC_{50} was >103 ppm a.i., which classifies AE C421200 (a metabolite of thidiazuron) as practically non-toxic to Rainbow Trout (*Oncorhynchus mykiss*) on an acute toxicity basis. The NOEC (for mortality and sub-lethal effects) was 103 ppm a.i., the highest concentration tested.

96-Hour

LC₅₀: >103 ppm a.i. NOEC: 103 ppm a.i. LOEC: >103 ppm a.i. Endpoints affected: None

III. REFERENCES:

- Organization for Economic Co-Operation and Development (OECD). 1993. Guideline for the Testing of Chemicals. *Guideline 203: Fish Acute Toxicity Test*, Adopted by Council on 12 July 1992.
- U.S. Environmental Protection Agency. 1996. Fish Acute Toxicity Test, Freshwater and Marine. Series 850 Ecological Effects Test Guidelines (draft), OPPTS Number 850.1075.
- ASTM Standard E729-88a. 1994. Standard Guide for Conducting Acute Toxicity Tests with Fishes, Macroinvertebrates, and Amphibians. American Society for Testing and Materials.
- APHA, AWWA, WPCF. 1998. Standard Methods for the Examination of Water and Wastewater. 20th Edition, American Public Health Association. American Water Works Association. Water Pollution Control Federation, New York.