

US EPA ARCHIVE DOCUMENT

BB-316  
TAR-4621



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

004621

8/20/85

OFFICE OF  
PESTICIDES AND TOXIC SUBSTANCES

MEMORANDUM

SUBJECT: EPA No. 35977-U. Fenoxycarb. Review of 21-Day Dermal Toxicity  
Study in Rats

Tox. Chem. No. 652C

FROM: Edwin R. Budd, Section Head  
Section II, Toxicology Branch  
Hazard Evaluation Division (TS-769c)

*Budd*  
*8/20/85*  
*WMB*  
*8/20/85*

TO: Tim Gardner, PM# 17  
Registration Division (TS-767c)

CONCLUSIONS

The subject 21-day dermal toxicity study (reviewed below) is acceptable in fulfillment of the requirement for same.

Toxicology Branch has no objection to the full registration of Logic® 1% Fire Ant Bait for use on non-agricultural turf (including residential sites). All toxicology data requirements have been satisfied. The label precautionary statements are acceptable.

1076

21-DAY DERMAL TOXICITY STUDY OF FENOXYCARB (Ro 13-5223/000) IN RATS

004621

Hazleton Laboratories Europe, Ltd; Report No. 4552-161/157; July 3, 1985

PROTOCOL: Groups of 5 male and 5 female Crl:CD(SD)BR rats (241-362 g) were dosed with 4 ml/kg/day of freshly prepared Ro 13-5223/000 (96.6% pure) in corn oil at dose levels of 0 (vehicle control), 20, 200, and 2000 mg/kg/day for 21 days. They were dosed on shaved nonabraded skin which represented at least 10% of the body surface area. Each dose was occluded for 6 hours (there was no mention of dose removal). Each rat was examined once daily for clinical signs and scored for skin irritation. Body weights and food consumption were measured at weekly intervals. Blood samples drawn from the orbital sinus of each fasted animal during the final study week were evaluated for the following parameters:

Hematology:

Erythrocyte count	Mean cell hemoglobin concentration
Hemoglobin concentration	Leukocyte count
Packed cell volume	Differential leukocyte count
Mean cell volume	Platelet count
Mean cell hemoglobin	

Clinical Chemistry:

SGOT	Creatinine
SGPT	Albumin
Alkaline phosphatase	Albumin/globulin ratio
Bilirubin	Sodium
Total protein	Potassium
BUN	Calcium
Glucose	Phosphorus (inorganic)

All rats were sacrificed on day 21 and examined grossly. Organ weights were measured for livers, kidneys, adrenals, ovaries, and testes. Sections of liver, kidney, treated skin, untreated skin, and any gross lesions were examined microscopically for the vehicle control and high-dose rats. Food and water were available *ad libitum* except prior to bleeding and necropsy. Dose concentration analysis was performed on days 1 and 15.

RESULTS: Dose concentration analysis demonstrated that samples were within 89.5-93.5% of nominal on day 1, and within 93.5-95.5% of nominal on day 15; this is clearly within an acceptable range. There were no compound-related effects on body weights, food consumption, or clinical pathology. Absolute and relative liver weights were moderately elevated (22% and 19%, respectively) in both sexes dosed at 2000 mg/kg/day, compared to the control group. Slight erythema was observed in 3 low-dose males and 1 mid-dose female during the first week of dosing; these lesions were probably not due to the test article. No other clinical signs were reported in any rats. Mild to severe abdominal alopecia in all rats, and occasional concurrent mild to marked sores were observed during gross necropsy and were attributed to the method of occlusion. Mild to severe hydronephrosis of the right kidney was also seen in a few of the dosed animals, but was probably not dose-related. Microscopic evaluations of skin did not reveal any dose-related effects. Two males and 3 females in the high-dose group 4 had slight liver hypertrophy.

**BEST AVAILABLE COPY**

2

004621

CONCLUSIONS: There were no deaths during this study, so the dermal LD<sub>50</sub> is >2000 mg/kg/day. The NOEL is 200 mg/kg/day, and the LEL is 2000 mg/kg/day [slight liver hypertrophy]. Skin lesions including slight erythema and alopecia and sores on abdominal skin (dosing site?) were probably caused by the method of occlusion. The only dose-related toxicity was slight liver hypertrophy in the 2000 mg/kg/day dose group which resulted in elevated liver weights.

This study is CORE GUIDELINE. The report failed to describe the dosing sites used, the method of occlusion, and the means of removing the doses. The age of the rats was not supplied. During gross necropsy, all rats were reported as having mild to severe abdominal alopecia, and some had concurrent mild to marked sores. These lesions were certainly severe enough that they should have been recorded as clinical signs, yet they were not - thus making it impossible to determine the time of onset. This study was reviewed for Quality Assurance.

Reviewed by - John E. Whalan, Toxicologist  
Section II, Toxicology Branch

BEST AVAILABLE COPY

Fenoxycarb toxicology review

---

Page \_\_\_\_\_ is not included in this copy.

Pages 4 through 6 are not included in this copy.

---

The material not included contains the following type of information:

- Identity of product inert ingredients
  - Identity of product impurities
  - Description of the product manufacturing process
  - Description of product quality control procedures
  - Identity of the source of product ingredients
  - Sales or other commercial/financial information
  - A draft product label
  - The product confidential statement of formula
  - Information about a pending registration action
  - FIFRA registration data
  - The document is a duplicate of page(s) \_\_\_\_\_
  - The document is not responsive to the request
- 

• The information not included is generally considered confidential by product registrants. If you have any questions, please contact the individual who prepared the response to your request.

---