

US EPA ARCHIVE DOCUMENT

Page 2



UNITED STATES ENVIRONMENTAL PROTECTION
AGENCY
WASHINGTON D.C., 20460

OPP OFFICIAL RECORD
HEALTH EFFECTS DIVISION
SCIENTIFIC DATA REVIEWS
EPA SERIES 361

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

TXR No. 0050919

MEMORANDUM

August 26, 2002

SUBJECT: BAS510F Qualitative Risk Assessment Based On Wistar Chbb:THOM
(SPF) Rat Dietary Studies

P.C. Code 128008

TO: Alan Levy, Toxicologist
Registration Action Branch 2
Health Effects Division (7509C)

FROM: Lori L. Brunsman, Statistician
Science Information Management Branch
Health Effects Division (7509C)

Health Effects 8/26/02

THROUGH: Jess Rowland, Branch Chief
Science Information Management Branch
Health Effects Division (7509C)

Jess Rowland 8/27/02

Background

A carcinogenicity study in Wistar Chbb:THOM (SPF) rats was conducted by the Experimental Toxicology and Ecology BASF Aktiengesellschaft, Ludwigshafen/Rhein, Germany, for the BASF Corporation Agricultural Products Division, Research Triangle Park, North Carolina, and dated February 28, 2001 (Laboratory Project No. 82C0179/97090, BASF Registration Document No. 2001/1000115, MRID No. 45404828).

The study design allocated groups of 50 rats per sex to dose levels of 0, 100, 500, 2500 or 15,000 ppm (0, 4.6, 23.0, 116.1 and 768.8 mg/kg/day for males; 0, 6.0, 29.7, 155.6, and 1024.4 mg/kg/day for females) of BAS510F for 105 weeks. The 15,000 ppm dose group exhibited mortality in males and excessive weight loss in females and, therefore, this dose group was terminated at 17 months and not further analyzed.

A chronic toxicity study in Wistar Chbb:THOM (SPF) rats was conducted by the Experimental Toxicology and Ecology BASF Aktiengesellschaft, Ludwigshafen/Rhein, Germany, for the BASF Corporation Agricultural Products Division, Research Triangle Park, North Carolina, and dated February 28, 2001 (Laboratory Project No. 82C0179/97091, BASF Registration Document No. 2001/1000114, MRID No. 45404827).

The study design allocated groups of 20 rats per sex to dose levels of 0, 100, 500, 2500 or 15,000 ppm (0, 4.4, 21.9, 110.0 and 739.0 mg/kg/day for

males; 0, 5.9, 30.0, 150.3, and 1000.4 mg/kg/day for females) of BAS510F for 105 weeks. Both sexes in the 15,000 ppm dose group exhibited excessive weight loss and, therefore, these animals were terminated at 17 months and not further analyzed.

Survival Analyses

The statistical evaluation of mortality indicated no significant incremental changes with increasing doses of BAS510F in either male or female rats of the carcinogenicity study, or in male rats of the chronic toxicity study. Female rats of the chronic toxicity study indicated a statistically significant decreasing trend for mortality, as well as a statistically significant pair-wise comparison of the high dose with the control. Though there were survival disparities among the dose groups for the female rats of the chronic toxicity study, there were too few animals in this study to appropriately use the Peto's Prevalence test, since the first tumor was observed at week 102. The Fisher's Exact test and the exact test for trend were used for all mortality analyses. See Tables 1, 4 and 6 for male mortality test results. See Tables 9, 12 and 14 for female mortality test results.

The statistical evaluation of mortality was based upon the Thomas, Breslow and Gart computer program.

Tumor Analyses: Carcinogenicity Study

Male rats had significant increasing trends in thyroid follicular cell adenomas at $p < 0.01$, and adenomas and/or carcinomas combined at $p < 0.05$. There were no significant differences in the pair-wise comparisons of the dosed groups with the controls.

Female rats had a significant increasing trend in thyroid follicular cell adenomas at $p < 0.05$. There were no significant differences in the pair-wise comparisons of the dosed groups with the controls.

The statistical analyses of both sexes were based upon the Exact trend test and the Fisher's Exact test for pair-wise comparisons. See Tables 2, 3, 10 and 11 for carcinogenicity study tumor analysis results.

Tumor Analyses: Chronic Toxicity Study

There were no statistically significant trends or pair-wise comparisons of the dosed groups with the controls in either male or female rats of the chronic toxicity study.

The statistical analyses of both sexes were based upon the Exact trend test and the Fisher's Exact test for pair-wise comparisons. See Tables 5 and 13 for chronic toxicity study tumor analysis results.

Tumor Analyses: Carcinogenicity and Chronic Toxicity Studies Combined

Male rats had significant increasing trends in thyroid follicular cell adenomas at $p < 0.01$, and adenomas and/or carcinomas combined at $p < 0.05$. There was a significant difference in the pair-wise comparison of the 2500 ppm dose group with the controls for thyroid follicular cell adenomas at $p < 0.05$.

Female rats had a significant increasing trend for thyroid follicular cell adenomas at $p < 0.05$. There were no significant differences in the pair-wise comparisons of the dosed groups with the controls.

The statistical analyses of both sexes were based upon the Exact trend test and the Fisher's Exact test for pair-wise comparisons. See Tables 7, 8, 15, and 16 for carcinogenicity and chronic toxicity studies combined tumor analysis results.

4

Table 1. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity Rat Study
Male Mortality Rates* and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-26	27-52	53-78	79-108 ^f	
0	0/50	0/50	2/50	7/48	9/50 (18)
100	0/50	0/50	2/50	11/48	13/50 (26)
500	0/50	0/50	2/50	11/48	13/50 (26)
2500	0/50	0/50	2/50	14/48	16/50 (32)

*Number of animals that died during interval/Number of animals alive at the beginning of the interval.

^fFinal sacrifice at week 105.

() Percent.

Note: Time intervals were selected for display purposes only.

Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

4

Table 2. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity Rat Study

Male Thyroid Follicular Cell Tumor Rates* and Exact Trend Test
and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
Adenomas (%)	0/50 (0)	0/50 (0)	1/50 (2)	4 [*] /50 (8)
p =	0.0054 ^{**}	1.0000	0.5000	0.0587
Carcinomas (%)	1 ^b /50 (2)	0/50 (0)	0/50 (0)	0/50 (0)
p =	0.2500	0.5000	0.5000	0.5000
Combined (%)	1/50 (2)	0/50 (0)	1/50 (2)	4/50 (8)
p =	0.02145 [*]	0.5000	0.7525	0.1811

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

^{*}First adenoma observed at week 87, dose 2500 ppm.

^bFirst carcinoma observed at week 107, dose 0 ppm.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If ^{*}, then $p < 0.05$. If ^{**}, then $p < 0.01$.

Table 3. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity Rat Study

Male Thyroid C-Cell Tumor Rates^a and Exact Trend Test
and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
Adenomas (%)	5/50 (10)	5 ^a /50 (10)	6/50 (12)	3/50 (6)
p =	0.2019	0.6297	0.5000	0.3575
Carcinomas (%)	0/50 (0)	2 ^b /50 (4)	0/50 (0)	1/50 (2)
p =	0.4384	0.2475	1.0000	0.5000
Combined (%)	5/50 (10)	7/50 (14)	6/50 (12)	4/50 (8)
p =	0.2423	0.3798	0.5000	0.5000

^aNumber of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

^aFirst adenoma observed at week 84, dose 100 ppm.

^bFirst carcinoma observed at week 72, dose 100 ppm.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

7

Table 4. BAS510F - Wistar Chbb:THOM (SPF) Chronic Toxicity Rat Study

Male Mortality Rates* and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-26	27-52	53-78	79-106 ^f	
0	0/20	0/20	0/20	5/20	5/20 (25)
100	0/20	0/20	1/20	3/19	4/20 (20)
500	0/20	0/20	2/20	4/18	6/20 (30)
2500	0/20	0/20	1/20	5/19	6/20 (30)

*Number of animals that died during interval/Number of animals alive at the beginning of the interval.

^fFinal sacrifice at week 105.

() Percent.

Note: Time intervals were selected for display purposes only.

Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

7

8

Table 5. BAS510F - Wistar Chbb:THOM (SPF) Chronic Toxicity Rat Study

Male Thyroid Tumor Rates* and Exact Trend Test
and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
C-Cell Adenomas#	4/20	2/20	3 ^a /20	4/20
(%)	(20)	(10)	(15)	(20)
p =	0.3274	0.3307	0.5000	0.6526
<hr/>				
Follicular Cell Adenomas#	0/20	0/20	2 ^b /20	1/20
(%)	(0)	(0)	(10)	(5)
p =	0.1989	1.0000	0.4949	0.5000

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

^aFirst c-cell adenoma observed at week 79, dose 500 ppm.

^bFirst follicular cell adenoma observed at week 105, dose 500 ppm.

#No c-cell or follicular cell carcinomas were observed.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

8

9

Table 6. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity and Chronic Toxicity Combined Rat Studies

Male Mortality Rates* and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-26	27-52	53-78	79-108 ^f	
0	0/70	0/70	2/70	12/68	14/70 (20)
100	0/70	0/70	3/70	14/67	17/70 (24)
500	0/70	0/70	4/70	15/66	19/70 (27)
2500	0/70	0/70	3/70	19/67	22/70 (31)

*Number of animals that died during interval/Number of animals alive at the beginning of the interval.

^fFinal sacrifice at week 105.

() Percent.

Note: Time intervals were selected for display purposes only.

Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

9

10

Table 7. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity and Chronic Toxicity Combined Rat Studies

Male Thyroid Follicular Cell Tumor Rates* and Exact Trend Test and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
Adenomas (%)	0/70 (0)	0/70 (0)	3/70 (4)	5 ^a /70 (7)
p =	0.0045 ^{**}	1.0000	0.1223	0.0290 [*]
Carcinomas (%)	1 ^b /70 (1)	0/70 (0)	0/70 (0)	0/70 (0)
p =	0.2500	0.5000	0.5000	0.5000
Combined (%)	1/70 (1)	0/70 (0)	3/70 (4)	5/70 (7)
p =	0.0130 [*]	0.5000	0.3098	0.1043

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

^aFirst adenoma observed at week 87, dose 2500 ppm.

^bFirst carcinoma observed at week 107, dose 0 ppm.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If ^{*}, then $p < 0.05$. If ^{**}, then $p < 0.01$.

10

11

Table 8. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity and Chronic Toxicity Combined Rat Studies

Male Thyroid C-Cell Tumor Rates* and Exact Trend Test and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
Adenomas (%)	9/70 (13)	7/70 (10)	9 ^a /70 (13)	7/70 (10)
p =	0.3625	0.3957	0.5993	0.3957
Carcinomas (%)	0/70 (0)	2 ^b /70 (3)	0/70 (0)	1/70 (1)
p =	0.4382	0.2482	1.0000	0.5000
Combined (%)	9/70 (13)	9/70 (13)	9/70 (13)	8/70 (11)
p =	0.3927	0.5993	0.5993	0.5000

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

^aFirst adenoma observed at week 79, dose 500 ppm.

^bFirst carcinoma observed at week 72, dose 100 ppm.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If ', then $p < 0.05$. If "", then $p < 0.01$.

11

12

Table 9. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity Rat Study

Female Mortality Rates* and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-26	27-52	53-78	79-107 [†]	
0	0/50	0/50	2/50	12/48	14/50 (28)
100	0/50	0/50	2/50	6/48	8/50 (16)
500	0/50	1/50	1/49	9/48	11/50 (22)
2500	0/50	0/50	1/50	9/49	10/50 (20)

*Number of animals that died during interval/Number of animals alive at the beginning of the interval.

[†]Final sacrifice at week 105.

() Percent.

Note: Time intervals were selected for display purposes only.

Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

12

13

Table 10. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity Rat Study

Female Thyroid Follicular Cell Tumor Rates* and Exact Trend Test
and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
Adenomas* (%)	0/50 (0)	1*/50 (2)	0/49 (0)	3/50 (6)
p =	0.0342*	0.5000	1.0000	0.1212

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

*First adenoma observed at week 106, dose 100 ppm.

*No carcinomas were observed.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

13

Table 11. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity Rat Study

Female Thyroid C-Cell Tumor Rates and Exact Trend Test
and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
Adenomas (%)	8/50 (16)	8/50 (16)	7 ^a /49 (14)	5/50 (10)
p =	0.1664	0.6071	0.5171	0.2768
Carcinomas (%)	0/50 (0)	0/50 (0)	1 ^b /49 (2)	0/50 (0)
p =	0.7487	1.0000	0.4949	1.0000
Combined (%)	8/50 (16)	8/50 (16)	8/49 (16)	5/50 (10)
p =	0.1566	0.6071	0.5900	0.2768

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

^aFirst adenoma observed at week 77, dose 500 ppm.

^bFirst carcinoma observed at week 105, dose 500 ppm.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose
level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

15

Table 12. BAS510F - Wistar Chbb:THOM (SPF) Chronic Toxicity Rat Study

Female Mortality Rates^a and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-26	27-52	53-78	79-106 ^b	
0	0/20	0/20	1/20	6/19	7/20 (35)*
100	0/20	0/20	0/20	3/20	3/20 (15)
500	0/20	0/20	1/20	6/19	7/20 (35)
2500	0/20	0/20	1/20	0/19	1/20 (5)*

^aNumber of animals that died during interval/Number of animals alive at the beginning of the interval.

^bFinal sacrifice at week 105.

() Percent.

Note: Time intervals were selected for display purposes only.

Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

16

Table 13. BAS510F - Wistar Chbb:THOM (SPF) Chronic Toxicity Rat Study

Female Thyroid Tumor Rates* and Exact Trend Test
and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
C-Cell Adenomas#	3/20	5 ^a /20	7/20	4/20
(%)	(15)	(25)	(35)	(20)
p =	0.4434	0.3474	0.1367	0.5000
<hr/>				
Follicular Cell Adenomas#	0/20	0/20	1 ^b /20	0/20
(%)	(0)	(0)	(10)	(0)
p =	0.7500	1.0000	0.5000	1.0000

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

^aFirst c-cell adenoma observed at week 102, dose 100 ppm.

^bFirst follicular cell adenoma observed at week 106, dose 500 ppm.

#No c-cell or follicular cell carcinomas were observed.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

17

Table 14. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity and Chronic Toxicity Combined Rat Studies

Female Mortality Rates[†] and Cox or Generalized K/W Test Results

Dose (ppm)	<u>Weeks</u>				Total
	1-26	27-52	53-78	79-107 [‡]	
0	0/70	0/70	3/70	18/67	21/70 (30)
100	0/70	0/70	2/70	9/68	11/70 (16)*
500	0/70	1/70	2/69	15/67	18/70 (26)
2500	0/70	0/70	2/70	9/68	11/70 (16)*

[†]Number of animals that died during interval/Number of animals alive at the beginning of the interval.

[‡]Final sacrifice at week 105.

() Percent.

Note: Time intervals were selected for display purposes only.

Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If †, then $p < 0.05$. If **, then $p < 0.01$.

17

18

Table 15. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity and Chronic Toxicity Combined Rat Studies

Female Thyroid Follicular Cell Tumor Rates* and Exact Trend Test and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
Adenomas [†] (%)	0/70 (0)	1*/70 (1)	1*/69 (1)	3/70 (4)
p =	0.0439*	0.5000	0.4964	0.1223

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

*First adenomas observed concurrently at week 106 at doses 100 and 500 ppm.

*No carcinomas were observed.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

18

Table 16. BAS510F - Wistar Chbb:THOM (SPF) Carcinogenicity and Chronic Toxicity Combined Rat Studies

Female Thyroid C-Cell Tumor Rates* and Exact Trend Test and Fisher's Exact Test Results (p-values)

	<u>Dose (ppm)</u>			
	0	100	500	2500
Adenomas (%)	11/70 (16)	13/70 (19)	14 ^a /69 (20)	9/70 (13)
p =	0.1903	0.4115	0.3153	0.4049
Carcinomas (%)	0/70 (0)	0/70 (0)	1 ^b /69 (1)	0/70 (0)
p =	0.7491	1.0000	0.4964	1.0000
Combined (%)	11/70 (16)	13/70 (19)	15/69 (22)	9/70 (13)
p =	0.1803	0.4115	0.2443	0.4049

*Number of tumor bearing animals/Number of animals examined, excluding those that died before week 53.

^aFirst adenoma observed at week 77, dose 500 ppm.

^bFirst carcinoma observed at week 105, dose 500 ppm.

Note: Significance of trend denoted at control.

Significance of pair-wise comparison with control denoted at dose level.

If *, then $p < 0.05$. If **, then $p < 0.01$.

References

- Cox, D.R. (1972) Regression Models and Life Tables (with discussion). J. Royal Stat. Soc. Ser. B. 34, 187-220.
- Gart, J.J., D. Krewski, P.N. Lee, R.E. Tarone, and J. Wahrendorf (1986) The Design and Analysis of Long-Term Animal Experiments. In: Statistical Methods in Cancer Research, Volume III. IARC Scientific Publications No. 79. Lyon, France: International Agency for Research on Cancer, p. 18.
- Peto, R., M. Pike, N. Day, R. Gray, P. Lee, S. Parish, J. Peto, S. Richard, and J. Wahrendorf (1980) Guidelines for Simple, Sensitive, Significant Tests for Carcinogenic Effects in Long-Term Animal Experiments. In: Monographs on the long-term and short-term screening assays for carcinogens: a critical appraisal. IARC Monographs, Supplement 2. Lyon, France: International Agency for Research on Cancer, pp. 311-426.
- Thomas, D.G., N. Breslow, and J.J. Gart (1977) Trend and Homogeneity Analyses of Proportions and Life Table Data. Computers and Biomedical Research 10, 373-381.