

US EPA ARCHIVE DOCUMENT

Candace Brassard

Amendment to DER
Bobwhite Reproduction Study on Cyhalothrin

The study authors have responded to the data discrepancies that were identified in a data evaluation record dated 3/10/88.

The major concern identified in the study was the high incidence of eggs cracked in the control groups. According to the Environmental Research Laboratory, Corvallis, Oregon, (EPA), eggs cracked in the control should be 2%. EEB has received other acceptable data ranging from 5 to 9% for eggs cracked in the control.

The handling procedures at the Huntington Research facility need vast improvement. Though we have accepted data in the past with levels as high as 15.6% (as the study authors reported for fomesafen), the research facility should not assume that this is a good laboratory standard. Measures should be taken in the future that will correct this. We suggest you contact Rick Bennett, EPA Laboratory in Corvallis, Oregon, at (503)757-4601 for guidance. After personal communications with Rick Bennett, (4/24/89) EEB believes that in this case, that if there was a chemical effect on the percent eggs cracked, that there would have been greater effect in the treated groups (percent eggs cracked) when compared to the control.

An interesting note - it appears Huntington Research Lab has selectively chosen the historical control data that was submitted. EEB is aware that there is at least one other bobwhite reproduction study (1984) which showed percent eggs cracked for the control as being 9%, which Huntington failed to include in their Table 1.

Adequacy of Study

- Classification- This study is classified as CORE for the following parameters:

eggs laid	NOEL= < 50 ppm
eggs cracked	NOEL= < 50 ppm
eggs set	NOEL= < 50 ppm
viable embryos	NOEL= < 50 ppm
normal hatchlings	NOEL= < 50 ppm
14-day survivors	NOEL= < 50 ppm

The study authors should be aware that the decision to accept the eggs cracked data is on a case-by-case basis, and in no way should the research facility believe that future studies with eggs cracked in this range will be acceptable data.

- Rationale- The company has addressed the data discrepancies that were identified in the earlier review.

- Repairability- N/A

Reviewed by: Candace Brassard
Ecological Effects Branch/EFED

Approved by: Douglas J. Urban, Head Section-III
Ecological Effects Branch/EFED

Candace Brassard
4/25/89

Douglas J. Urban
4/26/89

Page 2 is not included in this copy.

Pages _____ through _____ are not included in this copy.

The material not included contains the following type of information:

- Identity of product inert ingredients.
 - Identity of product impurities.
 - Description of the product manufacturing process.
 - Description of quality control procedures.
 - Identity of the source of product ingredients.
 - Sales or other commercial/financial information.
 - A draft product label.
 - The product confidential statement of formula.
 - Information about a pending registration action.
 - FIFRA registration data.
 - The document is a duplicate of page(s) _____.
 - The document is not responsive to the request.
-

The information not included is generally considered confidential by product registrants. If you have any questions, please contact the individual who prepared the response to your request.

AVIAN REPRODUCTION
DER EXAMPLE
BOBWHITE

DATA EVALUATION RECORD

- 1. Chemical: Cyhalothrin
- 2. Test Material: Cyhalothrin 92.2% w/w
 Chemical Structure: (R,S) -cyano-3-phenoxy
 benzyl (+)-cis-3, 3(2-2-chloro-3,3,3-
 trifluoroprop-1-en)-2,2-dimethylcyclo-
 propanecarboxylate
 cis:trans ratio: 96.8:3.2
- 3. Study Type: Avian Reproduction
Species Tested: Bobwhite Quail
 (Colinus virginianus)
- 4. Study ID: Roberts, N.L., Fairley, C., Chanter, D.O.,
 McAllister, A., and Almond, R.H. (1982) The
 Effect of the Dietary Inclusion of Cyhalothrin
 on Reproduction in the Bobwhite Quail. Prepared
 by Huntingdon Research Centre, Huntingdon, England
 - PE 18 6ES; Submitted by ICI Americas, Inc. EPA
 Accession No. 073989.

- 5. Reviewed By: Candy Brassard
 ERP/HRD
 Signature: *Candy Brassard*
 Date: 3-7-88
- 6. Approved By: Douglas J. Urban
 Head-Section III
 ERP/HRD
 Signature: *Douglas J. Urban*
 Date: 3/10/88

7. Conclusion:
 Based on the submitted data it appears that cyhalothrin does not cause reproductive impairment for the number of eggs laid, eggs set, viable embryos, live embryos, normal hatchlings and 14-day survivors at < 50 ppm Cyhalothrin. The statistical analysis indicated the NOEL was < 50 ppm cyhalothrin for eggs cracked as well, however, the percent eggs cracked (and damaged) was reported to be as high as 17 percent for the control. Therefore, the results for this parameter appears to be unreliable. The study appears to be scientifically sound, however there are data discrepancies that cause concerns.

8. Recommendations:

The study author/company should satisfy discrepancies outlined in section 14A. Specifically, the cause for high percent eggs cracked for the control should be addressed.

9. Background:

The study was submitted to support registration of Karate or PP321 on cotton and soybeans.

10. Discussion of Individual Tests: N/A

11. Materials and Methods:

- a. Test Animals - Young adult bobwhite quail that were approaching first laying season, were obtained from Lincolnshire Pheasantries Limited, Boston, Lincolnshire. The birds were acclimated for over 2 months prior to dosing. A total of 51 males and 102 females were used in the study.
- b. Test System - Adults - (excerpted from submission)

Adult birds were housed in treatment replicate groups each consisting of one male and two females. The groups were housed in tiered cages of polythene coated steel wire, each measuring approximately 31.5 cm x 38.5 cm x 24 cm. Each cage contained a stainless steel food hopper and a nipple drinker and had a sloping floor with a 10 cm egg catcher. The maximum and minimum ambient temperature together with the relative humidity was recorded daily throughout the study, with the following values:

	<u>Mean</u>	<u>Range</u>
Relative humidity (%)	67	48 to 85
Maximum temperature (°C)	18	10 to 25
Minimum temperature (°C)	14	8 to 22

Ventilation fans were adjusted as necessary. The following controlled artificial lighting pattern was adopted:

<u>Days of study</u>	<u>Hours light</u>	<u>Hours dark</u>
1 to 70	7	17
71 to 77	8	16
78 to 91	9	15
92 to 98	12	12
99 to 105	13	11
106 to 217	14	10

The birds were fed both basal diet with and without test compound. The diet consisted of the following ingredients:

<u>Ingredient</u>	<u>% w/w</u>
Ground wheat	38.25
Ground maize	30.00
Weatings (Wheatfeed)	5.00
Provimi 66 fishmeal	10.00
Soya bean meal	10.00
Limestone flour	5.50
Pantoribin 537*	1.25

* Mineral, vitamin and trace element supplement
(B.P. Nutrition (U.K.) Ltd.).

Water was available ad libitum from nipple drinkers.

Diet Preparation - The test substance was mixed with corn oil in the final diet. Corn oil at a final rate of 0.1% w/w, was incorporated in treatment as well as the control diets. Diets were mixed on a daily basis. Residue analysis was conducted on the diet to test for stability and homogeneity. Throughout the report, nominal concentrations of cyhalothrin are given.

Eggs - Eggs were incubated on a weekly interval using a Western Incubator. The temperature and humidity were recorded daily as follows: Temperature - 37.7 °C (mean) and humidity ranging from 34 to 69 percent with mean 63.0 percent. Eggs were turned every 45 minutes through an angle of 90° throughout the incubation period. Eggs were incubated for approximately 20 days before transferred to hatcher.

Hatching - The hatchers were Air Bristol Incubator models PH 90 and PH 150, which operated at 37.5 °C. Hatcher trays were made from wooden frames with wire mesh floors. Chicks hatched 24 to 26 days after eggs were first set in the incubator.

Chicks - (following excerpted from submission)

Chicks were housed in wooden pens with concrete floors. Each pen contained two drinkers and two food hoppers. Wood shavings, supplied by the Sawdust Marketing Company Limited, were used as bedding. Each pen contained two 300 watt infra-red lamps placed at bird level to supply additional heat to the chicks. The minimum and maximum ambient temperature together with the relative humidity were recorded once daily and had the following values:

<u>Ingredient</u>	<u>% w/w</u>
Ground wheat	38.25
Ground maize	30.00
Weatings (Wheatfeed)	5.00
Provimi 66 fishmeal	10.00
Soya bean meal	10.00
Limestone flour	5.50
Pantoribin 537*	1.25

* Mineral, vitamin and trace element supplement
(B.P. Nutrition (U.K.) Ltd.).

Water was available ad libitum from nipple drinkers.

Diet Preparation - The test substance was mixed with corn oil in the final diet. Corn oil at a final rate of 0.1% w/w, was incorporated in adults as well as the control diets. Diets were mixed on a daily basis. Residue analysis was conducted on the diet to test for stability and homogeneity. Throughout the report, nominal concentrations of cyhalothrin are given.

Eggs - Eggs were incubated on a weekly interval using a Western Incubator. The temperature and humidity were recorded daily as follows: Temperature - 37.7 °C (mean) and humidity ranging from 34 to 69 percent with mean 63.0 percent. Eggs were turned every 45 minutes through an angle of 90° throughout the incubation period. Eggs were incubated for approximately 20 days before transferred to hatcher.

Hatching - The hatchers were Air Bristol Incubator models PH 90 and PH 150, which operated at 37.5 °C. Hatcher trays were made from wooden frames with wire mesh floors. Chicks hatched 24 to 26 days after eggs were first set in the incubator.

Chicks - (following excerpted from submission)

Chicks were housed in wooden pens with concrete floors. Each pen contained two drinkers and two food hoppers. Wood shavings, supplied by the Sawdust Marketing Company Limited, were used as bedding. Each pen contained two 300 watt infra-red lamps placed at bird level to supply additional heat to the chicks. The minimum and maximum ambient temperature together with the relative humidity were recorded once daily and had the following values:

	<u>Mean</u>	<u>Range</u>
Relative humidity (%)	48	31 to 72
Maximum temperature (°C)	38	28 to 46
Minimum temperature (°C)	31	22 to 39

A continuous artificial lighting pattern was adopted for the chicks. Ventilation fans were adjusted as required.

Feeding - The chicks were given standard HRC chick diet in meal form made by Joseph Odam, Ltd., Eye Mill, Peterborough, Cambridgeshire, England.

Diet composition

<u>Ingredient</u>	<u>% w/w</u>
Ground wheat	30.00
Ground maize	25.00
Ground barley	10.00
Provimi 66 fishmeal	15.00
Extracted soya bean meal	13.75
Weatings (Wheatfeed)	5.00
Pantoribin 537*	1.25

* Mineral, vitamin and trace element supplement
(B.P. Nutrition (U.K.) Ltd.).

- c. Dose - A control and two treatment levels were used - 5 ppm and 50 ppm cyhalothrin.
- d. Design - There were 14 pens per treatment group, each containing two females and one male, a total of 28 females and 14 males per dose level. In addition, a number of spare birds (three males and six females per treatment group) were maintained on each of the three treatments for use as replacement birds if necessary during the pre-egg production period.

Observations -

Adult Bird Observations were as follows:

- Mortalities - daily.
- Bird health - daily.
- Gross mean food consumption - weekly per replicate.

- Individual body weights - days 0*, 14, 28, 42, 56, 70, 84, 98, 112, and 217.
- Gross macroscopic postmortem examination - all birds were examined postmortem for gross abnormalities.

* At the start of day 1 immediately prior to the introduction of test diets.

Egg Observations were as follows:

- Egg collection - Eggs were collected daily throughout the 12-week egg production period.
- Egg weights - At 7-day intervals the collected eggs were weighed. Broken eggs were not weighed.

Egg Quality - Eggs cracked (and damaged) - At 7-day intervals the collected eggs were candled after weighing, to check for cracks and breakages. Any other shell abnormalities were noted at this stage.

Egg Shell Thickness - All eggs collected in the first 2 days of weeks 1, 3, 5, 7, 9, and 11 were examined. The eggs were cracked open at the widest point and the contents washed out with tap water. The shells were then "left to dry at room temperature for a minimum of 48 hours. The shell thickness of each egg was measured at four points around the circumference of the shell using a micrometer calibrated to 0.01 mm."

Incubation (excerpted from submission):

Incubation - All eggs remaining after the cracked, broken, abnormal eggs and those taken for shell thickness had been removed were placed in the incubator at weekly intervals.

Candling and Hatching (excerpted from submission):

The incubated eggs were examined at days 11 and 18 of the incubation period by passing over a light source in a darkened room (candling). The following parameters were recorded:

1. Infertile eggs - Appearing as "clears" at the Day 11 candling.

2. Early embryonic mortalities - At the Day 11 candling any embryos observed to be dead were removed. The embryos at this stage were not fully differentiated.
3. Late embryonic mortalities - At the Day 18 candling any embryos observed to be dead were removed. At this stage the embryos were fully differentiated.
4. "Dead in shells" - Any eggs which failed to hatch after the embryonic deaths and infertiles had been removed were recorded as "dead in shells." These eggs generally contained chicks which appeared to be fully formed and viable but failed to get out of the shell. Pipped eggs, i.e., chicks which had been able to crack the shell but had been too weak to get out, were also noted.
5. Chicks hatched - The chicks hatched alive were recorded. In addition, those which hatched but were found dead in the hatcher were also recorded. Any abnormalities were also noted at this stage.

Chicks - All chicks hatched alive were reared until they were 14 days old and the following parameters were recorded:

1. Individual bodyweight - Within 24 hours of hatching and 14 days later.
2. Bird health - Daily.
3. Mortalities - Daily.
4. Gross macroscopic postmortem examination - Only sporadic mortalities were examined for gross abnormalities. No examination was made at termination.

Summary of Study Duration (excerpted from submission):

Adults 19-week pre-egg production period
 12-week egg production period.

Incubation Eggs collected over the 12-week egg
 production period were incubated
 weekly. The incubation period lasted
 approximately 24 to 26 days.

Chicks

The weekly hatches of chicks from the 12-week egg production period were reared until they were 14 days old.

The total study duration from the start of the adult observation period to the final chick observations was approximately 36 weeks.

e. Statistical Analysis - (excerpted from submission)

A statistical analysis of the following responses was carried out:

1. Adult food consumption
2. Adult bodyweight
3. Number of eggs laid and proportion damaged
4. Egg weight
5. Egg shell thickness
6. Number of infertilities, embryonic mortalities and hatchings
7. Number of 14-day old surviving chicks
8. Chick bodyweights

12. Reported Results:

Mortalities and Bird Health

No evidence of any treatment-related response in the number of mortalities occurring in each group. Birds which died during pre-egg production period (Days 1 through 133) were replaced by sparebirds. No replacements were made during egg production period (Days 134 through 217). (See Table 1.)

<u>Bird No. and Sex</u>	<u>Replicate/ Group</u>	<u>Day of Death</u>	<u>Replacement Bird No.</u>
325M	9A	127	461M
338F	13A	76	462F
412M*	38A	98	446M
425F	42A	6	466F
322M	8B	35	467M
340M	14B	67	455M
310M	4C	22	452M
345F*	15C	9	453F

* Signs of "bullying" recorded prior to death.

Bird health observations are in Appendix 2.

Adult Body Weights - All body weights were within normal limits and no treatment-related effects were found. (See Table 2.)

Food Consumption - Food consumption was within normal limits in all groups throughout the study and no-treatment-related effects were observed. (See Tables 3 and 4.)

Gross Postmortem Examination - Sporadic mortalities. Two birds appeared to have died from bullying. Bird No. 412M (Replicate group 38A) and 345F (Replicate 15C).

Terminal Findings - Pale livers were noted in a number of birds during postmortem. This was observed in all groups and was not considered to be an abnormality.

The following observations were also made:

- Ovary underdeveloped - four in control, four in lowest dose level, and one in highest dose level.
- Testes underdeveloped - one in lowest dose and one in highest dose.
- Liver blotchy - five in control, one in each dose level (5 ppm and 50 ppm).
- Ovary developed but no eggs in oviduct - one in control, one in lowest dose level.
- Egg bound - one at highest dose tested.

Eggs Laid - The total number of eggs laid were similar for the control and lowest dose (5 ppm) and slightly higher for highest dose (50 ppm). Statistical analysis of the results showed no significant difference between treatments. (See Table 5.)

Eggs Cracked (and Broken) (excerpted from submission) - No significance differences between treatments were detected in the proportions of cracked and broken eggs. (See Table 6.)

Egg Weights (excerpted from submission) - Statistical analysis of the egg weight data showed that there were no significant differences between treatments in total egg mass or mean egg weights. (See Table 7.)

Egg Shell Thickness (excerpted from submission) - Egg shell thickness was within normal limits for both control and test groups and statistical analysis of the results showed no differences between treatments. (See Table 8.)

Infertile eggs (excerpted from submission) - The proportions of eggs incubated which were found to be infertile at Day 11 candling varied within treatment groups during the 12-week egg production period. Overall, the proportion of infertile eggs was lower in Group C (Cyhalothrin at 50 ppm) than in Groups A and B (Control and Cyhalothrin at 5 ppm). Statistical analysis of the results showed no significant differences between treatments. (See Tables 9 and 10.)

Early embryonic mortalities - The proportions of early embryonic mortalities occurring in fertile eggs were variable, but similar overall in all treatment groups. No significant differences between treatments were detected. (See Tables 9 and 10.)

Late embryonic mortalities - The proportions of late embryonic mortalities were small in all treatment groups. Statistical analysis of the results was not practicable. (See Tables 9 and 10.)

Hatching (excerpted from submission) - The proportions of fertile eggs which subsequently hatched were generally high and were similar overall in all groups. Statistical analysis of the results showed no significant treatment differences. (See Table 11.)

Chick Health and Mortalities (excerpted from submission) - Chick health was generally good and the numbers of mortalities occurring were within normal limits. Details of mortalities are given in Appendix 7.

The following observations on chick health were made:

<u>Bird No.</u>	<u>Replicate/Group</u>	<u>Week of hatch</u>	<u>Observations</u>
2G (blue)	19B	2	Sacrificed on Day 1 as very weak, with splayed legs.
62N (green)	38A	12	Bird subdued on Day 13, prior to death on Day 14.

Number of 14-Day Survivors (excerpted from submission) - The proportion of chicks surviving to 14 days was generally high in all groups and statistical analysis showed no significant differences between treatments. (See Table 12.)

Bodyweights (excerpted from submission) - All mean chick bodyweights at hatching and after 14 days were within normal limits and no statistically significant differences were found between treatments. (See Table 13.)

Postmortem Examination

No abnormalities other than those mentioned in chick health section, were detected in any chicks during postmortem examination.

13. Study Authors' Conclusions/OA Measures (excerpted from submission):

Under the conditions of this study there was no evidence that dietary administration of cyhalothrin at dose levels of 5 ppm and 50 ppm had any adverse effects on reproduction in the Bobwhite quail.

To the best of my knowledge and belief, this study was conducted in compliance with Good Laboratory Practice regulations as set forth in "Title 21, of the U.S. Code of Federal Regulations, Part 58", with the exception of possible minor items, none of which is considered to have an impact on the validity of the data or the interpretation of the results in the report. (Signed by N.L. Roberts.)

14. Reviewer's Discussion and Interpretation of the Study:

The following discrepancies were noted in the study:

- a. Test Procedures - The percent cracked of eggs laid in the control is of real concern. Typically, 0.6 to 2.0 percent is seen and may be as high as 5-6percent, and this study author reported the percent cracked and damaged to be as high as 17 percent. The percentage eggs cracked, not including broken, was as high as 15.0 percent. While it is true that there is no statistically significant difference between the control and two treatment groups the percent eggs cracked of eggs laid in the control is considerably higher than the normal limit. No historical control data were submitted. Another study submitted by this Laboratory to EPA, specifically for Rifenhrin, reported 9 percent eggs cracked in the control (review completed by Les Touart on October 13, 1987). Rick Bennett with U.S. EPA Environmental Research Laboratory, Corvallis, OR, reported the percent eggs cracked for the bobwhite, with sloped cage facilities, to be less than 5 percent and usually only 1 to 2 percent. The study author should report the cause for such a high percent eggs cracked in the control, and should also submit the historical control data.
- Exact age not reported - only that the birds were approaching their first laying season.

- The percent calcium and phosphorous in the adult diet should be indicated.
 - Typically, a 12 week pre-egg production period is used with bobwhite reproduction studies. The study author should indicate why a 19-week pre-egg production period was used.
 - It was not reported if the adult diet was available ad libitum.
 - The study author reported temperature ranges of 8 to 25 °C and relative humidity ranges of 48 to 85 percent. The recommended levels (McLane, D. 1986) are 21 °C and 55 percent relative humidity. The study author should account for the temperature variation.
 - Provisions made to avoid food spillage were not reported.
 - The study author did not indicate the size of chick pens.
 - In appendix 5, week 1 for the control - Group A, there was a 32A for a replicate number. FFR is assuming that the study author intended it to read 38A. This should be clarified.
 - The percent live embryos when compared to viable embryos should be around 97 to 99 percent. This study was reported to be as low as 94.2 percent for the control.
- b. Statistical Analysis - The following parameters were evaluated using an ANOVA program and Duncan's multiple range test: eggs laid, eggs cracked, eggs set, viable embryos, live embryos, normal hatchlings, and 14-day-old survivors.

The results are as follows:

Eggs laid = NOEL > 50 ppm

Eggs cracked = NOEL > 50 ppm

Eggs set = NOEL > 50 ppm (These results may not be reliable.)

Viable embryos = NOEL > 50 ppm

Live embryos = NOEL > 50 ppm

Normal hatchings = NOEL > 50 ppm

14-day survivors = NOEL > 50 ppm

RRR analyzed the reproductive effects. (See Table A.)

c. Discussion/Results - The percent eggs cracked for the control is considerably high, even for sloped cage facilities. The study author should account for this discrepancy, along with submitting historical control data for this study. The study author should report the reason for such a wide variation in the temperature used for the accommodations of the birds. The study author should also indicate why the unusually long pre-egg production period was used. There are numerous other minor data discrepancies listed in section 14 of this review.

d. Adequacy of Study

- 1) Classification - Supplemental for 92.2% w/w cyhalothrin.
- 2) Rationale - This study appears to be scientifically sound; however, the data discrepancies outlined in 14.c. detract from the study.
- 3) Repairability - Repairability pending the data submitted to satisfy discrepancies outlined in Section 14.a. and c.

Table A
Analysis of Reproductive Effects

	<u>Concentrations of Cyhalothrin</u>		
	<u>Control</u>	<u>5 ppm</u>	<u>50 ppm</u>
<u>Eggs laid</u>	1280	1287	1345
Eggs laid/hen/season*	46.9	47.6	49.6
<u>Eggs cracked**</u>	217	211	277
Eggs cracked/hen/season	6.9	7.8	10.1
Percent of eggs laid	17%	16.4%	20.6%
<u>Eggs set</u>	963	967	966
Percent of eggs laid	75.2%	75.1%	71.8%
<u>Viable Embryos (11-Day)</u>	768	780	848
Percent of eggs laid	60%	60.6%	63.0%
Percent of eggs set	79.7%	80.6%	87.7%
<u>Live 18-Day Embryos</u>	724	714	804
Percent of viable embryos	94.2%	91.5%	94.8%
<u>Hatchlings</u>	664	660	732
Percent of eggs laid	51.8%	51.3%	54.4%
Percent of eggs set	69.0%	68.3%	75.8%
Percent of viable embryos	86.5%	84.6%	86.3%
Percent of 18-day embryos	91.7%	92.4%	91.0%
<u>14-Day Survivors***</u>	550	525	613
Percent of normal hatchlings	83%	80%	84%
<u>Average hatchweight (g)</u>	7	7	7
<u>Average 14-Day-Old Survivors weight (g)</u>	19	19	20
<u>Adult Body Weight (g/Bird) (at study termination)</u>			
Females	226	225	224
Males	209	205	208

	<u>Control</u>	<u>5 ppm</u>	<u>50 ppm</u>
<u>Adult Body Weight (g/Bird)</u>			
<u>Increase compared to Day 0</u>			
Females	+33	+33	+31
Males	+ 9	+ 6	+ 8
<u>Mean Eggshell Thickness</u>	0.195	0.19	0.198
<u>Mean Egg Weight</u>	10.0	10.0	10.0
<u>Average Feed Consumption</u>			
Pre-egg production period	17.2	17.1	18.0
Egg production period	22.1	24.3	25.3
Mean Total	19.1	19.9	20.8

* The number of females per week were used to estimate number per pen. Therefore, the mortalities were included.

** Eggs cracked include all broken, damaged and cracked eggs.

***Number of survivors per hen could not be calculated since there were mortalities within each treatment level (and control).

Page _____ is not included in this copy.

Pages 19 through 25 are not included in this copy.

The material not included contains the following type of information:

- Identity of product inert ingredients.
 - Identity of product impurities.
 - Description of the product manufacturing process.
 - Description of quality control procedures.
 - Identity of the source of product ingredients.
 - Sales or other commercial/financial information.
 - A draft product label.
 - The product confidential statement of formula.
 - Information about a pending registration action.
 - FIFRA registration data.
 - The document is a duplicate of page(s) _____.
 - The document is not responsive to the request.
-

The information not included is generally considered confidential by product registrants. If you have any questions, please contact the individual who prepared the response to your request.

	ISS	TRT	EL	EC	ES	VE	LE	NH
426								
427								
428								
429	1	A	93	24	70	66	65	52
430	2	A	112	18	72	67	67	56
431	3	A	65	4	56	56	56	47
432	4	A	95	14	61	26	26	21
433	5	A	123	16	94	77	77	68
434	6	A	74	1	67	18	16	13
435	7	A	73	4	63	60	54	49
436	8	A	83	15	60	22	21	16
437	9	A	93	13	73	62	59	54
438	10	A	110	19	93	59	59	55
439	11	A	122	13	97	87	88	74
440	12	A	79	27	43	33	33	26
441	13	A	91	20	65	64	64	56
442	14	A	72	23	46	40	41	35
443	15	B	85	20	59	42	37	31
444	16	B	44	14	28	28	29	28
445	17	B	133	24	99	67	67	64
446	18	B	72	7	57	52	51	47
447	19	B	108	6	91	9	8	7
448	20	B	53	5	44	23	23	23
449	21	B	94	19	61	46	51	49
450	22	B	101	14	79	70	70	65
451	23	B	93	25	61	57	57	56
452	24	B	130	27	91	86	86	73
453	25	B	120	14	93	59	63	61
454	26	B	126	18	99	94	92	87
455	27	B	34	3	29	18	19	14
456	28	B	94	15	71	63	62	54
457	29	C	102	23	73	41	41	38
458	30	C	100	26	84	67	66	58
459	31	C	97	17	76	66	70	63
460	32	C	78	9	63	52	61	47
461	33	C	40	4	33	22	20	16
462	34	C	108	23	75	64	63	58
463	35	C	117	14	90	73	73	69
464	36	C	119	30	77	73	71	54
465	37	C	127	27	70	53	53	51
466	38	C	120	39	72	70	70	69
467	39	C	67	16	45	44	44	42
468	40	C	102	14	88	71	70	60
469	41	C	101	20	74	66	66	64
470	42	C	67	9	51	46	45	44

*Big Bird
 Cylalobus
 Bobwhite*

1. ANALYSIS OF EL DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988

GENERAL LINEAR MODEL PROCEDURE

CLASS LEVEL INFORMATION

CLASS LEVELS VALUES

TRT 3 A B C

26

 GENERAL LINEAR MODELS PROCEDURE

486
 487
 488
 489 DEPENDENT VARIABLE: RESP

490
 491 SOURCE

DF	SUM OF SQUARES	MEAN SQUARE	F VALUE	PR > F	R-SQUARE	DF
2	181.85714286	90.92857143	0.14	0.8715	0.007070	27.850
39	13869.14285714	355.6191037				
41	13869.14285714					

492
 493 MODEL
 494 ERROR
 495 CORRECTED TOTAL

496 SOURCE

DF	TYPE I SS	F VALUE	PR > F	DF	TYPE III SS	F VALUE	PR
2	181.85714286	0.14	0.8715	2	181.85714286	0.14	0.871

1. ANALYSIS OF EL DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988

 GENERAL LINEAR MODELS PROCEDURE

DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESP
 NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONWISE ERROR RATE,
 NOT THE EXPERIMENTWISE ERROR RATE

ALPHA=0.05 DF=39 MSE=658.648

NUMBER OF MEANS: 2 3
 CRITICAL RANGE: 19.5197 20.6254

MEANS WITH THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT.

DUNCAN	GROUPING	MEAN	N	TRT
	A	96.071	14	C
	A	91.929	14	B
	A	91.429	14	A

2. ANALYSIS OF EC DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988

 GENERAL LINEAR MODELS PROCEDURE

CLASS LEVEL INFORMATION

CLASS	LEVELS	VALUES
TRT	3	A B C

NUMBER OF OBSERVATIONS IN DATA SET = 42

2. ANALYSIS OF EC DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988

NOEL 7502011

27

527
528
529
530
531
532
533
534
535
536
537
538
539

GENERAL LINEAR MODELS PROCEDURE

CLASS LEVEL INFORMATION

CLASS	LEVELS	VALUES
TRT	3	4 3 3

NUMBER OF OBSERVATIONS IN DATA SET = 42
1. ANALYSIS OF EC DATA

10:53 WEDNESDAY, FEBRUARY 24, 1988

540
541
542
543

GENERAL LINEAR MODELS PROCEDURE

544 DEPENDENT VARIABLE: RESP

545

SOURCE	DF	SUM OF SQUARES	MEAN SQUARE	F VALUE	PR > F	R-SQUARE	CLV
MODEL	2	171.42857143	85.71428571	1.23	0.3020	0.059555	50.493
ERROR	39	2707.97142857	69.41208791				RESP MEA
CORRECTED TOTAL	41	2878.50000000				3.33139172	16.5000000

553
554

SOURCE	DF	TYPE I SS	F VALUE	PR > F	DF	TYPE III SS	F VALUE	PR >
TRT	2	171.42857143	1.23	0.3020	2	171.42857143	1.23	0.302

558

2. ANALYSIS OF EC DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988

559
560

GENERAL LINEAR MODELS PROCEDURE

DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESP

NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONWISE ERROR RATE,
NOT THE EXPERIMENTWISE ERROR RATE

ALPHA=0.05 DF=39 MSE=69.4121

NUMBER OF MEANS	2	3
CRITICAL RANGE	6.36716	8.09565

MEANS WITH THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT.

DUNCAN GROUPING	MEAN	N	TRT
A	15.757	14	C
A	15.071	14	A
A	15.071	14	B

3. ANALYSIS OF ES DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988

EC
HOELT
28

581

GENERAL LINEAR MODELS PROCEDURE

CLASS LEVEL INFORMATION

CLASS LEVELS VALUES

TRT 3 A B C

NUMBER OF OBSERVATIONS IN DATA SET = 42
3. ANALYSIS OF ES DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988

GENERAL LINEAR MODELS PROCEDURE

DEPENDENT VARIABLE: RESP

SOURCE	DF	SUM OF SQUARES	MEAN SQUARE	F VALUE	PR > F	R-SQUARE	C.V.
MODEL	2	30.61904762	15.30952381	0.04	0.9627	0.001950	29.704
ERROR	39	15671.21428571	401.82600733				RESP MEA
CORRECTED TOTAL	41	15701.83333333			20.04559820		69.8333333

SOURCE	DF	TYPE I SS	F VALUE	PR > F	DF	TYPE III SS	F VALUE	PR > F
TRT	2	30.61904762	0.04	0.9627	2	30.61904762	0.04	0.962

3. ANALYSIS OF ES DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988 1

GENERAL LINEAR MODELS PROCEDURE

DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESP
NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONWISE ERROR RATE,
NOT THE EXPERIMENTWISE ERROR RATE

ALPHA=0.05 DF=39 MSE=401.826

NUMBER OF MEANS 2 3
CRITICAL RANGE 15.3244 16.11

MEANS WITH THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT.

DUNCAN	GROUPING	MEAN	N	TRT
	A	70.786	14	C
	A	70.000	14	A
	A	68.714	14	B

4. ANALYSIS OF VE DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988 1

*Egg-net
750 p/12*

29

641
642
643
644
645
646
647
648
649

650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668

669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691

692
693

CLASS LEVEL INFORMATION

CLASS LEVELS VALUES

TRT 3 A B C

NUMBER OF OBSERVATIONS IN DATA SET = 42
4. ANALYSIS OF VE DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988

GENERAL LINEAR MODELS PROCEDURE

DEPENDENT VARIABLE: RESP

SOURCE	DF	SUM OF SQUARES	MEAN SQUARE	F VALUE	PR > F	R-SQUARE	DF
MODEL	2	343.00000000	171.50000000	0.39	0.6785	0.019891	38.814
ERROR	39	17076.07142857	437.84798535				RESF MEA
CORRECTED TOTAL	41	17419.07142857			20.92481745		53.7857142

SOURCE	DF	TYPE I SS	F VALUE	PR > F	DF	TYPE III SS	F VALUE	PR >
TRT	2	343.00000000	0.39	0.6785	2	343.00000000	0.39	0.678

4. ANALYSIS OF VE DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988 1

GENERAL LINEAR MODELS PROCEDURE

DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESP
NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONWISE ERROR RATE,
NOT THE EXPERIMENTWISE ERROR RATE

ALPHA=0.05 DF=39 MSE=437.848

NUMBER OF MEANS 2 3
CRITICAL RANGE 15.9766 16.9166

MEANS WITH THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT.

DUNCAN	GROUPING	MEAN	N	TRT
	A	57.714	14	C
	A	52.643	14	A
	A	51.000	14	B

5. ANALYSIS OF LE DATA

13:58 WEDNESDAY, FEBRUARY 24, 1988 1

GENERAL LINEAR MODELS PROCEDURE

*Multiple Comparisons
Note: 750000*

30

CLASS LEVELS VALUES

TRT 3 A B C

NUMBER OF OBSERVATIONS IN DATA SET = 42

5. ANALYSIS OF LE DATA 15

GENERAL LINEAR MODELS PROCEDURE

DEPENDENT VARIABLE: REEF

SOURCE	DF	SUM OF SQUARES	MEAN SQUARE	F VALUE	PR > F
MODEL	2	411.76190476	205.88095238	0.47	
ERROR	39	53.06666667	1.36068376		
CORRECTED TOTAL	41	464.82857143			

SOURCE	DF	TYPE III SS	F VALUE	PR > F
TRT	2	411.76190476	0.47	0.6315

5. ANALYSIS OF LE DATA 13:58 WEDNESDAY, FEBR988 16*****

GENERAL LINEAR MODELS P DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RES

NOT THE EXPERIMENTWISE ERROR RATE

NUMBER OF MEAN 16.0841 16.9086

MEANS WITH THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT.

DUNCAN GROUPING MEAN 741

A 58.071 14 C A 51.857 14 A

6. ANALYSIS OF NH DATA

GENERAL LINEAR MODELS PROCEDURE

CLASS LEVEL LEVELS TRT 3 A B C

ANALYSIS OF NH DATA 13:58 WEDNESDAY, FEBRUARY 24, 1988 18 R MODELS PR 763

SOURCE	DF	SUM OF SQUARES	MEAN SQUARE	F VALUE	PR > F
MODEL	17	47.95238095	2.82072241	0.61	
ERROR	25	14523.57142857	580.94102674		

SOURCE	SS	F VALUE	PR > F
TRT	456.33333333	0.5470	

6. ANALYSIS OF NH D

GENERAL LINEAR MODELS PROCEDURE

31 ***** 31

NOT THE EXPERIMENTWIS

ALPHA=0.05 DF=39 MSE=372.399

NUMBER OF MEANS 2 3
CRITICAL RANGE 14.7526 15.5089

MEANS WITH THE SA

DUNCAN GROUPING

MEAN N

TRT

795
796 14 C
797
798 A

800 A 47.071 799
44.429 14 A

7. ANALYSIS OF ES/EL DATA

13:5, FEBRUARY 24, 1988

GENERAL LINEAR MODELS PROCEDURE

CLASS LEVEL INFORMATION

809
810 NUMBER OF OBSERVATIONS IN DATA SET = 42 TRT 3 A B C
814 RUARY 24, 1988 21

L LINEAR MODELS PROCEDURE

819 DEPENDENT VARIABLE: RESPONSE

820 WEIGHT: WT

821

822 SOURCE

DF

SUM OF SQUARES

MEAN SQUARE

F VALUE

PR > F

R-SQUARE

C.V

823
824 MODEL 1424.26660590 0.55 0.5787 0.027658 84.1400
825

826 ERR 60.22452366 ROOT MSE NSE MEANTOTAL 41 102990.44411397

829

830

831 SOURCE

DF

TYPE I SS

F VALUE

PR > F

DF

TYPE III SS

F VALUE

PR

832

833 TRT 2 1 0.55 0.5787

834

7. ANALYSIS OF ES/EL DATA

13:5

836

837

838

839

841

844

845

847

848

851

852

854 60.17 14 B

855

856 14 C

857

7. ANALYSIS OF ES/EL DATA

13:

859

860 VARIABLE

N

MEAN

STANDARD

MINIMUM

MAXIMUM

STD ERROR

SUM

VARIANCE

C.V

861

862

DEVIATION

VALUE

VALUE

OF MEAN

32

GENERAL LINEAR MODELS PROCEDURE

DEPENDENT VARIABLE: RESP

SOURCE	DF	SUM OF SQUARES	MEAN SQUARE	F VALUE	PR > F	R-SQUARE	CV
MODEL	2	298.61904762	149.30952381	0.46	0.6326	0.003211	14.811
ERROR	39	12567.0000000	322.23076923				
CORRECTED TOTAL	41	12865.61904762			17.95078743		40.0000000

SOURCE	DF	TYPE I SS	F VALUE	PR > F	DF	TYPE III SS	F VALUE	PR
TRT	2	298.61904762	0.46	0.6326	2	298.61904762	0.46	0.632

SAS 15:54 WEDNESDAY, FEBRUARY 24, 1958

GENERAL LINEAR MODELS PROCEDURE

DUNCAN'S MULTIPLE RANGE TEST FOR VARIABLE: RESP
 NOTE: THIS TEST CONTROLS THE TYPE I COMPARISONWISE ERROR RATE,
 NOT THE EXPERIMENTWISE ERROR RATE

ALPHA=0.05 DF=39 MSE=322.231

NUMBER OF MEANS 2 3
 CRITICAL RANGE 15.723 14.4264

MEANS WITH THE SAME LETTER ARE NOT SIGNIFICANTLY DIFFERENT.

DUNCAN	GROUPING	MEAN	N	TRT
	A	43.786	14	C
	A	39.571	14	A
	A	37.357	14	B

14 DAY SURVIVAL
DAY =
Cyhalothrin
Babuvate