


US EPA ARCHIVE DOCUMENT

DATA EVALUATION REPORT

1. Chemical: Mycostop™ - Streptomyces griseoviridis
2. Test Material: Technical
3. Study/Action Type: Nontarget Honey Bee (Apis mellifera) Testing (154A-24)
4. Study Identification: A Dietary Pathogenicity and Toxicity Study with the Honey Bee. By K. A. Hoxter, G. J. Smith and M. Jaber. Prepared By Wildlife International LTD, November 1990. Project No. 293-101. Submitted By Kemira Oy. Helsinki, Finland EPA Acc. No. 4184211-23.
5. Reviewed By: David C. Bays
Microbiologist
EFED/EEB

Les W. Touart
Head, Section 1
EFED/EEB

Signature: 
Date: 9/8/92

Signature:
Date:
6. Conclusions: The study is scientifically sound and demonstrated an $LC_{50} > 2400$ ppm. However, due to nonspecific mortality in the controls, the study will be considered supplemental. The results were reliable enough to indicate that Mycostop is practically nontoxic to Honey Bee.
7. Recommendations: N/A
8. Background: This study was submitted to support the request for the registration of Streptomyces griseoviridis.
10. Materials and Methods:
 - A. Test Organisms: The test bees were obtained from the Wildlife International Ltd. hives located in Easton, Maryland. One frame of pupae was taken from the hives (3 days before test initiation) and placed in a Marsh Roll-X automatic incubator for 3 day/s to allow the adult bees to emerge. The bees used in the test were 1 to 3 days old and were healthy in appearance.

- B. Dosage Form: The test diets were prepared by mixing together a calculated amount of Mycostop (specific activity of 9.8×10^8 cfu/g diet) and honey. The nominal concentrations used were 240, 276 and 2400 ppm with no adjustment for purity of the test substance. The attenuated control was prepared by autoclaving a portion of the test substance (highest concentration tested) at 121C for 30 minutes.
- C. Referenced Protocol: The test insects were placed in disposable one pint rolled paper containers (87 mm in diameter/85 mm high) that were covered with a disposable plastic petri dish (90 mm in diameter). The test diet (available ad libitum) was placed in a 20 ml glass vial which was covered with cheese cloth, and then inserted into the container's cover. A moist sponge, which was misted daily, was placed on the top of each container to increase humidity within the test chamber.

Two replicates, containing 25 insects each, were randomly assigned to each of 3 treatment levels (240, 760 and 2400 ppm) along with the attenuated and negative (untreated honey) controls. The bees were immobilized with nitrogen at the start of the study and the test diets were placed atop the test chambers. The test insects were observed for mortality and signs of toxicity twice on the day the experiment started (first observation immediately following the introduction of the test diets) and once a day thereafter until the end of the study. The study was terminated when the negative control mortality exceeded 20%. The environmental conditions were as follows: 8 hours of light/day, a temperature of 22-24C, and an average relative humidity of 82%.

- D. Statistical Analysis: After study completion, an estimation of the LC_{50} value was made by visual inspection of the mortality data. A calculation of the LC_{50} value was not necessary because of the lack of mortalities associated with the test substance found in this study.

12. Reported Results:

<u>Dosage</u>	<u>ppm</u>	<u>Replicate</u>	<u>Number Dead/Number Exposed</u> <u>(At 5 Days After Dosing)</u>
Negative control	0	A	7/25
		B	8/25
Attenuated control	2400	A	15/25
		B	16/25

Treatment	240	A	4/25
		B	6/25
	760	A	11/25
		B	2/25
	2400	A	5/25
		B	9/25

LC₅₀ > 2400 ppm

Mortalities occurred in both of the control groups (negative and attenuated) and in all 3 of the treatment groups. The mortalities in the negative and attenuated control groups were 30% and 62%, respectively, while those in the 240, 760, and 2400 ppm concentrations averaged 20%, 26% and 28%, respectively. The pattern of mortality was found not to be dose responsive and did not appear to be treatment related. The LC₅₀ was determined to be greater than 2400 ppm and the no effects concentration was 2400 ppm which was the highest concentration tested.

13. Study Author's Conclusions/Quality Assurance Measures:

LC₅₀ > 2400 ppm

"This study was conducted so as to conform with Good Laboratory Practices as published by the U.S. Environmental Protection Agency, Office of Pesticide Programs in 40 CFR Part 160." Signed by study director, Kimberly A. Hoxter.

14. Reviewer's Discussion and Interpretation of the Study:

- A. Test Procedures: The procedures used follow those recommended by EPA in the 1989 Pesticide Testing Guidelines for Microbial and Biochemical Pest Control Agents, Subdivision M.
- B. Statistical Analysis: None was needed since the pattern of mortality did not facilitate the calculation of an LC₅₀ value.
- C. Discussion/Results: An LC₅₀ > 2400 ppm indicates that Mycostop is practically non-toxic to Honey Bee.

D. Adequacy of the Study:

1. Validation Category: Supplemental
2. Rationale: Nonspecific mortality occurred in the controls.

15. Completion of the One-liner: