

US EPA ARCHIVE DOCUMENT

Study Title

Gas Chromatographic Method for Determination of Residues of
Tebuconazole in Crops, Processed Products, Soil and Water

Data Requirement

EPA Ref: 171-4(c), Residue Analytical Method-Plants

Author

W. Maasfeld

BB-16

MRD 422095-05

Revisions By

R. G. Minor

Completion Date

January 15, 1992

Originating Laboratory

BAYER AG
Crop Protection-Research
Chemical Product Development
and Environmental Biology
Institute for Product Information
and Residue Analysis
D-5090 Leverkusen - Bayerwerk, Germany

Submitting Laboratory

Miles Inc.
Agriculture Division
Research and Development Department
Miles Research Park
17745 S. Metcalf Ave
Stilwell, KS 66085

Miles Report Number

101341

Tebuconazole is the common name for FOLICUR®
FOLICUR® is a Reg. TM of Bayer AG, Germany

Statement of Confidentiality

No claim of confidentiality is made for any information contained in this study on the basis of its falling within the scope of FIFRA #10(d)(1)(A), (B) or (C).

Company: Miles Inc.
Agriculture Division
Research and Development Department
Environmental Research Section

Company Agent: D. R. Flint Date: 1-17-97
D. R. Flint, Ph.D.
Manager, Environmental Research

These data are the property of the Agriculture Division of Miles Inc., and as such, are considered to be confidential for all purposes other than compliance with FIFRA 10. Submission of these data in compliance with FIFRA does not constitute a waiver of any right to confidentiality which may exist under any other statute or in any other country.

Certification of Good Laboratory Practice

Good laboratory practice requirements of 40 CFR Part 160 do not apply to the study described in this document.

Submitter: Miles Inc.
Agriculture Division
Research and Development Department
Environmental Research Section

D. R. Flint
D. R. Flint, Ph.D.
Manager, Environmental Research

Date: 1-17-92

Sponsor
Representative:

J. J. Murphy
J. J. Murphy, Ph.D.
Manager, Metabolism/Residue Methodology

Date: 1/17/92

Preparer:

R. G. Minor
R. G. Minor
Chemical Specialist

Date: 1/17/92

Certification of Authenticity

Preparer: R. G. Minor Date: 1/17/92
R. G. Minor
Chemical Specialist

Approved By: J. J. Murphy Date: 1/17/92
J. J. Murphy, Ph.D.
Manager, Metabolism/Residue Methodology

D. R. Flint Date: 1-17-92
D. R. Flint, Ph.D.
Manager, Environmental Research

Inquiries

Inquiries should be directed to:

J. J. Murphy, Ph.D.
Miles Inc.
Agriculture Division
Miles Research Park
17745 South Metcalf
Stilwell, KS 66085

Telephone: (913) 897-9330

Revisions

Date _____ Revision _____

Table of Contents

	<u>Page No.</u>
Title Page.....	1
Statement of Confidentiality.....	2
Certification of Good Laboratory Practice.....	3
Certification of Authenticity.....	4
Revisions.....	5
Table of Contents.....	6
1.0 Summary.....	8
2.0 Introduction.....	8
3.0 Experimental.....	9
3.1 Materials.....	9
3.1.1 Apparatus.....	9
3.1.2 Reagents/Chemicals/Supplies.....	10
3.1.3 Standard Solutions Required.....	10
3.2 Method.....	11
3.2.1 Calibrations.....	11
3.2.1.1 GPC Elution Profile Determination.....	11
3.2.1.2 Silica Gel Column Profile Determination.....	11
3.2.1.3 SPS Hplc Column Profile Determination.....	12
3.2.2 Extraction.....	13
3.2.2.1 Grain (green foliage, kernels, straw); Peanut (green foliage, kernels, hay).....	13
3.2.2.2 Peanut (shells).....	14
3.2.2.3 Plant Material with High Water Content.....	15
3.2.2.4 Must and Wine.....	15
3.2.2.5 Soil.....	15
3.2.2.6 Water.....	16
3.2.2.7 Refined and Crude Peanut Oil.....	17
3.2.2.8 Peanut Soapstock.....	17
3.2.3 Gel Permeation Chromatography (gpc).....	18
3.2.4 Silica Gel Column Chromatography.....	18
3.2.5 C-18 Bond Elut® Cleanup.....	18
3.2.6 SPS Hplc Column Cleanup.....	19
3.2.7 Gas Chromatographic Analysis.....	20
3.2.7.1 Standard Procedure.....	20
3.2.7.2 Confirmatory Procedure.....	23
4.0 Conclusion.....	23
5.0 Bibliography.....	24
Tables	
1. Final volumes for sample prior to gas chromatographic analysis.....	25

Table of Contents (con't)

	<u>Page No.</u>
Addenda	
1. Extraction efficiency of the analytical residue method for tebuconazole in crops (Miles Ag Report No. 101342; Study No. FR111606).....	26
2. A competitor product interference study for the analytical residue method for tebuconazole residues in plants (Miles Ag Report Nos. 95680 and 95680-1; Study Nos. FR141601 and FR141603).....	27
3. The independent laboratory method validation for the analysis of tebuconazole (FOLICUR®) in crops and processed products (Miles Ag Report No. 101991; Project No. A012.003).....	28
4. Confirmatory procedure for tebuconazole in various peanut matrices.....	29

Gas Chromatographic Method for Determination of Residues of Tebuconazole in Crops, Processed Products, Soil and Water

1.0 Summary

The analytical method for determining the residue of tebuconazole [FOLICUR®, HWG 1608, α -[2-(4-chlorophenyl)ethyl]- α -(1,1-dimethylethyl)-1H-1,2,4,-triazole-1-ethanol] in plant material, soil and water (Miles Ag Report, No. 94295) has been revised based on suggestions following analysis of field-treated peanut meats and hay and minor modifications which were needed during an independent laboratory validation of the method. These revisions involve extraction, cleanup and gas chromatography procedures. In addition, during the analysis of tebuconazole residue in processed peanut products, different extraction procedures were needed. The extraction procedures developed for peanut processed products (refined and crude oil, soapstock) are included in this revised analytical method.

Extraction of residue of tebuconazole from grain (green foliage, kernels, and straw) and peanuts (foliage, hay, and kernels) involves an initial homogenization of the samples in acetone/water (3:1). In the case of peanut shells, the shells are not homogenized in acetone/water, but are first finely ground and then subjected to an overnight reflux in acetone/water (3:1). Plant material containing a high water content or aqueous samples (i.e. wine) are extracted with acetone only. The resultant acetone/water extract is saturated with sodium chloride, and the tebuconazole residue is partitioned into dichloromethane.

Extraction of the tebuconazole residue from refined and crude peanut oil involves extraction with hexane followed by partitioning the residue of tebuconazole into acetonitrile. Soapstock is extracted with ethyl acetate. The ethyl acetate is partitioned sequentially against 1N aqueous HCl and water. Residue of tebuconazole partitioning into ethyl acetate are further purified using a hexane/acetonitrile cleanup.

With the exception of peanut shells, all plant extracts are purified using gel permeation chromatography and silica gel column chromatography. Peanut shell extracts are subjected to an acetone precipitation procedure. With the exception of peanut oil, extracts are further purified using either an SPS hplc column or a C-18 Bond Elut® column prior to gas chromatography with a nitrogen/phosphorus detector.

The residue of tebuconazole is extracted from water samples using dichloromethane and from soil by refluxing with methanol/water (4:1). Water and soil samples are purified using gel permeation chromatography. In the case of extracts of soil and water, the silica gel column cleanup can be eliminated.

2.0 Introduction

The analytical residue method devised by Maasfeld¹ (Miles Ag Report No. S4295) for the gas chromatographic determination of tebuconazole [FOLICUR®, HWG 1608, α -[2-(4-chlorophenyl)ethyl]- α -(1,1-dimethylethyl)-1H-1,2,4,-triazole-1-ethanol] in plant material, soil and water has been revised and rewritten.

During the analysis of tebuconazole residue in crops^{2, 3} and during an independent laboratory validation (PR Notice 88-5) of the method⁴, minor modifications were suggested to both the cleanup and gas chromatography procedures.

In a study using aged radioactive residues from [¹⁴C] tebuconazole metabolism studies to determine the extraction efficiency of the procedures described in Miles Ag Report No. 94295, the extraction of [¹⁴C] tebuconazole was found to be adequate from wheat straw, peanut forage, and peanut kernels but was marginal from peanut shells.⁵ An overnight reflux in acetone/water (3:1) was found to improve the extraction of aged [¹⁴C] tebuconazole residue from peanut shells. The residue method for peanut shells which includes an overnight reflux has been validated.⁶

Analysis of tebuconazole residue in processed peanut products has also been conducted using the analytical method described in Miles Ag Report No. 94295. Modifications of the crop method which were used to analyze for tebuconazole residue in peanut oil involved adding a hexane/acetonitrile partitioning followed by gel permeation chromatography and silica gel column chromatography.⁶ Modifications of the crop method when used to analyze for tebuconazole residue in soapstock involved extracting with ethyl acetate, partitioning with hexane/acetonitrile followed by gel permeation chromatography and silica gel column chromatography, and using an SPS hplc column for final cleanup.⁶ Earlier residue analyses of soapstock were performed using either a Lobar low pressure reverse phase column or a C-18 Bond Elut® column with adequate recovery.^{7, 8} The current residue method for soapstock was validated when using the SPS hplc column for final cleanup.⁶

All of the modifications referred to above have been included in this revised method.

3.0 Experimental

3.1 Materials

3.1.1 Apparatus

Beaker, 1-liter metal

Chromatography column: 19 mm id, length 340 mm, elongated discharge with Teflon cock or equivalent

Gas chromatograph with thermionic N/P detector, Varian 3700 or HP 5890 or equivalent

Gel permeation chromatograph GPC Autoprep 1002 (Analytical Biochemistry Laboratories, Inc., Columbia, MO) or equivalent, equipped with a chromatography column (25 mm id, length 600 mm) packed with 60 g of Bio Beads® S-X 3, 200-400 mesh, pretreated in cyclohexane/ethyl acetate 1:1 (v/v) to allow for swelling. Elution rate: 5.0 ml/min. (For each new gel column or after approx. 500 samples have been run, the elution profile should be determined, see 3.2.1.1)

Glass powder funnel, 100 mm diameter

Graduated cylinders, 100 ml, 250 ml, 500 ml

Graduated pipets, 1 ml, 3 ml, 5 ml, 10 ml

Heating mantels, for 1000 ml round-bottomed flasks, or equivalent

High rpm blender, Polytron (KINEMATICA, Kriens, Switzerland) or equivalent

Injection syringe with Luer-Lock connection, 10 ml

Injection syringe, 10 μ l
 Integrator, Spectra Physics 4100 or LAS on HP 1000 A900 or equivalent
 Magnetic stirrer/hot plate or equivalent.
 N-Evap analytical evaporator (Organomation Association, Inc., South Berlin, MA) or equivalent.
 Porcelain Buchner funnel, 110 mm id
 Round-bottomed flasks with ground-glass joints, 25 ml, 100 ml, 125 ml, 250 ml, 500 ml, 1000 ml
 Reflux condenser
 Rotary vacuum evaporator with water bath, bath temperature 30°C
 Separatory funnels with ground-glass stopper, 500 ml, 1000 ml
 Test tubes with ground-glass joints, 10 ml
 Vacuum filter flasks, 500 ml, 1000 ml
 Volumetric flasks with ground-glass joints, 25 ml, 50 ml, 100 ml

3.1.2 Reagents/Chemicals/Supplies

Acrodisc® CR PTFE, non-sterile, 0.45 μ m and 1.0 μ m pore size, No. 4219 and No. 4226, respectively (Gelman Sciences, product supplied by Baxter Scientific Product, McGaw Park, IL)
 Bond Elut® C-18 octadecyl disposable column, 500 mg capacity (Varian, Harbor City, CA)
 Filter aid, Celite® 545, not acid washed (Fisher Scientific, Fair Lawn, NJ) or equivalent
 Filter paper, No. 41, 90 mm (Whatman, Hillsboro, OR)
 Gases: helium, air, nitrogen, hydrogen.
 Glass microfibre filters, GF/A, 90 mm (Whatman, Hillsboro, OR), for use on soapstock only
 Glass wool
 Hydrochloric acid (concentrated), reagent grade
 Polystyrene gel: Bio Beads® S-X3, 0.037 to 0.074 mm, 200 to 400 mesh, No. 152-2750 (Analytical Biochemistry Laboratories, Inc., Columbia, MO)
 Silica gel 60: 0.063 to 0.200 mm, 70 to 230 mesh (MERCK No. 7734), activated overnight at 130 to 150°C, then 5% (95 g silica gel + 5 g water) deactivated with water. (For each new batch of silica gel, the elution profile must be determined see 3.2.1.2)
 Sodium chloride, reagent grade
 Sodium sulfate (anhydrous), reagent grade
 Solvents, pesticide grade: acetonitrile, acetone, cyclohexane, dichloromethane, ethyl acetate, n-hexane, methanol, toluene, water (hplc grade)
 Solvent mixtures, acetone/water 3:1 (v/v), hexane/ethyl acetate 1:4 (v/v), hexane/ethyl acetate 1:1 (v/v), cyclohexane/ethyl acetate 1:1 (v/v), methanol/water 7:3 (v/v), methanol/water 9:1 (v/v), 1N HCl (v/v)

3.1.3 Standard Solutions Required

Analytical standards of tebuconazole may be obtained from Miles Inc., Agriculture, Metabolism and Residue Methodology, Miles Research Park, 17745 S. Metcalf Ave, Stilwell, Kansas 66085.

Prepare a standard solution of tebuconazole at 250 μ g/ml in ethyl acetate. From this standard solution prepare 0.0625 μ g/ml, 0.125 μ g/ml, 0.25 μ g/ml, 0.625 μ g/ml, 1.25 μ g/ml, 2.50 μ g/ml, 25 μ g/ml solutions of tebuconazole in ethyl acetate.

Store standards under refrigerated conditions (2°C). Stable under refrigerated conditions for at least 1 month.

3.2 Method

3.2.1 Calibrations

3.2.1.1 GPC Elution Profile Determination

1. Process a control sample through the method to the gpc step (3.2.3).
2. Fortify this control with 2 ml of the 250 µg/ml standard of tebuconazole.
3. Evaporate the solvent from the fortified control sample using a rotary vacuum evaporator, and re-dissolve the dry residue in 10 ml of the gpc solvent, cyclohexane/ethyl acetate (1:1, v:v).
4. Fill the 5-ml loop of the gpc, and inject 5 ml of the sample onto the gpc column.
5. Collect twenty-three, 10-ml fractions, and evaporate a 1-ml aliquot from each fraction to dryness.
6. Re-dissolve the dry residue of each fraction in 10 ml of ethyl acetate, and analyze each fraction by gc (Gas Chromatographic Analysis 3.2.6) to determine the amount of tebuconazole in the gpc column eluate by comparing against a 2.5µg/ml tebuconazole standard (this standard represent 100% recovery of tebuconazole from the column).
Note: the amount of tebuconazole in 1 ml is 1/10th of the fraction (1 ml from 10 ml).
7. Establish the "dump" fraction and "collect" fraction ranges by plotting the percent of tebuconazole present in each of the twenty-three, 10-ml fractions. Starting from the last 10-ml fraction which contained tebuconazole and moving toward the first fraction, total the percent recovered until at least 90% has been reached. This set of fractions represents the "collect" fraction. In this manner, the majority of the control sample matrix will be excluded from the "collect" fraction and will be in the "dump" fraction.

3.2.1.2 Silica Gel Column Profile Determination

1. Process a control sample through the method to the silica gel cleanup procedure (3.2.4).
2. Prepare the chromatography column in the following sequence: 10 ml toluene, glass wool plug, 20 g silica gel slurried in toluene (filling height approximately 170 mm), approximately 10 mm sodium sulfate layer. Drain the toluene until the solvent is just above the sodium sulfate.
3. Fortify the processed control sample with 1 ml of the 250 µg/ml standard of tebuconazole.

4. Evaporate the solvent from the fortified control sample using a rotary vacuum evaporator, and re-dissolve the dry residue in 10 ml of toluene.
5. Transfer the dissolved sample to the column with a pipet, and allow the toluene to drain down to the sodium sulfate layer.
6. Rinse the flask with 20 ml of hexane/ethyl acetate (1:1, v/v), and transfer the solvent to the column.
7. Repeat Step 6 three times for a total of 80 ml of rinse solvent. Note: collect all of the eluate up to and including this step in a 250-ml graduated cylinder. This fraction represents the "discard" fraction. Adjust the volume of the eluate to 200 ml and retain for Gas Chromatographic Analysis, 3.2.7.
8. Elute tebuconazole from the column with 200 ml of hexane/ethyl acetate (1:4, v/v). This fraction represents the "collect" fraction. Adjust the volume of the eluate to 200 ml, and retain for Gas Chromatographic Analysis, 3.2.7.
9. Analyze each "collect" and "discard" fraction by gc (Gas Chromatographic Analysis, 3.2.7) to determine the amount of tebuconazole in each fraction. Compare each fraction to a 1.25 $\mu\text{g/ml}$ tebuconazole standard (the 1.25 $\mu\text{g/ml}$ standard represents 100% recovery of tebuconazole from the column).
10. If greater than 10% of the tebuconazole is found in the "discard" fraction, decrease the moisture content of the silica gel to 2.5% water (97.5 g silica gel + 2.5 g water), and repeat the elution profile. If less than 90% of the tebuconazole is found in the "collect" fraction, increase the moisture content of the silica gel to 7% or 10% water (e.g. 90 g silica gel + 10 g water), and repeat the elution profile.

3.2.1.3 SPS Hplc Column Profile Determination

1. Prepare a 100 $\mu\text{g/ml}$ standard of tebuconazole in methanol/water (85:15).
2. Initiate the chromatography process using the following hplc parameters:
 - Instrument: High performance liquid chromatograph (hplc), Beckman Model 338 (Fullerton, CA) or equivalent, capable of solvent gradient elution and equipped with a variable wavelength uv flow-through detector.
 - Mobile phase solvents: Methanol and hplc grade water
 - Solvent flow rate: 2 ml/min
 - UV detector wavelength: 220 nm
 - Solvent program: Start with a linear gradient of 60% methanol to 80% methanol in 20 min, followed by a linear gradient of 80% methanol to 100% methanol in 15 min. Maintain 100% methanol for 15 min. Reequilibrate at 60% methanol for 25 min before making a second injection.

Column: Semi-Permeable Surface (SPS), C8, 5 μ m, 25 cm X 10 mm i.d.
(Regis Chemical Company, Morton Grove, IL)
Pre-Column: Spheri-5, RP-18, 5 μ m, 30 mm X 4.6 mm (Applied
Biosystems, Inc., San Jose, CA)
Sample loop: 1.0 ml

3. Inject 100 μ g of tebuconazole in 1 ml methanol/water (85:15).
4. Determine the retention time for tebuconazole in order to establish when to collect the column eluate. If the retention time for tebuconazole is about 23 min, the collection window should be about 22 to 25.5 min. (The "collection window" should be 1 min before and 2.5 min after the observed elution time for tebuconazole)

3.2.2 Extraction

3.2.2.1 Grain (green foliage, kernels, straw); Peanut (green foliage, kernels, hay)

1. Add 400 ml of acetone/water (3:1) to 50 g of green foliage, or 50 g of kernels, or 25 g of straw or hay, contained in a 1-liter beaker; homogenize the sample for 3 min with a Polytron or equivalent blender.
2. Add approximately 15 g of filter aid to the beaker, and swirl the contents several times.
3. Filter the mixture through a porcelain Buchner funnel under gentle vacuum.
4. Rinse the beaker twice with 100-ml portions of acetone/water (3:1), and filter the rinse through the solids retained in the Buchner funnel under gentle vacuum.
5. Discard the filtered solids.
6. Transfer the filtrate to a 1000-ml separatory funnel.
7. Saturate the filtrate with approximately 40 g of sodium chloride, and partition the mixture with 100 ml of dichloromethane.
8. Allow the layers to separate, and discard the lower aqueous phase.
9. Transfer the organic phase to a 1000-ml boiling flask, and concentrate the organic phase to a volume of 40 ml or less using a rotary vacuum evaporator. *30'*
10. Add 100 ml of dichloromethane to the flask. While slowly swirling the flask, add 150 to 200 g of sodium sulfate to the flask. *dichloromethane*
11. Immediately filter the sample through a glass funnel containing a glass wool plug and approximately 3 cm of sodium sulfate into a 500-ml round-bottomed flask.

12. Rinse the 1000-ml boiling flask (Step 10) three times with 50-ml portions of dichloromethane, and pass the rinses through the sodium sulfate into the 500-ml round-bottomed flask in step 11.
13. Evaporate the filtrate plus rinses to dryness on a rotary vacuum evaporator.
14. Proceed to Gel Permeation Chromatography (3.2.3).

3.2.2.2 Peanut (shells)

1. Add 400 ml of acetone/water (3:1) to 25 g of finely ground shell contained in a 1000-ml boiling flask. Reflux the sample overnight (approx. 16 hours). Allow the sample to cool to room temperature prior to proceeding to Step 2.
2. Add approximately 15 g of filter aid (Celite or Hyflo-Super Cel) to the flask, and swirl the contents several times.
3. Vacuum filter the mixture through a Whatman #41 filter paper (Whatman, Hillsboro, OR) supported on a Buchner funnel into a 1000-ml separatory funnel equipped with a ground glass or Teflon stopcock.
4. Rinse the flask twice with 100-ml portions of acetone/H₂O (3:1), and vacuum filter the rinse through the solids retained on the Buchner funnel into the separatory funnel in Step 3.
5. Discard the filtered solids.
6. Saturate the filtrate with approximately 40 g of sodium chloride, and partition the mixture with 100 ml of dichloromethane.
7. Allow the layers to separate, and discard the lower aqueous layer.
8. Transfer the organic phase to a 1000-ml boiling flask, and concentrate the organic phase to dryness using a rotary vacuum evaporator. Use acetonitrile (approx. 150 to 300 ml) to form an azeotrope with any water present.
9. Add about 30 ml of acetone to the flask from Step 8, and sonicate (approx. 1 min) the flask to loosen the dry residues.
10. Slurry 10 g of filter aid (Celite 545®, Fisher Scientific, Fair Lawn, NJ) in acetone, and vacuum filter the slurry through a 5.5 cm Whatman #2 filter paper supported on a Buchner funnel. Discard the acetone.
11. Vacuum filter the sample from Step 9 through the Celite 545® prepared in Step 10 into a clean 500-ml boiling flask.
12. Repeat Step 9 two more times, and vacuum filter the acetone rinses into the 500-ml boiling flask in Step 11.

13. Evaporate the acetone filtrate to dryness using a rotary vacuum evaporator.
14. Dissolve the dry residues from Step 13 in methanol, and transfer the solution to a 13-ml centrifuge tube using methanol.
15. Adjust the volume of the methanol solution to 10.0 ml.
16. Filter the methanol solution through a 1.0 μm Acrodisc® CR PTFE (Gelman Sciences, Ann Arbor, MI) into a 13-ml centrifuge tube.
17. Proceed to SPS Hplc Column Cleanup (3.2.6).

3.2.2.3 Plant Material with High Water Content

1. Add 300 ml of acetone to 100 g of bananas, or 50 g of banana peel, or 100 g of grapes in a 1000-ml beaker; homogenize the sample for approximately 3 min with a blender.
2. Proceed to Step 2 of 3.2.2.1 for further processing of the sample.

3.2.2.4 Must and Wine

Note: Must is first mixed with approximately 15 g of filter aid and vacuum filtered through filter paper contained in a porcelain Buchner funnel to prevent emulsions.

1. Add 200 ml of acetone to 100 g must or wine contained in a 1000-ml separatory funnel.
2. Saturate the water solution with approximately 40 g of sodium chloride.
3. Add 100 ml of dichloromethane to the separatory funnel, and shake the separatory funnel for approximately 3 min.
4. Allow the layers to separate, and discard the lower aqueous phase.
5. Proceed to Step 9 of Section 3.2.2.1 for further processing of the sample.

3.2.2.5 Soil

1. Place 50 g of soil in a 1000-ml boiling flask containing 300 ml of methanol/water (7:3).
2. Attach a condenser to the boiling flask, and heat the flask contents at reflux for 4 hours.
3. Allow the sample to cool to room temperature; filter the sample using a gentle vacuum through filter paper contained in a Buchner funnel containing approximately 15 g of filter aid.

4. Rinse the flask twice with 50-ml portions of methanol/water (7:3), and filter the rinse through the solids retained in the Buchner funnel.
5. Concentrate the filtrate to the aqueous residue (approximately 100 ml) on a rotary vacuum evaporator.
6. Transfer the aqueous residue to a 250-ml separatory funnel.
7. Partition the aqueous residue with 100 ml of dichloromethane.
8. Allow the layers to separate, and drain the dichloromethane through sodium sulfate (glass funnel with glass wool plug and a 3 cm deep layer of sodium sulfate) into a 500-ml boiling flask.
9. Repeat Steps 7 and 8 twice with 50-ml portions of dichloromethane.
10. Rinse the sodium sulfate contained in the glass funnel three times with 25-ml portions of dichloromethane.
11. Evaporate the dichloromethane to dryness on a rotary vacuum evaporator.
12. Proceed to Gel Permeation Chromatography (3.2.3). Note: The silica gel column cleanup is not necessary.

3.2.2.6 Water

Note: For other volumes of water, the volumes of dichloromethane must be adjusted accordingly.

1. Add 200 ml of water to a 500-ml separatory funnel, and partition the water twice with 100-ml portions of dichloromethane.
2. Allow the layers to separate, and drain the dichloromethane through sodium sulfate (glass funnel with glass wool plug and approximately 3 cm deep layer of sodium sulfate) into a 500-ml boiling flask.
3. Repartition the water layer with 50 ml of dichloromethane, and drain the dichloromethane through the sodium sulfate into the 500-ml boiling flask from Step 2.
4. Rinse the sodium sulfate with approximately 25 ml of dichloromethane.
5. Evaporate the dichloromethane to dryness on a rotary vacuum evaporator.
6. Proceed to Gel Permeation Chromatography (3.2.3). Note: The silica gel column cleanup is not necessary.

3.2.2.7 Refined and Crude Peanut Oil

1. Weigh 25 g of peanut oil into a 500-ml beaker.
2. Dissolve the oil in 200 ml of hexane and transfer to a 500-ml separatory funnel with two, 25-ml portions of hexane.
3. Partition the hexane solution with 100 ml of acetonitrile pre-saturated with hexane. Collect the acetonitrile phase in a 500-ml flask.
4. Repeat Step 3. Combine both acetonitrile fractions in the 500-ml flask, and evaporate the acetonitrile to dryness at room temperature using a rotary vacuum evaporator.
5. Proceed to Gel Permeation Chromatography (3.2.3).

3.2.2.8 Peanut Soapstock

1. Weigh 12.5 g of soapstock into a 1-liter beaker.
2. Add 200 ml of ethyl acetate to the beaker, and extract the soapstock using a Polytron for 30 sec.
3. Add 10 to 15 g of Celite® to the extract, and filter the mixture through a Whatman GF/A filter into a 500-ml separatory funnel. Rinse the beaker twice with 20-ml portions of ethyl acetate, and filter the rinse through the filter cake.
4. Partition the ethyl acetate filtrate with 100 ml of 1N HCl. Discard the aqueous layer.
5. Partition the filtrate two times with 200-ml portions of water (pre-saturated with ethyl acetate). Discard the aqueous layer.
6. Drain the ethyl acetate layer into a 500-ml boiling flask through a powder funnel containing a plug of glass wool and filled with sodium sulfate. Rinse the sodium sulfate three times with 20-ml portions of ethyl acetate.
7. Evaporate the sample to dryness using a rotary vacuum evaporator.
8. Transfer the sample to a 1000-ml separatory funnel using two, 25-ml portions of acetonitrile (pre-saturated with hexane) and two, 100-ml portions of hexane.
9. Shake the separatory funnel for 30 sec, and drain the lower acetonitrile phase into a 250-ml boiling flask.
10. Partition the hexane fraction with an additional 50 ml of acetonitrile (pre-saturated with hexane), and drain the acetonitrile into the same 250-ml boiling flask from Step 9.

11. Evaporate the combined acetonitrile fractions to dryness on a rotary vacuum evaporator.
12. Proceed to Gel Permeation Chromatography (3.2.3).

3.2.3 Gel Permeation Chromatography (gpc)

1. Dissolve the dry residue in 10 ml of cyclohexane/ethyl acetate (1:1, v/v).
2. Using a 10-ml glass syringe, inject 7 to 8 ml into a 5-ml sample loop on the gel chromatograph.
3. Based on the elution profile determined in 3.2.1.1, collect tebuconazole from the gpc column in a 125-ml boiling flask, and evaporate the solution to dryness on a rotary vacuum evaporator.
4. Proceed to Silica Gel Column Chromatography (3.2.4). Note: For soil and water samples proceed to Gas Chromatographic Analysis (3.2.7).

3.2.4 Silica Gel Column Chromatography

1. Use the column as prepared in 3.2.1.2.
2. Dissolve the dry residue in 10 ml of toluene, transfer this solution to the column with a pipet, and allow the toluene to drain down to the sodium sulfate layer.
3. Elute the column as determined in 3.2.1.2.
4. Collect the tebuconazole fraction in a 250-ml round-bottomed flask, and evaporate the solution to dryness on a rotary vacuum evaporator.
5. Proceed to C-18 Bond Elute Cleanup (3.2.5). Note: For peanut oil proceed to Gas Chromatographic Analysis (3.2.7), and for soapstock samples proceed to the SPS Hplc Column Cleanup (3.2.6).

3.2.5 C-18 Bond Elut® Cleanup

1. Add 1 ml of methanol to the dry residue, and swirl the flask.
2. Add 9 ml of water to the sample, and swirl the flask.
3. Condition a C-18 Bond Elut® column with approximately 2 ml of methanol followed by approximately 2 ml of water.
4. Add the aqueous sample to the C-18 column using a pipet, and elute the column dropwise. Discard the eluate.
5. Wash the column with 1 ml of 10% methanol in water, and elute the column dropwise. Discard the eluate.
6. Add 10 ml of methanol/water (7:3), and elute the column dropwise. Collect the eluate.

7. Evaporate the methanol/water fraction to dryness on a rotary vacuum evaporator. Add acetonitrile to form an azeotrope with the water.
8. Transfer the dry residue to a 13-ml centrifuge tube with acetone.
9. Evaporate the residue to dryness using a gentle stream of nitrogen.
10. Proceed to Gas Chromatographic Analysis (3.2.6).

3.2.6 SPS Hplc Column Cleanup

1. Establish a "collection window" as described in 3.2.1.3 (SPS Column Profile Determination).
2. Sample preparation.
 - A. Peanut shells
 - (1). Pipet 5.0 ml of the methanol filtrate from Step 16 of 3.2.2.2 into a 13-ml centrifuge tube, and evaporate the methanol filtrate to 1.7 ml using a gentle stream of nitrogen.
 - (2) Add 300 μ l of distilled water to bring the volume in the tube to 2.0 ml, and mix the contents of the tube well and centrifuge if necessary.
 - (3) Reserve the sample for hplc injection (Step 3).
 - B. Soapstock
 - (1) Transfer the dry residues from Step 4 of 3.2.4 to a 5-ml centrifuge tube.
 - (2) Evaporate the sample to the oily residue using a gentle stream of nitrogen.
 - (3) Pipet 1.7 ml of methanol to the sample and mix the contents of the centrifuge tube to dissolve the residue.
 - (4) Pipet 0.3 ml of distilled water to the sample and mix the contents again.
 - (5) Centrifuge the contents of the centrifuge tube for 10 min at approximately 1200 rpm.
 - (6) Reserve the sample for hplc injection (Step 3).
3. Inject 1.2 ml of the sample solution from Step 2 into the 1.0 ml sample loop of the hplc.

4. Collect the eluate during the "collection window" established in Section 3.2.1.3 (SPS Column Profile Determination) in a 125-ml pear-shaped flask.
5. Evaporate the solvent in the 125-ml flask to an aqueous solution using a rotary vacuum evaporator.
6. Add 30 ml of acetonitrile to the flask, and evaporate the solution to dryness using a rotary vacuum evaporator.
7. Dissolve the dry residues in ethyl acetate, and transfer the solution to a 13-ml centrifuge tube.
8. Refer to Table I for the correct sample volume needed for gc analysis.
9. Proceed to Gas Chromatographic Analysis (3.2.7)

3.2.7 Gas Chromatographic Analysis

3.2.7.1 Standard Procedure

A. Instrument Conditions

Primary:

Instrument: Varian 3700

Data Acquisition: Hewlett-Packard 1000 Mini-Computer with Beckman
CALC System Software

Detector: TSD (Nitrogen-Phosphorous)

Column: 6 ft x 0.25 inch x 2 mm id 3% OV-101 on 80/100
Supelcoport

Gases:

Carrier gas: Helium, 12 ml/min

Makeup: Helium, 17 ml/min

Combustion: Hydrogen, 5 ml/min; Air, 125 ml/min

Temperatures:

Injector: 250°C or 280°C

Detector: 330°C

Column program: 220°C isothermal

Alternate:

Instrument: 5890A Hewlett Packard gas chromatograph

Data Acquisition: 3393A Integrator

Detector: N/P (nitrogen/phosphorous)

Column: Fused silica HP-5, 10 m x 0.53 mm id with 2.65 μ m film
thickness

Gases:

Carrier gas: Helium, 5 ml/min

Makeup: Helium, 25 ml/min

Combustion: Hydrogen, 3.5 ml/min; Air, 110 ml/min

Temperatures:

Injector: 280°C

Detector: 300°C

Column program: 60°C for 1 min

60°C to 180°C at 30°C/min

180°C to 240°C at 15°C/min

240°C for 15 min

240°C to 280°C at 10°C/min*

280°C for 5 min*

(*To be used with peanut meat if late eluting peaks are present.)

Alternate column temperature: 220°C

Alternate:

Instrument: Varian 3400 gas chromatograph

Data Acquisition: Perkin-Elmer Nelson System

Detector: TSD (nitrogen/phosphorous)

Column: HP-1, 25 m x 0.32 mm id with 0.17 μ m film thickness

Gases:

Carrier gas: Helium, 2 ml/min

Makeup: Nitrogen, 26 ml/min

Combustion: Hydrogen, 4.5 ml/min; Air, 170 ml/min

Temperatures:

Injector: 250°C

Detector: 300°C

Column program: 180°C for 1 min

180°C to 230°C at 10°C/min

230°C for 15 min

- B. Standard Curve (e.g. peanut oil, weight following gpc = 12.5 g, final volume = 1 ml)
1. Inject 3 to 5 μ l of each tebuconazole standard, 0.0625 μ g/ml, 0.125 μ g/ml, 0.25 μ g/ml, 0.625 μ g/ml, 1.25 μ g/ml and 2.50 μ g/ml prepared in ethyl acetate and in control matrix. The six standard concentrations above represent 0.005 ppm, 0.01 ppm, 0.02 ppm, 0.05 ppm, 0.1 ppm, and 0.2 ppm based on a final weight of 12.5 g following gpc and a final volume of 1 ml.
 2. Plot area/height versus concentration to confirm a linear response of tebuconazole. The 1.25 μ g/ml tebuconazole standard represents a 0.1 ppm standard based on a sample size of 12.5 g and a final volume of 1 ml.
- C. Procedure:
1. Dissolve the dry residue in an appropriate volume of ethyl acetate (see Table 1).
 2. Inject a 5 μ l aliquot of the 0.1 ppm equivalent standard solution of tebuconazole (1.25 μ g/ml), and determine the area of the tebuconazole peak.

3. Inject a 5 μ l aliquot of the sample, and determine the area of any peak at the retention time of tebuconazole.
4. Repeat sequence of standard, sample, standard, sample, etc. for injections.
5. For the purpose of calculating the ppm recovery in a sample, use the average value of the peak area of the two standards analyzed before and after the sample.
6. If the content of tebuconazole is greater than the response for the highest linearity value, dilute the sample appropriately with ethyl acetate to be within the linear range for tebuconazole, and re-inject the diluted sample.

D. Calculations

Example calculation for determining residues in peanut oil:

Extracts are diluted to a final volume of 1.0 ml and quantitated versus a 1.25 μ g/ml tebuconazole standard. The 1.25 μ g/ml tebuconazole standard is equivalent to a 0.1 ppm sample residue as shown below.

$$1.25 \mu\text{g/ml} \times \frac{\text{(Sample Volume)} \quad \text{(Pre-GPC Vol)}}{\text{(Sample Wt)} \quad \text{(GPC Aliquot)}} = 0.1 \mu\text{g/g (0.1 ppm)}$$

$$1.25 \mu\text{g/ml} \times \frac{1.0 \text{ ml}}{25 \text{ g}} \times \frac{10 \text{ ml}}{5 \text{ ml}} = 0.1 \mu\text{g/g (0.1 ppm)}$$

Example calculation for determining residues in peanut shell:

Extracts are diluted to a final volume of 0.5 ml and quantitated versus a 1.25 μ g/ml tebuconazole standard. The 1.25 μ g/ml tebuconazole standard is equivalent to a 0.1 ppm sample residue as shown below:

$$1.25 \mu\text{g/ml} \times \frac{\text{(Sample Volume)} \quad \text{(Pre-Filtration)} \quad \text{(Pre-Hplc Vol)}}{\text{(Sample Wt)} \quad \text{(Filtrate Aliquot)} \quad \text{(Hplc Aliquot)}} = 0.1 \mu\text{g/g (0.1ppm)}$$

$$1.25 \mu\text{g/ml} \times \frac{0.5 \text{ ml}}{25\text{g}} \times \frac{10 \text{ ml}}{5 \text{ ml}} \times \frac{2}{1} = 0.1 \mu\text{g/g (0.1ppm)}$$

Example calculation for determining residues in peanut kernels:

Extracts are diluted to a final volume of 2.0 ml and quantitated versus a 1.25 μ g/ml tebuconazole standard. The 1.25 μ g/ml tebuconazole standard is equivalent to a 0.1 ppm sample residue as shown below:

$$1.25 \mu\text{g/ml} \times \frac{\text{(Sample Volume)} \quad \text{(Pre-Filtration)}}{\text{(Sample Wt)} \quad \text{(Filtrate Aliquot)}} = 0.1 \mu\text{g/g (0.1ppm)}$$

$$1.25 \mu\text{g/ml} \times \frac{2.0 \text{ ml}}{50 \text{ g}} \times \frac{10 \text{ ml}}{5 \text{ ml}} = 0.1 \mu\text{g/g (0.1ppm)}$$

Residue levels in sample extracts are calculated by the following equation.

$$\frac{\text{Peak Response (Sample)}}{\text{Avg Peak Response (Standard)}} \times \text{Std Conc (0.1 ppm)} \times \text{Dilution Factor} = \text{ppm in sample}$$

3.2.7.2 Confirmatory Procedure

A. Instrument Conditions

Instrument: Varian 6000

Data Acquisition: Nelson Data Integration System (Model 2600 Chromatography Software, Rev 4.1, Nelson Analytical)

Detector: TSD (Nitrogen-Phosphorous)

Column: Rt.-200 15 m x 0.53 mm with a 1.0 μm trifluoropropylmethyl film thickness (Restek Corporation, Bellefonte, PA)

Gases:

Carrier gas: Nitrogen, 3 ml/min

Makeup: Nitrogen, 22 ml/min

Combustion: Hydrogen, 4.6 ml/min; Air, 170 ml/min

Temperatures:

Injector: 300°C

Detector: 300°C

Column: 210°C isothermal

B. Standard Curve - see Standard Procedure (3.2.7.1)

C. Procedure - see Standard Procedure (3.2.7.1)

D. Calculations - see Standard Procedure (3.2.7.1)

4.0 Conclusion

The analytical residue method (Miles Ag Report No. 94295) for determining residues of tebuconazole in plant material, soil and water has been rewritten to include modifications in extraction procedures, cleanup procedures, and gas chromatography analysis. In addition, extraction procedures for peanut processed products (refined and crude oil, soapstock) have also been included in this revised method.

5.0 Bibliography

1. Maasfeld, W. 1986. Method for gas chromatographic determination of residues of the fungicide FOLICUR (HWG 1608) in plant material, soil and water. Miles Ag Report No. 94295. (EPA MRID 40700963)
2. Williams, B. 1988. FOLICUR - Magnitude of the residue on peanuts 3.6F. Miles Ag Report No. 99129. (EPA MRID 51263316)
3. Koch, D. A. and B. B. Williams. 1988. FOLICUR (1.2 EC and 45 DF) - Magnitude of the residues on peanuts. Miles Ag Report No. 96728. (EPA MRID 40995936)
4. Erhardt-Zabik, S. and L. O. Ruzo. 1989. Validation of Miles Method No. 94295 - method for gas chromatographic determination of residues of the fungicide FOLICUR (HWG 1608) in plant materials, soil and water. Miles Ag Report No. 99624. (EPA MRID 41263319)
5. Minor, R. G. and P. L. Freeseaman. 1991. Extraction efficiency of the analytical residue method for tebuconazole in plants. Miles Ag Report No. 101342.
6. Bajzik, M. 1992. The independent laboratory method validation for the analysis of tebuconazole (FOLICUR®) in crops and processed products. Miles Ag Report No. 101991.
7. Koch, D. A. and B. B. Williams. 1989. FOLICUR - Magnitude of the residue on peanut and peanut processed products. Miles Ag Report No. 98541. (EPA MRID 41068501)
8. Leslie, W. L. 1988. FOLICUR - Magnitude of the residue on peanut processed products. Miles Ag Report No. 98308. (EPA MRID 40995945)

Table 1. Final volumes for sample prior to gas chromatographic analysis.
(Samples are compared to a 1.25 $\mu\text{g}/\text{ml}$ tebuconazole standard)

<u>Matrix</u>	<u>Sample Wt. (g)</u>	<u>Post-GPC Sample Wt. (g)</u>	<u>Post-Hplc Sample Wt. (g)</u>	<u>Final Vol. (ml)</u>
Wheat				
Straw	25	12.5		1 ¹
Green Forage	50	25		2 ¹
Kernels	50	25		2 ¹
Peanuts				
Hay	25	12.5		1 ¹
Green Forage	50	25		2 ¹
Kernels	50	25		2 ¹
Peanut shell	25	-	6.25	0.5 ¹
Peanut oil	25	12.5		1.0 ¹
Soapstock	12.5	6.25	3.125	0.25 ¹
Bananas				
Bananas	100	50		4 ¹
Banana Peels	50	25		2 ¹
Grapes				
Grapes	100	50		4 ¹
Others				
Soil	50	25		2 ¹
Wine/Must	100	50		0.4 ²
Water/Leachate	200	100		0.8 ²

¹Based on the sample weight and final volume of the sample the 1.25 $\mu\text{g}/\text{ml}$ standard represents a 0.1 ppm standard

²Based on the sample weight and final volume of the sample the 1.25 $\mu\text{g}/\text{ml}$ standard represents a 0.01 ppm standard

Addendum 1. Extraction efficiency of the analytical residue method for tebuconazole in crops (Miles Ag Report No. 101342; Study No. FR111606).

Introduction

The extraction efficiency of the analytical residue method for detecting tebuconazole in crops was evaluated with matrices containing aged radioactive residues from the [¹⁴C] tebuconazole peanut and the [¹⁴C] tebuconazole wheat metabolism studies.

Results

The extraction efficiency of the analytical residue method was found to be adequate (80% to 107%) for aged radioactive residues in peanut forage, peanut kernels, and wheat straw. A slight modification in the analytical residue method was found to be necessary to increase the extraction efficiency of [¹⁴C] tebuconazole residue from peanut shells. An overnight reflux in the recommended extraction solvent (acetone/water) successfully extracted the aged radioactive residue of tebuconazole from peanut shells.

<u>Matrix</u>	<u>Extraction Efficiency (%)</u>
Peanut	
forage	80, 107
shells	62, 63, 103 ¹
kernel	98
Wheat	
straw	96

¹An overnight reflux in acetone/water (4:1) was substituted for the blending with acetone/water (4:1).

Conclusion

The tebuconazole analytical residue method proved to be generally efficient in extracting aged [¹⁴C] tebuconazole residue from peanut and wheat matrices. The extraction efficiency from peanut shells was improved by substituting a reflux for the solvent blending. Based on these findings, the residue method was modified to include a reflux extraction of the peanut shells.

Addendum 2. A competitor product interference study for the analytical residue method for tebuconazole residues in plants (Miles Ag Report Nos. 95680 and 95680-1; Study Nos. FR141601 and FR141603)

Introduction

As of January, 1987, 94 nitrogen and/or phosphorous containing compounds had a registered tolerance on wheat, grapes, barley, peanuts, or seed grasses as listed in the Pesticide Chemical News Guide (Food Chemical News, Inc., Duggan and Duggan Editors, Washington D. C.). An additional study was conducted in September, 1991 to investigate the 25 compounds which have been granted a registered tolerance since January, 1987.

Results

Of the 25 compounds tested in this study, only nitralin gave a response near the retention time of tebuconazole using the standard gas chromatographic procedure. When nitralin was reanalyzed using the confirmatory gas chromatographic procedure, nitralin was shown not to interfere with the analysis of tebuconazole.

Conclusion

As of September, 1991, 121 nitrogen and/or phosphorous containing compounds had registered tolerances in wheat, grapes, barley, peanuts or seed grasses as listed in the "The Pesticide Chemical News Guide". None of the compounds tested interfered with the analysis of tebuconazole under the conditions described in the analytical residue method for crops. Two compounds were not available for testing.

Addendum 3. The independent laboratory method validation for the analysis of tebuconazole (FOLICUR®) in crops and processed products (Miles Ag Report No. 101991; Project No. A012.003).

Introduction

Huntingdon Analytical Services located in Middleport, New York was selected to validate the analytical residue procedure for the determination of tebuconazole residues in peanut shells, peanut oil, and soapstock according to the provisions outlined in PR Notice 88-5.

The study was conducted from September, 1991 through December, 1991.

Results

Peanut shells, peanut oil and soapstock were extracted by the procedures described in the tebuconazole analytical residue method (Miles Ag Report No. 101341). Control values were <0.1 ppm for peanut shells and <0.05 ppm for peanut oil. Control values for soapstock were <0.01 ppm.

A summary of the results of the independent laboratory validation is shown below:

<u>Matrix</u>	<u>Fortification Level</u>	<u>Recovery</u>
Peanut Shell	2 ppm (1X)	88, 94 (91) ¹
	10 ppm (5X)	102, 112 (107) ¹
Peanut Oil	0.1 ppm (1X)	74, 91 (82) ¹
	0.5 ppm (5X)	87, 102 (94) ¹
Soapstock	0.3 ppm (1X)	70, 76 (73) ¹
	1.5 ppm (5X)	98, 116 (107) ¹

¹The value in parenthesis is the average recovery value

Conclusion

Huntingdon Analytical Services successfully performed the independent laboratory validation of the analytical residue method for the determination of tebuconazole residues in peanut shells, peanut oil, and soapstock.

Addendum 4. Confirmatory procedure for tebuconazole in various peanut matrices.

Introduction

A confirmatory residue procedure was developed for the determination of tebuconazole (FOLICUR) residues in crops, processed products, soil and water (Miles Ag Report No. 101341).

Experimental

Extracts of peanut meat, oil, meal, and soapstock from the tebuconazole peanut processing study (Miles Ag Report No. 101343; Study No. FR19PE03), and peanut dry hay and peanut shells from a residue field trial (Miles Ag Report No. 101344; Study No. FR19PE02)) were confirmed using a Varian 6000 gas chromatograph equipped with an N/P detector and a Restek Rt_x-200 capillary column.

Instrumental conditions are shown below:

Instrument:

Varian 6000 gc equipped with a N/P detector (TSD) and a Nelson Data Integration System (Model 2600 Chromatography Software, Rev 4.1, Nelson Analytical)

Confirmatory Column:

Restek Corporation (Bellefonte, PA)
Rt_x-200 (Crossbonded[®] Trifluoropropylmethyl)
15_xm x 0.53 mm x 1.0 μ film thickness (Miles Column No. 52)

Carrier Gas: Nitrogen
Column Flow: 3 ml/min
Makeup Flow: 22 ml/min

Detector Gases: Air (170 ml/min), Hydrogen (4.6 ml/min)

Temperatures:

Detector: 300°C
Injector: 300°C
Column: 210°C isothermal

Injection Volume: 3μl

Addendum 4 (cont.).

Results

Tebuconazole was shown to have a linear response from 0.75 ng to 6 ng in solvent; in the presence of peanut shell matrix the response was linear from 1.875 ng to 18.75 ng. Recoveries of tebuconazole were confirmed, as shown below, in peanut meat, oil, meal, soapstock, dry hay and shells. Raw data is contained in Miles Notebook No. 91-R-332. Recoveries obtained initially (Miles Ag Report Nos. 101343 and 101344) using the primary residue method are also listed below.

<u>Matrix</u>	<u>Fortification Level</u>	<u>Percent Recovery</u>	
		<u>Confirmatory Method</u>	<u>Primary Method</u>
Peanut			
meat	0.05	108	118
oil (refined)	0.05	94	80
oil (crude)	0.05	82	90
meal	0.05	98	86
soapstock	0.05	86	-
dry hay	5.00	82	98
shells	0.10	101, 82	80, 109

Conclusion

Residue of tebuconazole in crops and processed products can be confirmed using a Varian 6000 gas chromatograph equipped with a N/P detector (TSD) and a Restek Rt_x-200 capillary column.