


US EPA ARCHIVE DOCUMENT



SDMS DocID 000204598

CONTAMINANT DISTRIBUTION IN WHOLE BODY, FILLET AND OFFAL OF FISH AND HUMAN HEALTH EXPOSURE ASSESSMENT OF FISH CONSUMPTION – A PRELIMINARY CASE STUDY OF THE CENTREDALE MANOR RESTORATION PROJECT SUPERFUND SITE

Chau H. Vu¹, Deirdre T. Dahlen², and Karen L. Tracy²

1 U.S. Environmental Protection Agency, Region I, One Congress Street, Boston, MA 02114
2 Battelle Memorial Institute, 397 Washington Street, Duxbury, MA 02332

Centredale
17.07
204598

Introduction

As part of the Remedial Investigation and Feasibility Study (RI/FS) conducted for the Centredale Manor Restoration Project Superfund site in North Providence, Rhode Island by the U.S. Environmental Protection Agency (USEPA) Region I and the U.S. Army Corps of Engineers (USACE), a Baseline Human Health Risk Assessment (BHHRA) and a Baseline Ecological Risk Assessment (BERA) are being developed. The objectives of the BHHRA in the Remedial Investigation are to:

- evaluate the human health risks associated with contaminants at the Centredale site via exposure pathways such as consumption of fish caught from portions of the Woonasquatucket River at the site, incidental ingestion of and dermal contact with the river bank surface soil, surface water and sediment at the site, and
- develop the preliminary remedial goals for the contaminated media.

A BHHRA would consist of four steps: hazard identification, exposure assessment, toxicity assessment and risk characterization¹. This paper focuses on the first two steps of the BHHRA: hazard identification and exposure assessment. The next two steps of the BHHRA, toxicity assessment and risk characterization, would be future EPA work and papers.

The hazard identification or data collection and evaluation process has found elevated levels of dioxin (specifically the most toxic dioxin congener 2,3,7,8-tetrachlorodibenzo-p-dioxin or 2,3,7,8-TCDD); 1,2,4,5,7,8-hexachloroxanthene (HCX) and polychlorinated biphenyls (PCB Aroclors) at the site in soil, sediment, groundwater, surface water and biota. Besides these contaminants that were found at elevated levels at the site, a full suite of other chemicals, including volatile organic compounds (VOCs), semivolatile organic compounds (SVOCs) and inorganics have also been analyzed for samples collected at the site. This paper will include a discussion of data evaluation methods, data summaries of the distribution of the main contaminants (2,3,7,8-TCDD, HCX and PCBs) in sediment and fish at four different exposure areas of the site (Allendale, Lyman Mill, Manton and Dyerville) and two reference areas (Greystone Mill and Assapumpset) and of the correlation between contaminant levels in sediment and in fish².

The exposure assessment process presents the pathways by which the human population may be potentially exposed to the main contaminants at the site for current and future scenarios. Contaminated wastes directly released into the soil and into the river were believed to result in high levels of contaminants in soil, sediment, surface water and biota. Through the release

mechanism of infiltration, leaching and direct release to soil, sediment and surface water, it is believed that the contaminants are transported by erosion, runoff or advection and discharge to the contaminated media of bank soil, sediment and surface water. Contaminants can also be taken up from the sediment and surface water and bioaccumulated in biota. The main receptors at Centredale, i.e., residents, visiting anglers, and commercial/industrial workers, can be exposed to contaminants in these media via biota consumption, ingestion of or dermal contact with bank soil, sediment and surface water. This paper will focus only on the fish consumption pathway from available sediment and fish data collected at the site.


Methods and Materials

Samples for characterization include surface sediments (0-1 foot), American eel (*Anguilla rostrata*), white sucker (*Catostomus commerson*), and largemouth bass (*Micropterus salmoides*)³. American eel and white sucker were analyzed as whole body. Largemouth bass were analyzed as separate fillet (skin-on) and offal tissue samples; with the reconstituted whole body (RWB) concentration calculated based on the proportional sum of the fillet and offal data. Samples were analyzed by high-resolution gas chromatography/high resolution mass spectrometry (HRGC/HRMS) for dioxins, furans, and HCHX following U.S. EPA Method 1613 (Revision B)⁴. Samples were also analyzed by gas chromatography/electron capture detection (GC/ECD) for PCB, as Aroclor, following methods comparable to U.S. EPA Method 8082⁵.

The distribution of contaminants in sediment and in fish at the site was evaluated by comparing either mean (sediment) or individual (fish) concentrations at each of the exposure areas. Because of the large datasets (e.g., over 100 samples analyzed for Allendale), sediment data within a sampling location were averaged. However, fewer fish data were available (10 or fewer samples, by species and location), thereby allowing for comparison on an individual sample basis. The correspondence between contaminants in sediment and fish was evaluated by linear regression using normalized, mean concentrations of contaminants in sediment and fish, by sampling area and species. Sediment data were normalized to the fraction of Total Organic Carbon (TOC) and then averaged by sampling location. Similarly, fish data were normalized to the fraction of lipid and then averaged by sampling location and species. Wherever data were averaged, a value of one-half the detection limit was used in cases of non-detects.

Results and Discussion

Contaminant concentrations were among the highest, and were by far the most variable, in sediment nearest to the source area, i.e., the Allendale sampling area (Figure 1). The magnitude and variability in contaminant concentrations generally decreased downstream of the source area. Mean concentrations of 2,3,7,8-TCDD at reference locations (Assumpset and Greystone Mill) and areas downstream of Lyman Mill (Manton



and Dyerville) were Mill (Figure 1). Since Allendale, followed downstream of Lyman and Lyman Mill. The area with the highest detection limit and Dyerville

Consistent with sediment data, fish contained higher concentrations downstream of Lyman Mill. The highest concentrations were found in largemouth bass. This might be due to the higher concentrations of 2,3,7,8-TCDD in the order of magnitude of the sediment, suggesting that 2,3,7,8-TCDD and PCBs are lipid-soluble and mainly stored in the offal (offal) instead of the tissues (fillet) (Figure 1). Fish was also measured for concentrations of 2,3,7,8-TCDD in the source area and reference areas, between American eel and white sucker. For largemouth bass, however, 2,3,7,8-TCDD was generally undetectable in the fillet, and when offal, concentrations were above the detection limit.


2,3,7,8-TCDD concentrations approached or exceeded the Tolerance Level, suggesting the need for areas (representative of the Tolerance Level) American eel and largemouth bass consistently exceeded the Tolerance Level at Allendale (Figure 1). A fish consumption advisory for the River in North

Based on the sediment data, largemouth bass sediment (no

liment and surface water, it is
 or advection and discharge to the
 ontaminants can also be taken up
 a. The main receptors at
 strial workers, can be exposed to
 or dermal contact with bank soil,
 sh consumption pathway from


, American eel (*Anguilla*
 h bass (*Micropterus salmoides*)³.
 largemouth bass were analyzed as
 ituted whole body (RWB)
 et and offal data. Samples were
 mass spectrometry
 A Method 1613 (Revision B)⁴.
 ure detection (GC/ECD) for
 thod 8082⁵.

was evaluated by comparing
 of the exposure areas. Because
 e), sediment data within a
 available (10 or fewer samples,
 ndividual sample basis. The
 aluated by linear regression
 nt and fish, by sampling area
 tal Organic Carbon (TOC) and
 malized to the fraction of lipid
 ita were averaged, a value of



and Dyerville) were well below mean concentrations of 2,3,7,8-TCDD at Allendale and Lyman Mill (Figure 1). Similarly, the highest concentration of HCX in sediment was measured at Allendale, followed by Lyman Mill. Concentrations of HCX at the reference areas and areas downstream of Lyman Mill were one or more orders of magnitude lower than those at Allendale and Lyman Mill. However, PCB detection was more complicated than the other contaminants, with the highest detection still at Allendale but also some high levels at Greystone Mill, Lyman Mill and Dyerville.


Consistent with sediment findings, fish collected from Allendale and Lyman Mill areas generally contained higher concentrations of contaminants than fish collected at reference areas and areas downstream of Lyman Mill. Among the fish species analyzed, white sucker generally contained the highest concentrations of 2,3,7,8-TCDD and PCBs compared to American eel and largemouth bass. This might be usual with white sucker being a bottom feeding fish. The concentrations of 2,3,7,8-TCDD and PCBs measured in the offal tissues of largemouth bass were generally one order of magnitude higher than those concentrations measured in the corresponding fillet tissues, suggesting that 2,3,7,8-TCDD and PCBs are lipophilic and mainly stored in the fatty tissues. (offal) instead of the muscle tissues (fillet) (Figure 2). HCX was also measured at higher concentrations downstream of the source area than the reference areas, and was similar between American eel and white sucker. For largemouth bass, however, HCX was generally undetected in the fillet, and when detected in the offal, concentrations were near the detection limit.



2,3,7,8-TCDD concentrations in fish collected at Allendale and Lyman Mill areas frequently approached or exceeded FDA "Do Not Consume" levels of 50 ppt by one order of magnitude, suggesting the receptors at the site not eat the offal parts or whole body of fish from these two areas (representative fish, largemouth bass, shown in Figure 2). Exceedences of the FDA Tolerance Level for PCBs were less widespread, with no observed and limited exceedences for American eel and largemouth bass, respectively. However, PCB concentrations in white sucker consistently exceeded the FDA Tolerance Level at Lyman Mill (10 out of 10) and to a less extent at Allendale (3 out of 10). There is no advisory or tolerance level for HCX available at this time. A fish consumption advisory recommending not eating fish caught from the Woonasquatucket River in North Providence issued by the Rhode Island Department of Health is currently in effect.

Based on the limited data available (n = 5 for American eel; n = 3 for white sucker; and n = 4 for largemouth bass), there is a positive correlation between mean concentrations of 2,3,7,8-TCDD in sediment (normalized to fraction TOC) and fish (normalized to fraction lipid) for largemouth bass

and American eel, whereas the correlation is considerably weaker for white sucker (Figure 3). These findings show that largemouth bass has a relatively stronger correlation between sediment and fish 2,3,7,8-TCDD concentrations than American eel and white sucker. Yet interestingly, the data showed that white sucker contained the highest concentrations of 2,3,7,8-TCDD compared to largemouth bass and American eel. Some potential reasons for this are: 1) the differences in diet, habitat, and lipid contents among the species, 2) the differences in age of the fish sampled, or 3) the potential for chemical disequilibrium in the river. The correlation between sediment and fish HCH concentrations was also fairly good among all species analyzed, except for largemouth bass where HCH was undetected for the most part. Results from the correlation analysis between sediment and fish for PCBs are more convoluted, suggesting that there is no clear point source of PCB contamination, as for 2,3,7,8-TCDD and HCH.



Acknowledgements

The authors would like to thank MACTEC Engineering and Consulting, Inc. for sample collection and data management and Battelle scientists for the dioxin/furan, HCH and PCB analyses.

References

1. U.S. EPA; "Risk Assessment Guidance for Superfund, Volume I, Human Health Evaluation Manual (Part A)", Interim Final (1989), EPA/540/1-89/002. Office of Emergency and Remedial Response.
2. MACTEC, Inc. 2003. *Final Data Evaluation Report, Centredale Manor Restoration Project Site, North Providence, Rhode Island*. Prepared for U.S. Army Corps of Engineers New England District. January 15.
3. Eddy, S. and Underhill, J.C. 1984. *How to Know the Freshwater Fishes*, Third Edition. Wm. C. Brown Company Publishers, Dubuque, Iowa.
4. "U.S. EPA Method 1613: Tetra- through Octa- Chlorinated Dioxins and Furans by Isotope Dilution HRGC/HRMS", Revision B (1994), EPA 821-B94-0059. Office of Water, Engineering and Analysis Division.
5. "U.S. EPA Method 8082: Polychlorinated Biphenyls (PCBs) by Gas Chromatography" (1996), Test Methods for Evaluating Solid Waste. Office of Solid Waste and Emergency Response.

Disclaimer

Although the information in this document has been funded wholly (or in part) by the U.S. EPA, it has not been subjected to Agency review. Therefore, it does not necessarily reflect the views of the Agency and no official endorsement should be inferred.

ACCUMULA (TACHYCINI RESTORA

1 USGS, Upper Mid
2 U.S. Environmental

Introduction

From at least 1921 to
Site, North Providence
1971, the site was us
square feet of contam
tetrachlorodibenzo-

The U.S. Environm
Site. The tree swall
swallows are widely
contamination¹⁻⁵. T
locations of interest
Residues in their tis
biota⁸. Swallows w
are available on dio
North America^{1-5,9}.

Because tree swall
elsewhere. Calcula
because nestlings a
is factored out.

The objective of th
the Centredale Ma

Methods

Fifty-nine swallow
Allendale Mill Po
hereafter called G
Lyman, in 2001.
Woonasquatucket
RI. This is the lo
294172E) is the f
pond upstream of