


US EPA ARCHIVE DOCUMENT



R A C V

R E S P O N S E A C T I O N C O N T R A C T F O R

Remedial, Enforcement Oversight, and
Non-Time Critical Removal Activities at Sites of Release
or Threatened Release of Hazardous Substances in Region V



PREPARED FOR

U.S. Environmental Protection Agency



PREPARED BY

CH2M HILL

Ecology and Environment, Inc.

TN & Associates, Inc.

Tucker, Young, Jackson, Tull, Inc.

**FINAL
ECOLOGICAL RISK ASSESSMENT**

**KRESS CREEK/WEST BRANCH DUPAGE RIVER SITE
SEWAGE TREATMENT PLANT SITE
West Chicago Illinois**

Remedial Investigation / Feasibility Study Oversight

WA No. 216-RSBD-05ZZ/Contract No. 68-W6-0025

May 2004

Contents

1	Introduction	1-1
	1.1 Report Organization.....	1-1
	1.2 Project Background	1-2
	1.2.1 Kress Creek/West Branch of DuPage River Site	1-2
	1.2.2 Sewage Treatment Plant Site	1-2
	1.3 Site History	1-3
2	Overview of the Ecological Risk Assessment Process.....	2-1
3	Screening Level Problem Formulation.....	3-1
	3.1 Ecological Setting of the Kress Creek and Sewage Treatment Plant Sites.....	3-1
	3.2 Terrestrial Communities of the Kress Creek and Sewage Treatment Plant Areas	3-2
	3.2.1 Riparian and Other Wetland Communities	3-2
	3.2.2 Upland Woods.....	3-2
	3.2.3 Meadows or Old Fields	3-3
	3.2.4 Wildlife Species	3-3
	3.3 Aquatic Communities	3-4
	3.3.1 Aquatic Macroinvertebrates	3-5
	3.3.2 Fish Communities	3-5
	3.4 Rare, Threatened, and Endangered Species.....	3-6
	3.5 Summary of Available Analytical Data.....	3-6
	3.5.1 Preliminary Conceptual Site Model	3-7
	3.5.2 Identification of Preliminary Constituents of Potential Concern.....	3-7
	3.5.3 Exposure Pathways and Routes.....	3-12
	3.5.4 Ecological Receptors of Concern.....	3-15
	3.5.5 Screening Assessment Endpoints	3-17
4	Screening Level Exposure Estimate and Risk Characterization	4-1
	4.1 Exposure Point Concentrations	4-1
	4.1.1 Plants.....	4-1
	4.1.2 Soil Invertebrates.....	4-1
	4.1.3 Small Mammals	4-2
	4.2 Dietary Intakes	4-3
	4.2.1 Ingestion Screening Values.....	4-3
	4.3 Screening Level Risk Characterization	4-4
	4.3.1 Kress Creek	4-5
	4.3.2 Sewage Treatment Plant River	4-6
	4.3.3 Sewage Treatment Plant Upland Soils	4-7
5	Uncertainty Assessment	5-1
	5.1 Limiting the Analysis to Constituents of Potential Concern that Exceed Background and Established Benchmarks.....	5-1
	5.2 Use of Established Benchmark Values for Comparison.....	5-1

5.3 Inability to Quantitatively Evaluate All Detected Analytes..... 5-1

5.4 Limiting Evaluation of Potentially Complete Exposure Routes to Ingestion..... 5-2

5.5 Use of Default Value of 1.0 for Bioaccumulation Factor..... 5-2

5.6 Assumptions Regarding Conversion of Literature-Based Toxicity Data into Toxicity Reference Values..... 5-2

5.7 Assumptions Regarding Area Use, Bioavailability, Body Weight, Ingestion Rate, and Other Exposure Factors..... 5-2

5.8 Assumptions Regarding Potential Additive and Synergistic Effects 5-2

5.9 Use of the Lowest Reported Benchmark for Comparison..... 5-3

5.10 Data Limitations 5-3

 5.10.1 Specific Limitations of the RAD-BCG Model 5-3

6 Conclusions 6-1

 6.1 Kress Creek..... 6-1

 6.1.1 Radionuclides..... 6-1

 6.1.2 Chemical Contaminants..... 6-1

 6.2 Sewage Treatment Plant River 6-2

 6.2.1 Radionuclides..... 6-2

 6.2.2 Chemical Contaminants..... 6-2

 6.3 Sewage Treatment Plant Upland 6-2

 6.3.1 Radionuclides..... 6-2

 6.3.2 Chemical Contaminants..... 6-2

 6.4 Discussion..... 6-2

7 References..... 7-1

Tables

3-1 Wildlife Species Potentially Occurring and Habitat Associations

3-2 Macroinvertebrate Inventory Results - WBDR

3-3 Fish Survey Results

3-4 Identification of Constituents of Potential Concern, Process Summary, Sediment/Floodplain Soil - Kress Creek

3-5 Identification of Constituents of Potential Concern, Process Summary, Surface Water- Kress Creek

3-6 Identification of Constituents of Potential Concern, Process Summary, Sediment/Floodplain Soil - STP River

3-7 Identification of Constituents of Potential Concern, Process Summary, Surface Water - STP River

3-8 Identification of Constituents of Potential Concern, Process Summary, STP Upland Soil

3-9 Comparison of Maximum Detections with Background Concentrations, Kress Creek

3-10 Comparison of Maximum Detections with Background Concentrations, STP River

3-11 Comparison of Maximum Detections with Background Concentrations, STP Upland Soil

3-12 Comparison of Maximum Detections with Background Concentrations, Fish Tissues

3-13 Constituents Not Evaluated Quantitatively for Ecological Risk (No Benchmarks)

3-14 Summary of Radiological Parameters

3-15 Distribution Coefficients for Inorganic Constituents Detected in KCK/STP Media

4-1 Bioaccumulative Chemicals List and Log K_{ow} Values

- 4-2 Soil Bioconcentration Factors For Plants, Soil Invertebrates and Small Mammals
- 4-3 Exposure Parameters for Upper Trophic Level Ecological Receptors
- 4-4 Ingestion Screening Values for Mammals
- 4-5 Ingestion Screening Values for Birds
- 4-6 Results of Rad-BCG Screening, KCK Sediment Maximum Concentrations
- 4-7 Results of Rad-BCG Screening, KCK Sediment Mean Concentrations
- 4-8 Comparison of Concentrations of Detected Analytes in KCK Sediment to Ecological Benchmark Values
- 4-9 Comparison of Concentrations of Detected Analytes in KCK Surface Water to Ecological Benchmark Values
- 4-10 SERA Food Web Model for KCK
- 4-11 Results of Rad-BCG Screening, STP River Sediments and Surface Water Maximum Concentrations
- 4-12 Results of Rad-BCG Screening, STP River Sediments and Surface Water Mean Concentrations
- 4-13 Comparison of Concentrations of Detected Analytes in STP River Sediment to Ecological Benchmark Values
- 4-14 Comparison of Concentrations of Detected Analytes in STP River Surface Water to Ecological Benchmark Values
- 4-15 SERA Food Web Model Results for STP River
- 4-16 Results of Rad-BCG Screening, STP Upland Soils Maximum Concentrations
- 4-17 Results of Rad-BCG Screening, STP Upland Soils Mean Concentrations
- 4-18 Comparison of Concentrations of Detected Analytes in STP Upland Surface Soil to Ecological Benchmark Values
- 4-19 SERA Food Web Model Results for STP Upland

Figures

- 1-1 Site Location Map
- 3-1 Wildlife Survey Sampling Locations - Kress Creek/West Branch DuPage River to Warrenville Dam
- 3-2 Wildlife Survey Sampling Locations - West Branch DuPage River from STP to Confluence
- 3-3 Conceptual Site Model for Radionuclides, Kress Creek
- 3-4 Conceptual Site Model for Chemical Contaminants, Kress Creek
- 3-5 Conceptual Site Model for Radionuclides, STP River
- 3-6 Conceptual Site Model for Chemical Contaminants, STP River
- 3-7 Conceptual Site Model for Radionuclides, STP Upland
- 3-8 Conceptual Site Model for Chemical Contaminants, STP Upland

Acronyms and Abbreviations

$\mu\text{g/L}$	micrograms per liter
AEC	Atomic Energy Commission
BAF	bioaccumulation factor
BCF	bioconcentration factor
BCG	biota concentration guide
COPC	constituent of potential concern
CSM	conceptual site model
DDT	dichlorodiphenyl trichloroethane
DOE	U.S. Department of Energy
EE/CA	engineering evaluation/cost analysis
EIS	environmental impact statement
ERA	ecological risk assessment
HQ	hazard quotient
IAEA	International Atomic Energy Agency
IEPA	Illinois Environmental Protection Agency
KCK	Kress Creek
KM	Kerr-McGee
k_{oc}	organic carbon partition coefficient
k_{ow}	octanol-water partition coefficient
LD_{50}	lethal dose at 50% of the test population
LOAEL	lowest-observed adverse effect level
mg/kg	milligrams per kilogram
mg/kg-BW/day	milligrams per kilogram body weight of the receptor per day
mGy/d	milliGrays per day
mGy/h	milliGrays per hour
NCP	National Oil and Hazardous Substances Pollution Contingency Plan
NOAEL	no-observed effect level

ORNL	Oak Ridge National Laboratory
p,p'-DDD	p,p'-dichlorodiphenyl dichloroethane
PAHs	polycyclic aromatic hydrocarbons
PCBs	polychlorinated biphenyls
pCi/g	picoCuries per gram
Ra-226	radium-226
Ra-228	radium-228
RAD-BCG	DOE's RAD-BCG model
REF	Rare Earths Facility
RI/FS	remedial investigation/feasibility study
SERA	screening ecological risk assessment
STP	Sewage Treatment Plant
SVOCs	semivolatile organic chemicals
TRV	toxicity reference value
USEPA	U.S. Environmental Protection Agency
WBDR	West Branch of the DuPage River

SECTION 1

Introduction

This screening ecological risk assessment (SERA) was conducted for the Kerr-McGee Sewage Treatment Plant (STP) and Kress Creek/West Branch of the DuPage River (KCK) Sites, DuPage County, West Chicago, Illinois. It follows methodology outlined in the USEPA's Superfund Risk Assessment Guidance (1997).

The National Contingency Plan (NCP) (Section 300.430 (d)(1)) requires that a risk assessment be performed as part of an Remedial Investigation/Feasibility Study (RI/FS). The primary purpose of the ecological risk assessment (ERA) is to provide risk managers with an understanding of the actual and potential risks to the environment posed by a site and any uncertainties associated with the assessment. This information may be useful in determining whether a current or potential threat to the environment exists that warrants remedial action (USEPA 1990; 1991).

At the conclusion of the SERA, there are four possible decision points:

1. **No further action is warranted.** This decision is appropriate if the SERA indicates that sufficient data are available on which to base a conclusion of no unacceptable risk.
2. **Further evaluation is warranted.** This decision is appropriate if the SERA indicates that there is the potential for unacceptable risks for some pathways, receptors, and chemicals. In this instance, the ERA would progress to the baseline phase of the ERA process.
3. **Further data are required.** This decision is appropriate if the SERA indicates that there are insufficient data on which to base a risk estimate. This decision may also be appropriate if the potential for unacceptable risks is identified following the SERA and additional data to refine these estimates (e.g., additional analytical data, measures of bioavailability, etc.) are needed.
4. **Take remedial action.** This decision may be appropriate for circumstances in which the potential for unacceptable risks was identified following the SERA but these potential risks could best be addressed through remedial action (e.g., presumptive remedy, soil removal) rather than additional study.

Kress Creek (KCK) and the Sewage Treatment Plant (STP) are two of four sites in and around West Chicago, Illinois, that have been contaminated by materials generated and stored on the Kerr-McGee Rare Earths Facility (REF). This report presents the results of the SERA conducted for these two sites in light of the objectives presented above; media data collected in 1993 through 1995, and 1999 through 2001 were used to conduct this analysis. Additionally, the RI Report for the Kress Creek and STP Sites, prepared by BBL (2004) was used for project background information.

1.1 Report Organization

This report is divided into the following sections:

- **Section 1: Introduction.** Describes the purpose and scope of the SERA and outlines the report organization.
- **Section 2: Overview of the Ecological Risk Assessment Process.** Presents a brief discussion of the U.S. Environmental Protection Agency's (USEPA's) ecological risk assessment (ERA) approach.
- **Section 3: Screening Level Problem Formulation.** Describes the ecological setting of the site, including relevant transport pathways, receptors of concern, and the development of the conceptual site model (CSM).
- **Section 4. Screening Level Exposure Estimate and Risk Characterization.** Incorporates all of the qualitative and quantitative statements into one cohesive description of site risks and identifies the constituents of potential concern (COPCs).
- **Section 5. Uncertainty Assessment.** Identifies the sources of uncertainty in the SERA in the context of their potential impacts on the risk conclusions.
- **Section 6: Conclusions.** Presents the conclusions of this SERA.
- **Section 7: References.** Lists all references cited in the report.

Tables and figures are provided at the end of this document in respective sections.

1.2 Project Background

1.2.1 Kress Creek/West Branch of DuPage River Site

The Kress Creek site (KCK), located in DuPage County, Illinois, includes about 1.5 miles of Kress Creek and 5.2 miles of the West Branch DuPage River (WBDR), and contains contaminated sediments, banks, and/or floodplain areas. The site became contaminated by past surface water runoff from the REF that discharged into the creek via a storm sewer outfall located south of Roosevelt Road (Route 38), just east of the Elgin, Joliet, and Eastern railroad tracks. The KCK Site includes the creek from the storm sewer outfall to the creek's confluence with the WBDR, and the WBDR from the confluence to the McDowell Dam. The study area originally ended at the Warrenville Dam, but later was expanded further downstream to the McDowell Dam. See Figure 1-1.

1.2.2 Sewage Treatment Plant Site

The STP Site includes the West Chicago Sewage Treatment Plant (STP Upland), which is owned and operated by the City of West Chicago, and approximately 1.2 miles of the WBDR from the northern boundary of the STP property to the river's confluence with the creek (STP River). See Figure 1-1. The STP upland became contaminated from the use of thorium mill tailings as fill material. Kerr-McGee and the City of West Chicago conducted voluntary cleanup actions at the STP Upland during the mid-1980s (prior to the site's listing on the National Priorities List). The STP River has areas with contaminated sediments, banks and/or floodplains and became contaminated by runoff and erosion from contaminated areas of the STP Upland.

1.3 Site History

Detailed information on site history for the KCK and STP Sites is contained in the RI Report for the Kress Creek and STP Sites (BBL, 2004).

Overview of the Ecological Risk Assessment Process

The USEPA (USEPA 1997) has developed an 8-step process for conducting ERAs as follows:

- **Step 1:** Screening-Level Problem Formulation and Ecological Effects Evaluation
- **Step 2:** Screening-Level Exposure Estimate and Risk Calculation
- **Step 3:** Baseline Risk Assessment Problem Formulation
- **Step 4:** Study Design and Data Quality Objective Process
- **Step 5:** Field Verification of Sampling Design
- **Step 6:** Site Investigation
- **Step 7:** Risk Characterization
- **Step 8:** Risk Management

Steps 1 and 2 together constitute a SERA, the purpose of which is to determine the potential for risks based on conservative assumptions and methodologies. If such risks are possible, the results of the SERA are then used to focus subsequent steps of the ERA process (including the collection of any subsequent data) on the areas, chemicals, media, and receptors with the highest risk potential. Step 3 of the ERA process consists of a refined problem formulation and is the first step of the baseline ecological risk assessment (BERA). In Step 3, risk estimates are recalculated based on refined exposure assumptions, site-specific data, and/or detailed literature review. In Steps 4 through 6 of the process, methodologies for collecting and evaluating the data needed to answer these risk questions (test the hypotheses) are developed and the data are collected. These data are used to derive an estimate of potential risk (with an associated evaluation of the level of uncertainty of the estimate) in Step 7 using a weight-of-the-evidence type of approach relative to the assessment endpoints and risk questions. In Step 8, any identified risks are addressed through a risk management process. Each of these steps is conducted as the results of previous steps warrant. Under certain circumstances (e.g., sufficient data exist following Step 3 to adequately characterize risks), some steps of the process may be bypassed.

The steps reported herein include:

- **Screening Level Problem Formulation:** Summarization of the ecological characteristics of the site as well as background and site characterization data collected during field investigation activities, identification of detected analytes, compilation of existing, media-specific ecological benchmark values, selection of COPCs and receptor species for quantitative analysis in the ERA, selection of endpoints to screen for risk, and the development of a CSM.
- **Screening Level Risk Characterization:** Comparison of measured concentrations for COPCs to established benchmarks to determine the potential for adverse effects to receptor species, including a qualitative discussion of the major sources of uncertainty and conservatism inherent in the evaluation.

Screening Level Problem Formulation

For the screening level problem formulation, a conceptual site model is developed that addresses these five issues:

1. Environmental setting and contaminants known or suspected to exist at the site;
2. Contaminant fate and transport mechanisms that might exist at the site;
3. The mechanisms of ecotoxicity associated with contaminants and likely categories of receptors that could be affected;
4. What complete exposure pathways might exist at the site;
5. Selection of endpoints to screen for ecological risk.

These issues are discussed in the following sections.

3.1 Ecological Setting of the Kress Creek and Sewage Treatment Plant Sites

Information in this section was derived from technical memoranda from CH2M HILL to the USEPA summarizing ecological field activities at the KCK and STP sites (CH2M HILL 1993; 1994; 1995).

The KCK and STP sites lie within the Great Lake and Till Plains sections of the central Lowland Physiographic Province, about 30 miles west of Lake Michigan. This portion of DuPage County is characterized by gently rolling topography, with greater relief near rivers and creeks.

Major land uses and cover types of the KCK and STP areas are varied and interspersed. They range from high-density residential areas to floodplain forest. Portions of the project area lie within or abut the Blackwell Forest Preserve, which contains a mix of wildlife habitat types including forested wetlands, oak-hickory woodlands, and open fields and meadows.

Terrestrial and aquatic community surveys were conducted for the KCK and STP areas, as part of the initial site investigation work completed in 1993 and 1994. For the purpose of the ecological survey work for KCK, the study area at that time was defined as the area extending from the storm sewer outfall to the creek's confluence with the WBDR and from there downstream along the WBDR to the Warrenville Dam. (The KCK Site was later extended downstream to the McDowell Dam.) The study area for STP was defined as the area extending from the STP to the confluence of the WBDR with KCK. Total stream length within the study area for both sites at the time was approximately 4.75 miles.

From within this general study area, sample locations for the terrestrial and aquatic community investigations were selected. Final sampling areas were determined following a site reconnaissance to assess habitat condition, access, and physical conditions of the sites

(Figure 3-1 and Figure 3-2). Detailed information regarding the methods and results of the ecological characterization work are found the Source Characterization and Hydrological Assessment Technical Memoranda (CH2M HILL 1993; 1994; 1995).

3.2 Terrestrial Communities of the Kress Creek and Sewage Treatment Plant Areas

3.2.1 Riparian and Other Wetland Communities

Wetlands are found near Kress Creek and the WBDR. The two general categories of wetlands in the area are riverine and palustrine (CH2M HILL 1994). A riverine wetland includes wetlands and deepwater habitats contained within a channel, except those areas dominated by trees, shrubs, persistent emergents, emergent mosses, or lichens. The palustrine system includes nontidal wetlands dominated by trees, shrubs, persistent emergents, emergent mosses, or lichens. The three classes of palustrine wetlands found in the area include emergent, forested, and wetlands with unconsolidated bottoms (CH2M HILL 1994). The emergent wetland is characterized by erect, rooted, herbaceous hydrophytes. A forested wetland is characterized by woody vegetation that is at least 20 feet tall. Areas classified as unconsolidated bottoms include wetland and deepwater habitats with at least 25 percent cover of particles smaller than stones, and a vegetative cover less than 30 percent (U.S. Department of the Interior 1979).

Cattails and reed canary grass are common herbaceous plants found in the emergent wetlands of the KCK and STP areas. Box elder, elm, willow, green ash, cottonwood, silver maple, and red dogwood are woody species typically found in the forested wetlands. Near the creek and river, the wetlands classified as palustrine with unconsolidated bottoms are areas that have been excavated in the past but are now either permanently flooded or intermittently exposed. Vegetation is generally found around the edges of these areas and includes herbaceous species such as reed canary grass and cattails, plus woody plants including elm, box elder, and willow.

Vegetation found along the creek and river is typical of the wetland vegetation described above in low areas, plus wooded uplands, residential/urban areas, and parkland. The WBDR crosses through the western portion of the Blackwell Forest Preserve in an area of upland oak woods and forested wetland.

3.2.2 Upland Woods

Upland plant communities in the project area include oak woodlands, oak savanna, field (includes mowed parkland, yards, and old field), and agricultural land. Oak woodlands are found in the project area, especially in the Blackwell Forest Preserve along the WBDR. Trees commonly found in these oak woodlands include bur oak, white oak, red oak, shagbark hickory, and bitternut hickory. Small, remnant areas of oak savanna are present at the Blackwell Forest Preserve. Savannas are plant communities in which trees are present, but their density is so low that grasses and other herbaceous vegetation dominate the community. Bur and black oak are the predominant tree species in the oak savanna.

3.2.3 Meadows or Old Fields

Fields along the WBDR include some successional old fields; however, mowed grass in residential and parkland areas predominate. Grassy areas generally extend to the banks of the river or creek. Vegetation in the successional old fields includes grasses, goldenrods, brambles, and tree saplings. Agricultural fields are also found in the area, but are generally not adjacent to the creek or river.

3.2.4 Wildlife Species

A variety of wildlife species may potentially use the KCK and STP areas. Actual use will ultimately depend on the type and quality of wildlife habitat present. Habitat quality is a function not only of the type and distribution of the various plant community types described above, but on other factors such as the proximity to human disturbance. Preliminary information on wildlife occurrence was obtained from sources within DuPage County, including the Fermi National Accelerator Laboratory and the Forest Preserve District of DuPage County. Additional sources of information included the Illinois Natural History Survey and previously prepared Environmental Impact Statements (EISs) for Kerr-McGee's REF. This information was supplemented with the results of actual onsite surveys conducted in 1993 (CH2M HILL 1994).

Table 3-1 lists wildlife species potentially present at the sites. This information is based on wildlife inventory data provided by the Forest Preserve District of DuPage County and the results of field surveys; it was presented in the RI report for the Kerr-McGee Reed-Kepler Park Site (also in West Chicago, Illinois). The District's inventory has been developed through a series of faunal surveys of the various preserves of the county, beginning in 1981.

The Forest Preserve District of DuPage County also categorizes each species by an abundance status, such as abundant, rare, etc. As with general species occurrence, actual abundance within the KCK and STP areas will depend on habitat type and quality. Other sources of information, such as wildlife surveys of the Fermi National Accelerator Laboratory, are considered in the discussions below.

Birds

Seventy-five species of birds may potentially occur within the KCK and STP study areas based on information from the Blackwell Forest Preserve (Table 3-1). Within the nearby preserve, species such as the Mallard (*Anas platyrhynchos*), Canada Goose (*Branta canadensis*), American Robin (*Turdus migratorius*), Common Grackle (*Quiscalus quiscula*), and House Sparrow (*Passer domesticus*) are considered abundant, while many others are considered common. Species such as the Eastern Bluebird (*Sialia sialis*) and Blue-Gray Gnatcatcher (*Poliophtila caerulea*) were considered rare. Of the 75 known species to use the area, 32 were confirmed to be present within the KCK study area and 25 in the STP study area, based on the results of in-field surveys.

A 1988 survey of bird species occurrence at the Fermi National Accelerator Laboratory reported a significantly greater number of bird species (CH2M HILL 1994). Two hundred and twenty-four species were identified at Fermi Laboratory during 1987-1988 survey period, including 17 species on the state endangered list. Although the greater number and diversity of avian species at the Fermi Laboratory is certainly due to the greater extent and number of types of available habitat, results of the survey would suggest a diverse

population of avian species may occur in DuPage County either as breeding residents or migrants. The Forest Preserve District of DuPage County reports a total of 132 resident and 162 migrant bird species for the entire county.

Amphibians and Reptiles

Six species of amphibians and nine species of reptiles are reported for the Blackwell Forest Preserve. Three species, the bullfrog (*Rana catesbeiana*), green frog (*Rana clamitans*), and eastern garter snake (*Thamnophis semifaciatata*), were observed onsite during the 1993 surveys. Many of the other common species, such as American Toad (*Bufo americanus*) and Common Snapping Turtle (*Chelydra serpentina*), could be expected to occur within the KCK and STP study areas. The EIS relating to the Kerr-McGee REF (1982) listed four additional amphibian and reptile species as likely to occur on or near the Kerr-McGee facility. These included the Eastern Mud Turtle (*Terrapene carolina carolina*), the Gopher Tortoise (*Gopherus polyphemus*), the Spring Peeper (*Hyla crucifer*), and the Striped Chorus Frog (*Pseudacris triseriata*). The extent to which these species may or may not be present in the KCK and STP areas is unknown. The Forest Preserve District of DuPage County reports a total of 16 amphibian and 23 reptiles species for the entire county (Table 3-1).

Mammals

Ten mammal species or their sign were observed within the KCK and STP project area, while an additional 17 species of mammals are reported for the Blackwell Forest Preserve. These range from common species such as the Raccoon (*Procyon lotor*) to the rare Least Weasel (*Mustela rixosa*). Many of the more common mammal species were confirmed to be present in the study area. The EIS relating to the Kerr-McGee REF also reported the Deer Mouse (*Peromyscus maniculatus*), the Norway Rat (*Rattus norvegicus*), the Longtail Weasel (*Mustela frenata*), and the Prairie Vole (*Microtus ochrogaster*) as likely to occur in the area. A total of 45 mammal species for DuPage County are reported by the Forest Preserve District (Table 3-1).

3.3 Aquatic Communities

The physical characteristics of the KCK stream channel differ dramatically throughout the study area. This is the result of the extensive channelization and urbanization within its watershed. Upper portions of the creek were found to contain more silt, while the lower portions contained more gravel and cobble in the substrate. The water in the upper reaches appears to carry a much greater silt load. Water quality parameters, which were measured during the 1993 in-field characterization survey, varied, but were found to be within the range for the support of aquatic life.

The physical characteristics of the WBDR (including both STP River and KCK site portions of the River) were also found to vary, with the upper portion of the river containing more gravel, sand, and detritus than the lower portion, which contained more silt and sand as a result of the Warrenville Dam impoundment at the southern end. The water quality parameters measured throughout the WBDR were within the critical limits for the support of aquatic life.

Habitat assessment criteria, which were also evaluated during field surveys of the site, were used to qualitatively assess habitat quality. In general, the habitat within Kress Creek was found to contain many limiting factors. Based on the assessment, the habitat quality generally improved in a downstream direction, providing the best habitat for the support of aquatic life.

3.3.1 Aquatic Macroinvertebrates

The aquatic macroinvertebrate community structure within KCK was found to vary. The upper portions of the creek were dominated by organisms that are more tolerant of silt-laden substrates. Few intolerant species, such as mayflies and caddisflies, were seen in the upper portions. The lower portions of the creek experience a distinct change in community structure with an increase in the mayfly and caddisfly numbers due to the increase in flow velocity and change in substrate material from silt/sand to a more sand/gravel bottom. The community structure in the WBDR showed similarity in overall composition to the lower portions of the creek, but some differences were noted because of a change in stream order and substrate material.

An extensive collection of invertebrates from the WBDR was conducted by the Illinois Environmental Protection Agency (IEPA) and the Illinois Department of Conservation. These collections were made as part of an ecological assessment of the DuPage River Basin (IEPA/WPC/88-010 1988). Two sites, GBK-07 and GBK-05, were sampled on the WBDR just upstream and downstream of the confluence with KCK, respectively. Table 3-2 contains the macroinvertebrate inventory taken during the study. Results from this study also suggest a restrictive environment dominated by organisms that can handle tolerate enrichment, and are common in lotic, erosional environments.

3.3.2 Fish Communities

The fish community of KCK was found to be dominated by non-game species such as carp, sucker, and creek chub. Green sunfish were the only abundant game species present. Some bass and crappie were also collected in the creek, but in few numbers. The fish community structure in the WBDR was also found to be dominated by sucker and green sunfish. Physical anomalies noted in some fish included reddening of caudal fin areas and trematode infestation.

The IEPA (1988) has also conducted fish population assessments in the WBDR as part of their stream classification system. Two of their sampling stations (GBK-07 and GBK-12) were near the WBDR study area. By far, the most prevalent species are carp, minnows, and white suckers (Table 3-3). These results were consistent with the results of project specific fish community surveys conducted at the Kress Creek Site.

Based on Biological Stream Characterization ratings for the streams of Illinois (IEPA/WPC/89-275, 1989), Station GBK-07, upstream of the confluence of KCK on the WBDR, has been designated as Stream Class D, limited aquatic resource. This class has poor biotic resource quality, with the fish community dominated by tolerant forms. The species richness may be notably lower than expected for geographic area, stream size, or available habitat. Station GBK-12, downstream of the KCK confluence, was also designated as Stream Class D. GBK-05, also downstream of the confluence, was designated as Stream Class C,

which is moderate aquatic resource. This class has fair biotic resource quality, and fish consist primarily of bullheads, sunfish, and carp. The topic structure is skewed with increased frequency of omnivores and tolerant species.

3.4 Rare, Threatened, and Endangered Species

State or federally listed rare, threatened, or endangered species can be of particular concern in an ecological assessment due to their population status and sensitivity. At the time of the initial community assessment, the only federally listed threatened and endangered wildlife species known to the general project vicinity was the Indiana bat (*Myotis sodalis*). This species, which is currently listed as endangered, is known to occur in the county. Indiana bats inhabit floodplain and riparian woodlands during spring and summer months and over winter in caves. Nursery roosts are generally located under the shagging bark of dead or dying trees, where females bear usually one young. At the time of the original survey, the Forest Preserve District of DuPage County listed Lyman Woods and Waterfall Glenn Preserve as locations of known occurrence of the Indiana bat since 1980. This species was reportedly mist-netted at Lyman Woods on August 27, 1986.

Additionally, two federally threatened plant species are known to exist in DuPage County, the Eastern Prairie Fringed Orchid (*Platanthera leucophaea*), which occupies wet grassland habitat, and the Prairie Bush Clover (*Lespedeza leptostachya*), which occupies dry to mesic prairies with gravelly soils.

Information on state listed threatened and endangered species of the nearby Blackwell Forest Preserve was provided by preserve personnel. Known sightings include the Yellow Headed Blackbird (*Xanthocephalus xanthocephalus*), the Black Tern (*Chlidonias niger*), the Common Moorhen (*Gallinula chloropus*), and the Black-Crowned Night Heron (*Nycticorax nycticorax*) (W. Lamjsa, personal communication 1992). Countywide, the Forest Preserve District, in their inventory of the flora and fauna of the preserves of DuPage County, also lists three additional state listed species, including one endangered (Great Egret/*Casmerodius albus*), one threatened (Veery/*Catharus fuscescens*), and one watch species (Least Weasel/*Mustela rixosa*).

3.5 Summary of Available Analytical Data

Soil, sediment, and surface water samples were collected at the KCK and STP sites during 1993 through 1995, and 1999 through 2001. Additionally, fish tissues (white sucker and carp) were collected from the creek and the WBDR.

Sample analyses included radionuclides, metals, SVOCs, VOCs, PCBs, and pesticides, although not all media were analyzed for all of these constituents. More detailed information regarding the sampling and analyses performed at the KCK and STP Sites may be found in the RI Report for the Kress Creek and STP Sites (BBL, 2004). All analytical results also are presented in that report.

All positive analyte detections, including those with J qualifiers (i.e., estimated concentrations) were incorporated into this evaluation. Exposure point concentrations were developed using one-half the detection limit for non-detects, where applicable.

It should be noted that, for the purposes of this screening ERA, the data summarized for fish tissue (i.e., metals and radionuclides) were not considered quantitatively except to compare to background constituent concentrations. No ecologically-based benchmark values are available for fish tissue. However, the occurrence of COPCs in fish is indicative of the potential for food transfer and the attendant potential for impacts to higher trophic level organisms. Should a full baseline risk assessment be undertaken, these data could form the basis of dose estimates for piscivorous and omnivorous upper trophic level receptors.

3.5.1 Preliminary Conceptual Site Model

Information on the habitat features at the site and on the fate and transport of the chemicals detected at the site were used to construct CSMs (Figures 3-3 through 3-8). Key components of the CSM include the identification of potential sources of contamination (and identification of COPCs), transport pathways, exposure routes, and receptors. These components are described below.

Sources of Contamination

The waste materials transported from the REF contained a wide range of constituents, including tailings from processed ores, possibly untreated ores, and waste products from other process and manufacturing activities. Numerous sampling and analysis programs were conducted on the original waste materials at the REF. The radiological residuals include thorium, uranium, and their radioactive decay products. Additionally, there may also be natural sources of toxic and bioaccumulative substances in the river system such as weathering and erosion of terrestrial soils, bacterial decomposition of vegetation and animal matter, and long-range transport of substances originating from forest fires or other natural combustion sources.

3.5.2 Identification of Preliminary Constituents of Potential Concern

In order to focus the risk assessment on those constituents that are most likely to cause significant ecological effects, a tiered screen was performed on each medium of concern for the KCK and STP Sites, which considered nutritional status, frequency of detection, comparison to respective background concentration and a comparison with ecological benchmarks. The results of this elimination process are described below and illustrated on Tables 3-4 through 3-8; the risk screens are presented in subsequent sections.

Both chemical and radionuclide contaminants have been detected at all three investigation areas. Radionuclides are defined as contaminants that induce toxicity through the emission of ionizing radiation. Chemical contaminants are those that have toxic effects independent of radiological properties and include metals, SVOCs, VOCs, pesticides and PCBs.

Some chemical contaminants such as uranium possess both chemical and radiological toxicity. However, there are no ecological benchmark values for uranium for the aquatic and terrestrial receptors of concern at the KCK and STP Sites. As a result, the chemical toxicity of uranium was not evaluated in this document. Those chemical toxicity studies that do exist in the open literature are not robust and are not sufficiently representative of site conditions to be applicable for use in this document. It is also expected that, on the population level, the radiological effects of uranium would supercede any potential chemical effects to ecological receptors and, therefore, the radiological benchmarks should be considered adequately protective.

Essential Nutrients

The nutrients calcium, magnesium, potassium and sodium were removed from the constituent lists.

Detection Frequency

A constituent was eliminated if frequency of detection (i.e., the number of positive detections relative to the overall number of analyses) was less than 5 percent.

Background Comparison

Background data for inorganic chemicals and radionuclides were collected from surface soils and sediment from unimpacted areas within KCK and from an adjacent unimpacted area of the WBDR at the STP (performed during the 1993 sediment sampling program conducted by the USEPA). In addition, background concentrations of inorganics in surface water were obtained from these locations (data were unavailable for radionuclide analytes in surface water). Fish tissue was also collected for background comparison. These data sets were combined (i.e., KCK and STP, by medium) and formed the basis for a screen of measured site concentrations of detected analytes to naturally occurring background levels.

An analyte was considered to be not significantly different from background if the maximum of detected values was below the maximum of data from the combined background data set. Tables 3-9 through 3-12 summarize the comparisons of maximum analyte detections to background levels for surface soils, sediment, surface water and fish tissue. This was performed for KCK sediments, STP sediments, STP upland soils, KCK surface water, and STP surface water.

Benchmark Comparisons

A critical step in any risk assessment is the identification of the contaminants that will be included in the quantitative analysis of the potential for adverse effects to receptors. For the purposes of the ERA, the COPC selection process is straightforward and limited in scope.

As described previously in this document, a variety of analytical procedures were performed to characterize the suite of contaminants in surface soils, sediments and surface water, and some constituents were eliminated from further consideration, as described above. From the remaining group of constituents, two types of risk screening procedures were performed to further limit the list of COPCs to those contaminants that are projected to be the most deleterious to ecological receptors. One was performed for radionuclides; the other was performed for the following groups of chemical contaminants; semivolatile organic chemicals (SVOCs), pesticides, polychlorinated biphenyls (PCBs), and inorganics. A discussion of these two methods is provided below.

Radionuclide Screening. Radionuclide concentrations in the media of concern were screened for potential ecological effects using the U.S. Department of Energy's (DOE's) RAD-BCG model (DOE 2002). The rationale of the model is based on several biological principles derived from the literature (IAEA, 1992):

- Aquatic animals are no more sensitive than other organisms; however, because they are poikilothermic animals, temperature can control the time of expression of radiation effects.

- The radiosensitivity of aquatic organisms increases with increasing complexity, that is, as organisms occupy successively higher positions on the phylogenetic scale.
- The radiosensitivity of many aquatic organisms changes with age or, in the case of unhatched eggs, with the stage of development.
- Embryo development in fish and the process of gametogenesis appear to be the most radiosensitive stages of all aquatic organisms tested.
- The radiation-induced mutation rate for aquatic organisms appears to be in between that for fruit flies and mice.
- Appreciable effects in aquatic populations would not be expected at doses lower than 1 rad/d (10 mGy/d); limiting the dose to the maximally exposed individuals to less than 1 rad/d would provide adequate protection of the population.

Additionally, the IAEA (1992) summarized information about the effects of chronic ionizing radiation on terrestrial organisms as follows:

- Reproduction (encompassing the processes from genetic formation through embryonic development) is likely to be the most limiting endpoint in terms of survival of the population.
- Sensitivity to chronic radiation varies markedly among different taxa; certain mammals, birds, reptiles and a few tree species appear to be the most sensitive.
- In the case of invertebrates, indirect responses to radiation-induced changes in vegetation appear to be more critical than direct effects.
- Irradiation at chronic dose rates of 1 rad/d (10 mGy/d) or less does not appear likely to cause observable changes in terrestrial plant populations.
- Irradiation at chronic dose rates of 0.1 rad/d (1 mGy/d) or less does not appear likely to cause observable changes in terrestrial animal populations. The assumed threshold for effects in terrestrial animals is less than that for terrestrial plants, primarily because some species of mammals and reptiles are considered to be more radiosensitive.
- Reproductive effects on long-lived species with low reproductive capacity may require further consideration.

Additional summaries and reviews of radiation effects data on biota confirmed these findings; a discussion of these reviews may be found in DOE (2002).

Therefore, this model provides a graded approach to evaluate compliance with specified limits on radiation dose to populations of aquatic animals, terrestrial plants, and terrestrial animals. Specifically, these dose limits are:

- **Aquatic animals:** The absorbed dose to aquatic animals should not exceed 1 rad/d (10 milliGrays per day [mGy/d] or 0.4 milliGrays per hour [mGy/h]) from exposure to radiation or radioactive material releases into the aquatic environment. This dose limit is specified in DOE Order 5400.5.

- **Terrestrial plants:** The absorbed dose to terrestrial plants should not exceed 1 rad/d (10 mGy/d) from exposure to radiation or radioactive material releases into the terrestrial environment.
- **Terrestrial animals:** The absorbed dose to terrestrial animals should not exceed 0.1 rad/d (1 mGy/d) from exposure to radiation or radioactive material releases into the terrestrial environment.

Avoiding measurable impairment of reproductive capability is deemed to be the critical biological endpoint in establishing the dose limits for aquatic and terrestrial biota. As stated above, appreciable population effects would not be expected at doses lower than 1 rad/d and 0.1 rad/d, respectively, thereby establishing a level of adequate protection.

Internal and external sources of dose (and their contributing exposure pathways) were incorporated into the derivation of the graded approach methodology, and are based on the following general dose equation:

$$\text{Limiting Concentration} = \frac{\text{DoseRateLimit}}{(\text{InternalDoseRate}) + (\text{ExternalDoseRate}_{\text{soil / sed}}) + (\text{ExternalDoseRate}_{\text{water}})}$$

The limiting concentration in an environmental medium was calculated by first setting a target total dose (e.g., 1 rad/d) and then back-calculating the medium concentration necessary to produce the applicable dose from radionuclides in the organism (internal dose), plus the external dose components from radionuclides in the environment (external dose). The denominator of the generic equation represents the dose per unit media concentration and may be broken down into the base components of internal and external doses. Internal doses originate from radionuclides inside the organism's body. The internal dose is calculated as the product of the internal radionuclide concentration and an internal dose conversion factor. External doses originate from radionuclides external to the organism and are calculated as the product of the radionuclide concentration in the environmental medium in which the organism resides and an appropriate dose conversion factor.

The DOE defines a biota concentration guide (BCG) as the limiting concentration of a radionuclide in soil, sediment, or water that would not cause dose limits for protection of populations of aquatic and terrestrial biota to be exceeded. The BCGs used in the model are derived from the most sensitive potential receptor for which radionuclide toxicity data exist (for reproductive effects) for a given constituent.

Therefore, these receptors should be considered conservative indicators of risk and protective of less sensitive species. The receptors used are:

- "Riparian animal"
- "Terrestrial animal"
- "Aquatic animal"
- "Terrestrial plant"

The model compares a representative radionuclide concentration with generic BCGs and calculates a fraction, and in turn, these fractions are summed for each radionuclide in each medium. If the sum of all fractions is greater than 1.0, then the site does not pass the screen.

Because this approach is intended to be graded, performing the screen with the maximum detection of each radionuclide is considered the first tier, and therefore the most conservative evaluation. The second tier screen uses the arithmetic mean of constituent concentrations in order to be more realistic of actual site conditions.

For aquatic evaluations, the use of radionuclide concentration data from co-located sediment and surface water samples results in a less conservative, more realistic analysis. In the absence of one of the two media, the model derives the missing BCGs using a conservative sediment distribution coefficient (k_d) to calculate the environmental media radionuclide concentration. Results of the RAD-BCG screening are presented in the Risk Characterization section below.

Chemical Contaminant Screening. Chemical COPCs were selected on the basis of a comparison to existing ecologically-based benchmark values where applicable.

As noted above, existing media-specific benchmark values were identified as the preferred basis for comparison to constituent concentrations at the site. These benchmark values were obtained from several sources, each using unique methodologies and protocols in development of the respective values. The assumptions and methods followed in developing these benchmarks are described in the sources cited and the reader is referred to those publications for these details. In general, highly conservative assumptions are used in the development of these media and constituent-specific benchmarks. The intent of such an approach is to provide an estimate of a threshold concentration below which adverse effects are considered unlikely to even the most sensitive receptors, taking into account uncertainties associated with the data. The benchmarks utilized in the risk characterization may vary according to the differences in assumptions and methods followed. As an added measure of conservatism for COPC selection and risk characterization in the ERA, the lowest reported value was used in comparisons.

As noted on these data tables, the literature sources referenced for benchmarks for SVOC and/or metal constituents are as follows:

- Soils
 - Efroymson et al. 1997a; 1997b (Oak Ridge National Laboratory [ORNL])
 - Beyer 1990
- Sediment
 - Jones et al. 1987
 - Long et al. 1995
 - USEPA 1996a
 - Persaud et al. 1993
 - NOAA 1999
- Surface Water
 - USEPA 1996b
 - Suter and Tsao 1996 (ORNL)
 - NOAA 1999

Maximum constituent concentrations were compared to these benchmarks and a hazard quotient (HQ) was developed as follows:

$$HQ_i = C_i / TRV_i$$

Where:

- HQ = Hazard quotient for a given chemical in media i (unitless)
 C_i = Concentration of the chemical in media i (milligrams per kilogram [mg/kg] or micrograms per liter [µg/L])
 TRV = Toxicity reference value for a given chemical in media i (mg/kg or µg/L)

Chemicals with HQs greater than or equal to 1.0 were considered COPCs in the SERA.

Those constituents for which benchmarks do not exist were not analyzed quantitatively; a list of these is provided on Table 3-13; results of chemical contaminant screening are presented below in the Risk Characterization section.

3.5.3 Exposure Pathways and Routes

An exposure pathway links a source of contamination with one or more receptors via exposure to affected media. Exposure, and thus, potential for risk, can occur only if complete exposure pathways exist. As shown in Figures 3-3 through 3-8, the project area has potentially complete exposure pathways to ecological receptors.

The COPCs at the Sites originated at the REF many years ago. The primary transport mechanism for the COPCs (radionuclides, SVOCs, pesticides, PCBs) was adsorption to soil, sediment and suspended particulates. In Kress Creek and the WBDR, eventual redeposition downstream transported constituents into site media where they became available to biota. The current sources of contamination at the Sites are the contaminated soils at Upland STP and sediments and floodplain soils in Kress Creek and the WBDR. (See Figures 3-3 through 3-8.)

Exposure of ecological receptors to contaminants at the KCK and STP Sites is expected to occur primarily through ingestion and direct contact with contaminated surface soil, sediment, and surface water and through indirect exposure via ingestion of plant and prey items and subsequent bioaccumulation of contaminants through the food web.

It should be noted that a number of pathways at the site were potentially complete but not evaluated quantitatively in this SERA. For example, it is assumed that, while dermal contact with soil-bound contaminants could occur, the ingestion exposure route (both incidental ingestion of impacted media and food web exposures) likely accounts for the most significant dose for COPCs. Additionally, exposures associated with inhalation of and direct dermal contact with some COPCs may occur for some receptors, but it is assumed that such exposures are insignificant in relation to those associated with ingestion.

The following subsections provide brief discussions on the physicochemical characteristics as they relate to environmental disposition and the potential ecotoxicity of the COPCs at the KCK and STP sites.

Fate and Transport of the Constituents of Potential Concern

It is assumed that the surface water runoff from the REF was the primary mechanism for contamination reaching KCK. The outfall pool and the creek segment immediately below are the location of the highest concentrations of sediment contamination. Secondary

contaminant mechanisms include surface water runoff from contaminated properties within the KCK site watershed.

The primary migration mechanisms that may lead to the spread of contamination from the down stream area near the outfall are:

- Sediment transport and deposition within the stream to unaffected reaches
- Leaching of contaminants of concern from floodplain soils and sediments to surface water or groundwater
- Erosion of floodplain soils to the stream

The distribution coefficient for radium is approximately 250 mL/g and for thorium is 60,000 mL/g (Table 3-14). On the basis of these coefficients, radium and thorium particles are retained in soils and sediments. Because sediments and soils in the affected area tend to contain high percentages of fine materials (organic matter and clay), it is unlikely that radium and thorium will leach from the soil or sediments. Uranium has a lower distribution coefficient (45 mL/g), and thus may have a higher tendency to leach from soils and sediments.

Most metals have higher distribution coefficients (Table 3-15) and can be expected to react in the same way as radium and thorium. Exceptions are arsenic (1 to 18 mg/L), iron (1.4 to 1,000 mg/L), and selenium (1.2 to 8.6 mg/L), which have relatively low distribution coefficients and can be expected to leach.

The primary organic chemicals of concern identified at the site are PAHs, which generally have high partition coefficients (K_{oc}). They are not expected to leach from the soils or sediments.

Ecological Toxicity

Radionuclides. In general, the more primitive organisms are the most radioresistant taxonomic groups and the more advanced complex organisms, such as mammals, are the most radiosensitive. The early effects of exposure to ionizing radiation result primarily from cell death; cells that frequently undergo mitosis are the most radiosensitive, and cells that do not divide are the least. Thus, embryos and fetuses are particularly susceptible to ionizing radiation and very young animals are consistently more radiosensitive than adults (see review in Eisler 1994).

In addition to the evolutionary position and cell mitotic index, many extrinsic and intrinsic factors modify the response of a living organism for a given dose of radiation. Abiotic variables include the type and energy of radiation, exposure rate, length of exposure, total exposure and absorbed dose, dose rate, spatial distribution of dose, season, temperature, day length, and environmental chemicals; biotic variables include nutritional status, sensitizing or protective substances, competition, parasitism, and predation (Whicker and Schultz 1982; Hobbs and McClellan 1986; USCEAR 1988; Kiefer 1990).

Radiosensitivity of cells is related directly to their reproductive capacity and indirectly to their degree of differentiation (Hobbs and McClellan 1986). Early adverse effects of exposure to ionizing radiation are due mainly to the killing of cells. Cell death may result from the loss of reproductive integrity (i.e., inability to undergo mitosis). Reproductive death is important in rapidly dividing tissues such as bone marrow, skin, gut lining, and germinal epithelium. When the whole animal is exposed to a large dose of ionizing

radiation, some tissues are more prone to damage than others. Death rates of mammalian reproductive cells from ionizing radiation is modified by variations in the linear energy transfer of the radiation, the stage in the cell cycle, cell culture conditions (artifact), and sensitizing and protecting compounds (Barendsen 1990). The chemical form of the main stage of the acute radiation syndrome depends on the size and distribution of the absorbed dose. It is determined mainly by damage to blood platelets and other blood-forming organs at 4-5 Gy, to epithelial cells lining the small intestine at 5-30 Gy, and to brain damage at more than 30 Gy; death usually occurs within 48 hours at more than 30 Gy (McLean 1973).

Radioactive materials that gain entry to the body typically, through ingestion or inhalation, exert effects that are governed by their physical and chemical characteristics, which in turn influence their distributions and retention inside the body. In general, the radiation dose from internal emitters is a function of the effective half-time, energy released in the tissue, initial amount of introduced activity and mass of the organ (Hobbs and McClellan 1986). Retention of radionuclides by living organisms is quite variable and modified by numerous biologic and abiotic variables (Eisler 1994).

Chemical Contaminants. Several inorganics were positively detected in soil and sediment at the KCK and STP Sites. Of these, mercury is the only inorganic compound that both bioaccumulates and biomagnifies through the food chain. Mercury exposure could be important for the higher order predators that forage at the sites. The biological transformation of a variety of forms of mercury to methylmercury (the most toxic form) can take place in both terrestrial and aquatic environments (Olson and Cooper 1977 and Rogers 1976 *cited in* Heinz 1996). Other inorganic compounds detected at the sites that will bioaccumulate include lead, copper, and zinc. There are a variety of toxic mechanisms associated with metals.

PAHs are virtually ubiquitous in nature, primarily as a result of natural processes such as forest fires, microbial synthesis, and volcanic activity. They have been detected in animal and plant tissues, sediments, soils, air, surface water, drinking water, and groundwater. Anthropogenic sources of PAHs in the environment include high temperature combustion of organic materials typical of processes used in the steel industry, heating and power generation, and petroleum refining. PAHs in surface soils may be assimilated by plants, degraded by soil microorganisms, or accumulated to relatively high levels in the soil (Eisler 1987).

In some plants growing in highly contaminated areas, assimilation may exceed metabolism and degradation, resulting in accumulation in plant tissues. Laboratory experiments have demonstrated that plants can bioaccumulate PAHs to levels above those found in the environment, although this has not been conclusively demonstrated in field-grown plants. Uptake can be by both leaves (atmospheric deposition) and roots (soils and sediments) with subsequent translocation to other plant parts. Uptake is variable by plant species and soil conditions. Little data are available on bioaccumulation by vegetation and trophic transfer to higher level consumers in terrestrial and aquatic food chains (Eisler 1987).

PAHs are moderately persistent in the environment and therefore may potentially cause significant effects to vegetation, fish, and wildlife. The carcinogenicity of individual PAHs differs. Some lower weight compounds such as naphthalene, fluorenes, phenanthrenes, and anthracenes exhibit acute toxicity and other adverse effects to some organisms, but are

non-carcinogenic. In contrast, the higher molecular weight compounds are less acutely toxic, but many are carcinogenic, mutagenic, or teratogenic to a wide variety of organisms.

The pesticides detected at the sites are organochlorine compounds. The most serious environmental effects associated with exposure to organochlorine pesticides have occurred in birds. These effects include mortality, eggshell thinning, reduced reproductive success, population decline, and, in some cases, extirpation (Blus et al. 1996). Organochlorine pesticides are accumulated in lipids and biomagnify through the food chain.

The group known as PCBs contain congeners of differing persistence and toxicity in the environment. In general, PCB isomers with high lipophilicity and high numbers of substituted chlorines in adjacent positions constitute the greatest concern to wildlife due to their potential for bioaccumulation (Eisler 1986). Among sensitive avian species, PCBs disrupt normal patterns of growth, reproduction, metabolism, and behavior. In general, PCB accumulation is rapid and depuration proceeds at a much slower rate (NAS 1979).

Potential receptors at the sites include organisms that have significant direct contact with the soil. These could include plants, soil invertebrates, and animals that forage in soil or on organisms that have a high level of contact with the soil.

3.5.4 Ecological Receptors of Concern

A critical element of the problem formulation process is the identification of representative receptors that occur within the project area. As per USEPA guidance (USEPA 1997), these receptors should be conservative choices that are representative of the most highly exposed receptors to site media, groups essential to normal functioning of habitat, and federal or state threatened or endangered species.

Because of the complexity of natural systems, it is generally not possible to directly assess the potential impacts to all ecological receptors present within an area. Therefore, specific receptor species (e.g., great blue heron) or species groups (e.g., fish) are often selected as surrogates to evaluate potential risks to larger components of the ecological community (feeding guilds such as piscivorous birds) and are used to represent the assessment endpoints (e.g., survival and reproduction of piscivorous birds).

In the project area, the ecological receptors potentially at risk are those plants and animals that utilize terrestrial and aquatic habitats. Relevant groups of organisms include microbiota, aquatic and terrestrial plants, benthic/epibenthic macroinvertebrates, zooplankton, fish, reptiles, birds and mammals. These receptors were selected based on the habitat provided by the project area, the nature of COPCs (primarily soil- and sediment-associated contaminants and bioaccumulative compounds), and their high likelihood for exposure to COPCs. As such, they are considered the most at-risk receptors and are protective of receptor groups that would have less exposure to the affected environment. Relevant ecological receptors are discussed below.

Upper trophic level receptor species quantitatively evaluated in the ERA were limited to birds and mammals, the taxonomic groups with the most available information regarding exposure and toxicological effects. Lower trophic level receptor species were evaluated in the ERA based upon those taxonomic groupings for which screening values have been developed; these groupings and screening values are used in most ERAs. As such, specific species of aquatic biota

(e.g., macroinvertebrates) were not chosen as receptor species because of the limited information available for specific species and because aquatic biota are considered on a community level via a comparison to surface water and sediment screening values. Similarly, aquatic plants are considered protected by the federal Ambient Water Quality Criteria.

Additionally, other receptors that may be present onsite are threatened and/or endangered species such as the Indiana bat. Risks to this species cannot be estimated due to the paucity of toxicity values from the literature for this or related species. However, those receptors for which risk can be quantified would be expected to have greater potential exposure and, therefore, would be protective of other receptor groups with less exposure.

The following groups of receptors were evaluated using the hazard quotient screening technique described above.

Microbial Community

Microbial communities consist of bacteria, protozoans, and fungi and play several essential roles in ecosystems. They facilitate the degrading and transforming of detrital organic matter for ingestion by higher level consumers and serve as an important food source for a variety of larval and adult organisms. Additionally, microbes also play a role in the cycling and transformation of nutrients and sediments in the water column. The sediment microbial community would be at risk due to the direct exposure such communities might have to sediment-associated contaminants. Exposure of the microbial community to COPCs in the project area may significantly change or reduce community diversity. In turn, geochemical functions may be altered, reducing the productivity of these communities upon which many other receptors depend. Although specific information on the composition of the microbial community in the KCK/STP area is not available, this community is an essential component of the ecosystem.

Plants

As primary producers, plants are an important food source for herbivorous organisms and also provide essential habitat for a variety of aquatic and terrestrial species at all life stages. Plants are an essential component of ecosystems and exposure to contaminants may result in a loss of productivity within the ecosystem and limit the ability of the site to support ecological resources.

Invertebrates

Invertebrates (i.e., primary consumers) serve an important function in the aquatic and terrestrial food webs by consuming plants, detritus, etc., and are a food source for fish, birds and mammals. They represent an important link between organic matter and higher trophic level consumers. They are in intimate contact with sediments and soils and may be highly exposed to adsorbed contaminants. Reducing or impairing the function of invertebrate organisms may disrupt the flow of energy within the ecosystem. Therefore, impacts to this portion of the food web may have profound consequences to wildlife receptors, potentially resulting in decreases in fish, reptile, avian, and mammalian populations in and around the project area.

The following upper trophic level receptor species have been chosen for exposure modeling with media at the Sites based on the criteria listed above; it should be noted that these

receptors were evaluated for bioaccumulation and food web transfer of chemical constituents only. Radionuclide bioaccumulation was accounted for with similar receptors in the RAD-BCG screening models.

Mammals

Deer Mouse (*Peromyscus maniculatus*)—Terrestrial Mammalian Omnivore. Deer mice feed on seeds, berries, acorns, fruits, insects, and other small invertebrates, and serve as food for a variety of carnivores. They are the direct link in the terrestrial food chain between plants and higher trophic level organisms.

Least Shrew (*Cryptotis parva*)—Terrestrial Mammalian Insectivore. Shrews feed mainly on insects, earthworms, and other invertebrates, and would be expected to ingest significant amounts of soils incidentally through foraging and prey consumption.

Mink (*Mustela vison*)—Semi-Aquatic Mammalian Piscivore. Mink are top level carnivores that feed on fish, small mammals, birds, eggs, frogs, and macroinvertebrates. They are also known to be sensitive to environmental contaminants.

Raccoon (*Procyon lotr*)—Semi-Aquatic Mammalian Omnivore. Raccoons are most common in and around wetland areas, where they search for small aquatic animals like fish, crayfish, and freshwater mussels in the shallow water. Besides aquatic life and other animal matter, raccoons also eat a variety of fruits, berries, and seeds.

Birds

American Robin (*Turdus migratorius*)—Terrestrial Avian Omnivore. Robins live in a variety of habitats, including woodlands, swamps, suburbs, and parks. Robins forage on the ground in open areas, along edge habitats, or along the edges of streams. They forage along the ground for ground-dwelling invertebrates and search for fruit and foliage-dwelling insects in low tree branches (Malmborg and Willson 1988).

Mallard (*Anas platyrhynchos*)—Wetland/Aquatic Avian Omnivore. Mallards consume a wide variety of foods including vegetation, insects, worms, gastropods, and arthropods. Due to their feeding habits, mallards also tend to incidentally ingest significant amounts of sediment during feeding.

Great Blue Heron (*Ardea herodias*)—Wetland/Aquatic Avian Piscivore. Great blue herons represent carnivorous wading birds that feed on a variety of aquatic organisms, including fish, invertebrates, amphibians, and reptiles. Herons do not ingest significant amounts of sediment during feeding activities.

3.5.5 Screening Assessment Endpoints

The conclusion of the problem formulation stage includes the selection of assessment and measurement endpoints, based on the preliminary conceptual model. Endpoints in the SERA define ecological attributes that are to be protected (assessment endpoints) and measurable characteristics of those attributes (measurement endpoints) that can be used to gauge the degree of impact that has or could occur. Assessment endpoints most often relate to attributes of biological populations or communities, and are intended to focus the risk assessment on particular components of the ecosystem that could be adversely affected by

contaminants from the site (USEPA 1997). Assessment endpoints contain an entity (e.g., fish-eating birds) and an attribute of that entity (e.g., survival rate).

Because of the complexity of natural systems, it is generally not possible to directly assess the potential impacts to all ecological receptors present within an area. Therefore, receptor species (e.g., great blue heron) or species groups (e.g., fish) are often selected as surrogates to evaluate potential risks to larger components of the ecological community (feeding guilds; e.g., piscivorous birds) represented in the assessment endpoints (e.g., survival and reproduction of piscivorous birds). Selection criteria typically include those species that:

- Are known to occur, or are likely to occur, at the site
- Have a particular ecological, economic, or aesthetic value
- Are representative of taxonomic groups, life history traits, and/or trophic levels in the habitats present at the site for which complete exposure pathways are likely to exist
- Can, because of toxicological sensitivity or potential exposure magnitude, be expected to represent potentially sensitive populations at the site
- Have sufficient ecotoxicological information available on which to base an evaluation

Based on the habitat and types of contaminants present, seven assessment endpoints were chosen to evaluate the risk to ecological receptor populations from toxic components in KCK and STP site media. Each assessment endpoint and corresponding representative species or community is described below.

Survival and Reproduction of Terrestrial Plant Communities

Plants provide food, cover, and nesting material for many animals. The soils at the sites will support fewer birds and mammals if COPCs are limiting the survival and reproduction of plants.

Survival and Reproduction of Soil Invertebrate Communities

Soil invertebrates promote soil fertility by breaking down organic matter and releasing nutrients. They also improve aeration, drainage, and aggregation of soil, and serve as a forage base for many terrestrial species. The soils at the sites will support fewer insectivorous birds and mammals if chemical concentrations are limiting the survival and reproduction of soil invertebrates.

The endpoints that build on the above and were evaluated in this risk assessment are:

- Survival and reproduction of terrestrial mammalian omnivores (deer mouse)
- Survival and reproduction of terrestrial mammalian insectivores (least shrew)
- Survival and reproduction of terrestrial avian omnivores (American robin)

Assessment endpoints with aquatic bases that were evaluated herein are:

- Survival and reproduction of semi-aquatic mammalian piscivores (mink)
- Survival and reproduction of semi-aquatic mammalian omnivores (raccoon)
- Survival and reproduction of semi-aquatic avian omnivores (mallard)
- Survival and reproduction of semi-aquatic avian piscivores (great blue heron)

Endpoints specific to reptiles and amphibians were not selected, although potential exposure pathways may exist for these receptors. There is a lack of herpetofauna-specific toxicological data for most environmental contaminants. Reptiles and amphibians are indirectly assessed via the bird and mammals evaluations since they are not likely to be more sensitive than the receptors evaluated (Hall and Henry 1992). Birds and mammals have been selected that have similar diets to the herpetofauna that could potentially inhabit the KCK and STP sites.

The corresponding measurement endpoints associated with each assessment endpoint were defined as follows:

Assessment Endpoints	Measurement Endpoints
Survival and reproduction of soil invertebrate communities.	⇒ Comparison of HQs for soil invertebrates (earthworms) to a target HQ of 1. Medium-specific HQs are calculated for individual contaminants by dividing the maximum soil concentration by a soil benchmark that is intended to be protective of soil invertebrates.
Survival and reproduction of terrestrial plant communities.	⇒ Comparison of HQs for terrestrial plants to a target HQ of 1. Medium-specific HQs are calculated for individual contaminants by dividing the maximum soil concentration by a soil benchmark that is intended to be protective of terrestrial plants.
Survival and reproduction of avian terrestrial omnivores.	⇒ Comparison of HQs for American robin to a target HQ of 1. Receptor-specific HQs are calculated for individual contaminants by dividing an estimated level of exposure (dose) by a screening toxicity value that is associated with no adverse effects. Exposure estimates will include contributions from the consumption of plants, invertebrates, and soil.
Survival and reproduction of mammalian terrestrial insectivores.	⇒ Comparison of HQs for least shrew to a target HQ of 1. Exposure estimates will include contributions from the consumption of invertebrates, and soil.
Survival and reproduction of mammalian semi-aquatic omnivores.	⇒ Comparison of HQs for raccoon to a target HQ of 1. Exposure estimates will include contributions from the consumption of plants, invertebrates, fish and sediment.
Survival and reproduction of mammalian semi-aquatic piscivores.	⇒ Comparison of HQs for mink to a target HQ of 1. Exposure estimates will include contributions from the consumption fish and sediment
Survival and reproduction of mammalian terrestrial insectivores.	⇒ Comparison of HQs for least shrew to a target HQ of 1. Exposure estimates will include contributions from the consumption of invertebrates and soil.
Survival and reproduction of avian semi-aquatic omnivores.	Comparison of HQs for mallards to a target HQ of 1. Exposure estimates will include contributions from the consumption of plants, invertebrates, and sediment.
Survival and reproduction of avian semi-aquatic piscivores.	⇒ Comparison of HQs for great blue heron to a target HQ of 1. Exposure estimates will include contributions from the consumption of fish.

Screening Level Exposure Estimate and Risk Characterization

Upper trophic level receptor exposures to chemical contaminants at the Kress Creek and STP sites were determined by estimating the concentration of each chemical in each relevant dietary component.

4.1 Exposure Point Concentrations

The bioaccumulation of site-related constituents by plants and soil invertebrates (and hence, upper trophic level receptors) was estimated using models and maximum measured media concentrations. The methodology and models used to derive these estimates are described below. It is important to note that only those constituents listed on Table 4-2 of "Bioaccumulation Testing and Interpretation for the Purpose of Sediment Quality Assessment Status and Needs" (USEPA 2000) were included in the evaluation of bioaccumulation.

4.1.1 Plants

Tissue concentrations in the aboveground vegetative portion of plants were estimated by multiplying the maximum measured surface soil concentration for each chemical by chemical-specific soil-to-plant bioconcentration factors (BCFs) obtained from the literature. The BCF values used were based on root uptake from soil and on the ratio between dry-weight soil and dry-weight plant tissue. Literature values based on the ratio between dry-weight soil and wet-weight plant tissue were converted to a dry-weight basis by dividing the wet-weight BCF by the estimated solids content for terrestrial plants (15 percent [0.15]; Sample et al. 1997).

For inorganic chemicals without literature based BCFs, a soil-to-plant BCF of 1.0 was assumed. For organic chemicals without literature based BCFs, soil-to-plant BCFs were estimated using the algorithm provided in Travis and Arms (1988):

$$\log B_v = 1.588 - (0.578) (\log K_{ow})$$

where: B_v = Soil-to-plant BCF (unitless; dry weight basis)
 K_{ow} = Octanol-water partitioning coefficient (unitless)

The log K_{ow} values used in the calculations were obtained mostly from USEPA (1995; 1996c) and are listed in Table 4-1. The soil-to-plant BCFs used in the SERA are shown in Table 4-2.

4.1.2 Soil Invertebrates

Tissue concentrations in soil invertebrates (earthworms) were estimated by multiplying the maximum measured surface soil concentration for each chemical by chemical-specific BCFs

or bioaccumulation factors (BAFs) obtained from the literature. BCFs are calculated by dividing the concentration of a chemical in the tissues of an organism by the concentration of that same chemical in the surrounding environmental medium (in this case, soil) without accounting for uptake via the diet. BAFs consider both direct exposure to soil and exposure via the diet. Since earthworms consume soil, BAFs are more appropriate values and are used in the food web models when available. BAFs based on depurated analyses (soil was purged from the gut of the earthworm prior to analysis) are given preference over undepurated analyses when selecting BAF values since direct ingestion of soil is accounted for separately in the food web model.

The BCF/BAF values used were based on the ratio between dry-weight soil and dry-weight earthworm tissue. Literature values based on the ratio between dry-weight soil and wet-weight earthworm tissue were converted to a dry-weight basis by dividing the wet-weight BCF/BAF by the estimated solids content for earthworms (16 percent [0.16]; USEPA 1993). For inorganic chemicals without available measured BAFs or BCFs, an earthworm BAF of 1.0 was assumed. The soil-to-earthworm BCFs/BAFs used in the SERA are shown in Table 4-2.

4.1.3 Small Mammals

Whole-body tissue concentrations in small mammals (shrews, voles, and/or mice) were estimated using one of two methodologies. For chemicals with literature-based soil-to-small mammal BCFs, the small mammal tissue concentration was obtained by multiplying the maximum measured surface soil concentration for each chemical by a chemical-specific soil-to-small mammal BCF obtained from the literature. The BCF values used were based on the ratio between dry-weight soil and whole-body dry-weight tissue. Literature values based on the ratio between dry-weight soil and wet-weight tissue were converted to a dry-weight basis by dividing the wet-weight BCF by the estimated solids content for small mammals (32 percent [0.32]; USEPA 1993). BCFs for shrews were those reported in Sample et al. (1998) for insectivores (or for general small mammals if insectivore values were unavailable), for voles were those reported for herbivores, and for mice were those reported for omnivores.

For chemicals without soil-to-small mammal BCF values, an alternate approach was used to estimate whole-body tissue concentrations. Because most chemical exposure for these small mammal species is via the diet, it was assumed that the concentration of each chemical in the small mammal's tissues was equal to the chemical concentration in its diet, that is, a diet to whole-body BAF (wet-weight basis) of 1.0 was assumed. The use of a diet to whole-body BAF of 1.0 is likely to result in a conservative estimate of chemical concentrations for chemicals that are not known to biomagnify in terrestrial food chains (e.g., aluminum). For chemicals that are known to biomagnify (e.g., PCBs), a diet to whole-body BAF value of 1.0 will likely result in a realistic estimate of tissue concentrations based on reported literature values. For example, a maximum BAF (wet weight) value of 1.0 was reported by Simmons and McKee (1992) for PCBs based on laboratory studies with white-footed mice. Menzie et al. (1992) reported BAF values (wet-weight) for dichlorodiphenyl trichloroethane (DDT) of 0.3 for voles and 0.2 for short-tailed shrews. Reported BAF (wet-weight) values for dioxin were only slightly above 1.0 (1.4) for the deer mouse (USEPA 1990). Resulting tissue concentrations (wet-weight) were then converted to dry weight using an estimated solids

content of 32 percent (see above). The soil-to-small mammal BAFs used in the SERA are shown in Table 4-2.

4.2 Dietary Intakes

Dietary intakes for each receptor species were calculated using the following formula (modified from USEPA 1993):

$$DI_x = \frac{[[\sum_i (FIR)(FC_{xi})(PDF_i)] + [(FIR)(SC_x)(PDS)] + [(WIR)(WC_x)]]}{BW}$$

where: DI_x	=	Dietary intake for chemical x (mg chemical/kg body weight/day)
FIR	=	Food ingestion rate (kg/day, dry-weight)
FC_{xi}	=	Concentration of chemical x in food item i (mg/kg, dry weight)
PDF_i	=	Proportion of diet composed of food item i (dry weight basis)
SC_x	=	Concentration of chemical x in soil/sediment (mg/kg, dry weight)
PDS	=	Proportion of diet composed of soil/sediment (dry weight basis)
WIR	=	Water ingestion rate (L/day) (not applicable for this ERA)
WC_x	=	Concentration of chemical x in water (mg/L) (not applicable for this ERA)
BW	=	Body weight (kg, wet weight)

Exposure parameters for upper trophic level receptors are presented in Table 4-3.

4.2.1 Ingestion Screening Values

Ingestion screening values for dietary exposures were derived for each avian/mammalian receptor species and bioaccumulating chemical. Toxicological information from the literature for wildlife species most closely related to the receptor species was used, where available, but was supplemented by laboratory studies of non-wildlife species (e.g., laboratory mice) where necessary. The ingestion screening values are expressed as milligrams of the chemical per kilogram body weight of the receptor per day (mg/kg-BW/day).

Growth and reproduction were emphasized as assessment endpoints since they are the most relevant, ecologically, to maintaining viable populations and because they are generally the most studied chronic toxicological endpoints for ecological receptors. If several chronic toxicity studies were available from the literature, the most appropriate study was selected for each receptor species based on study design, study methodology, study duration, study endpoint, and test species. No Observed Adverse Effect Levels (NOAELs) based on growth and reproduction were used, where available, as the screening values. When chronic NOAEL values were unavailable, estimates were derived or extrapolated from chronic Lowest Observed Adverse Effect Levels (LOAELs) or acute values as follows:

- When values for chronic toxicity were not available, the median lethal dose (LD_{50}) was used. An uncertainty factor of 100 was used to convert the acute LD_{50} to a chronic NOAEL (i.e., the LD_{50} was multiplied by 0.01 to obtain the chronic NOAEL).

- An uncertainty factor of 10 was used to convert a reported LOAEL to a NOAEL.

Ingestion screening values for mammals and birds are summarized in Tables 4-4 and 4-5, respectively.

Other assumptions used in the models to determine the potential for food web transfer are as follows:

- **Area use:** The portion of a receptors home range that is impacted, assumed to be 100 percent
- **Bioavailability:** The percentage of the concentration of a COPC in an exposure medium that is taken up and metabolized by a receptor, assumed to be 100 percent.
- **Body weight:** The mean body weight of the population of receptors of a given species, assumed to be the minimum reported values by USEPA (1993).
- **Ingestion rates:** The average mass of food or environmental media ingested on a daily basis by the ROC, assumed to be the maximum value reported by USEPA (1993).
- **Bioaccumulation:** The degree to which a COPC concentrates in the tissues of biota at progressively higher trophic levels in the food web. The maximum value reported in the literature reviewed was used. In the absence of a chemical- and/or trophic level-specific value, a default value of 1.0 is assumed.
- **Dietary composition:** The percentage of diet comprised of various prey or forage material. If values are reported for various food items by USEPA (1993), these were incorporated into the calculations (Table 4-3). In the absence of species-specific information, it is assumed that the receptor's diet is comprised entirely of the most contaminated food items.

The results of the food web modeling are presented in the following section.

4.3 Screening Level Risk Characterization

The screening-level risk calculation is the final step in a SERA. In this step, the maximum exposure doses to upper trophic level receptor species are compared with the corresponding screening values to derive screening risk estimates. The outcome of this step is a list of COPCs for each media-pathway-receptor combination evaluated for a conclusion of acceptable or unacceptable risk.

To reiterate, KCK, STP river and STP upland data were screened using the RAD-BCG model to determine risks to aquatic and terrestrial receptors from exposure to radionuclides; the results of those screens are presented here. Chemical COPCs were selected using the HQ method, which entails dividing the exposure dose by the corresponding benchmark. The TRVs used here are media-specific values developed using conservative assumptions regarding toxicity and exposure and are intended to be levels protective of adverse impacts to even highly sensitive species. The lowest value derived from the literature was adopted for the purposes of comparison.

HQs exceeding one indicate the potential for risk since the chemical concentration or dose (exposure) exceeds the screening value (effect). However, screening values and exposure estimates are derived using intentionally conservative assumptions such that HQs greater

than 1.0 do not necessarily indicate that risks are present or impacts are occurring. Rather, such HQs identify chemical-pathway-receptor combinations requiring further evaluation. Following the same reasoning, HQs that are less than or equal to 1.0 indicate that risks are very unlikely, allowing a conclusion of no unacceptable risk to be reached with a high degree of confidence.

The results of the risk screening for each area of concern and each affected medium within the KCK and STP sites are presented in Tables 4-6 to 4-19 and are described further below.

4.3.1 Kress Creek

Radionuclides

A comparison of the maximum radionuclide concentrations detected in KCK site media with DOE's BCGs resulted in the following (see Table 4-6):

- Total sum of fractions in water and sediment was $2.96E+03$, and therefore the site screen failed.
- Radium-228 (Ra-228) had a partial fraction of 7.46 in sediment (and a calculated partial fraction of $2.8E+03$ in surface water) and appeared to be the risk driver.
- No radionuclide analyses were performed on collected surface water samples, and therefore the resultant screen was based on modeled water concentrations.

A comparison of the mean radionuclide concentrations detected in KCK site media with BCGs resulted in the following (Table 4-7):

- The site screen failed; however, this was based wholly on the calculated partial fractions of Ra-228 ($1.2E+02$) and radium-226 (Ra-226) ($1.4E+01$) in water; the total sum of fractions in sediment was below 1.0.
- The default distribution coefficient (k_d) values used in the model for both Ra-226 and Ra-228 are low (70 mL/g; USDOE 2002), indicating a theoretical propensity to migrate into the aqueous fraction; this is the reason for the high calculated partial fractions for these constituents. According to Langmuir (1997) and Oztunali and Roles (1984), radium (Ra) has a K_d value of 250 mL/g for soils similar to those in West Chicago. Therefore, under natural conditions, Ra-226 and Ra-228 remain bound to particulate fractions and measured surface water concentrations would be expected to be lower than those calculated by the model (see further discussion regarding fate and transport characteristics of these constituents above).

Chemical Contaminants

Table 4-8 and Table 4-9 summarize the comparison of maximum and mean analyte concentrations to available benchmark values for KCK sediment and surface water respectively. Constituents were considered risk drivers if the resultant HQ was greater than 10; results of this screen are presented below.

Sediment

Ten inorganic constituents, 13 semivolatile organics, *p,p'*-dichlorodiphenyl dichloroethane (*p,p'*-DDD), and Aroclor 1260 were detected in KCK sediments at concentrations exceeding respective ecological benchmarks (i.e., had HQs greater than 1.0). Of these, five metals

(arsenic, copper, lead, mercury, and zinc), six polycyclic aromatic hydrocarbons (PAHs) (acenaphthene, acenaphthylene, anthracene, benzo(a)anthracene, fluorene, and phenanthrene) and DDD and Aroclor 1260 had HQs greater than 10; HQs ranged from 13 to 179 for inorganics, 11 to 39 for SVOCs, 140 for DDD and 870 for Aroclor 1260.

Surface Water

Barium, cobalt, copper, lead and nickel exceeded ecological benchmarks for surface water (i.e. had HQ's greater than 1.0). Barium and copper had HQs greater than 10 (27 and 63, respectively).

Food Web Modeling Results

It appears likely that metals, Aroclor 1260, and some PAHs have the potential to bioaccumulate to significant levels in semi-aquatic receptors that are dependent upon KCK (see Table 4-10). Of these, modeled body burdens of aluminum, lead, mercury, zinc, chrysene, and pyrene in mink, great blue heron, and mallard were the highest relative to ecological benchmarks.

4.3.2 Sewage Treatment Plant River

Radionuclides

The following are the results of the RAD-BCG screen conducted with the maximum constituent concentrations in STP river sediment and surface water (Table 4-11):

- Total sum of fractions in water and sediment was 1.95, and therefore the site screen failed.
- Ra-228 had a partial fraction of 1.25 in sediment and 0.37 in water and was the risk driver.

However, when the mean concentrations were used, the total sum of fractions was below 1.0, and therefore, the site screen passed (Table 4-12).

Chemical Contaminants

STP River Sediment. Nine inorganics exceeded benchmarks in STP sediments (i.e. had HQ's greater than 1.0); only mercury had a maximum hazard quotient of greater than 10 (25). Additionally, four PAHs also had HQs greater than 1, but none were greater than 10 (see Table 4-13).

STP River Surface Water. Six inorganics exceeded benchmarks in STP surface water (i.e. had HQ's greater than 1.0); however, only barium had an HQ greater than 10 (24) (see Table 4-14).

Food Web Modeling Results

The food web model determined that concentrations of accumulated burdens of aluminum, mercury, chrysene, and pyrene in receptors exposed to STP river sediments and surface water exceeded ecotoxicological benchmarks (see Table 4-15). In particular, great blue heron and mallard (i.e., avian receptors that feed primarily on fish and aquatic invertebrates [as well as plants, in the case of the mallard]), had the highest modeled burdens of COPCs relative to benchmarks.

4.3.3 Sewage Treatment Plant Upland Soils

Radionuclides

The following are the results for the RAD-BCG screen conducted with the maximum constituent concentrations in STP Upland Soils (Table 4-16):

- Total sum of fractions in soil was 1.1E+01, and therefore, the site screen failed.
- Ra-228 had a partial fraction of 10 in soil and was the risk driver).

However, when the mean concentrations were used the total sum of fractions was below 1.0, and therefore the site screen passed (Table 4-17).

Chemical Contaminants

Fourteen inorganics exceeded benchmarks in STP surface soils (i.e, had HQs greater than 1.0) as well as 11 PAHs and one volatile organic (toluene). Of these, chromium, lead, manganese, iron, mercury, vanadium, and zinc had HQs greater than 10; HQs ranged from 20.85 (vanadium) to 5,400 (mercury) (see Table 4-18).

Food Web Modeling Results

For terrestrial receptors, metals were the primary accumulated COPCs, in particular lead, mercury, and zinc; cadmium and chromium were also important (Table 4-19). The least shrew and American robin had highest body burdens of these constituents relative to benchmarks, due to high accumulations in invertebrate (insects and earthworms) prey.

Uncertainty Assessment

Each step in the screening ERA process involves the use of assumptions and protocol that impart uncertainty to the final results. As noted above, whenever possible, assumptions that tend to increase conservatism are adopted to ensure that the likelihood for underestimating the potential for effects is minimized. In some cases, however, the absence of technical information concerning the toxicology of a given constituent or other factors precludes the consideration of a chemical or exposure route in the quantitative assessment. The exclusion of potential COPCs and potentially complete routes of exposure for receptors of concern will tend to be a source of downward bias to estimates of potential for effects. That is, such factors may offset some of the conservatism imparted to the process.

Some of the primary sources of uncertainty and their probable affect on the overall conservatism inherent in the analysis for this SERA are presented below.

5.1 Limiting the Analysis to Constituents of Potential Concern that Exceed Background and Established Benchmarks

More chemicals were detected in media at the sites than were quantitatively evaluated in the SERA. Assuming that the locations where background samples were collected represent unimpacted areas, this is a valid and accepted approach for screening chemicals as part of the risk assessment process. In some cases, background levels exceeded benchmark values, indicating that site concentrations would not increase risk to receptors beyond that experienced in the general environment in the region.

5.2 Use of Established Benchmark Values for Comparison

In general, these values have been developed using highly conservative assumptions regarding chemical fate and transport characteristics, physicochemical properties, ecotoxicological endpoints, and exposure conditions. Consistent with the general principles described by USEPA (1997b) for screening level ERAs, these values tend to incorporate significant margins of error.

5.3 Inability to Quantitatively Evaluate All Detected Analytes

Some chemical constituents could also not be quantitatively evaluated because of the paucity of available toxicological data. Therefore, the potential exists for disregarding constituents that could have an effect on the environment.

5.4 Limiting Evaluation of Potentially Complete Exposure Routes to Ingestion

Other routes of contact with COPCs may be complete for some receptors. In general, it is believed that ingestion of impacted media, forage, and prey items constitute the most significant route for most vertebrate receptors. Moreover, little if any technical information to support the quantitative evaluation of non-ingestion pathways in ecological receptors exists for most chemicals. As such, the uncertainties potentially associated with assumptions that would be necessary to do so would make the results highly questionable.

5.5 Use of Default Value of 1.0 for Bioaccumulation Factor

This may be an overestimate or underestimate, depending on the chemical, the medium, and the trophic level under consideration.

5.6 Assumptions Regarding Conversion of Literature-Based Toxicity Data into Toxicity Reference Values

A significant degree of subjectivity and uncertainty is involved with this process, particularly when short-term studies or lethal endpoints are involved. The degree to which the assumptions can be considered conservative is dependant upon the chemical under consideration.

5.7 Assumptions Regarding Area Use, Bioavailability, Body Weight, Ingestion Rate, and Other Exposure Factors

In the absence of any USEPA-approved information to the contrary, the most conservative assumptions were adopted across the board for these factors, leading to a highly conservative estimate of the potential for exposure.

5.8 Assumptions Regarding Potential Additive and Synergistic Effects

The response of an organism to combinations of toxicants may be increased or decreased because of toxicological responses at the site of action. These responses may be “additive” – the combined effect of two chemicals is equal to the sum of each individual agent (for example, $2 + 2 = 4$), or synergistic – the combined effects of two chemicals is far greater than the sum of the effects of each agent alone (for example $2 + 2 = 40$). Because these types of responses are difficult to quantify in a non-laboratory setting, they are generally not evaluated in an ecological risk assessment. Therefore, the conclusions drawn herein may be underestimates of actual biological responses.

5.9 Use of the Lowest Reported Benchmark for Comparison

These values can sometimes vary over orders of magnitude for the same COPC (e.g., arsenic in surface water, fluoranthene in sediment). Selecting the lowest value would tend to increase conservatism.

5.10 Data Limitations

For certain areas and media, data were available for a limited set of analytes, and for a limited sample size (i.e., generally less than 10). In some cases, only inorganic analyses were available (e.g., surface water) or radionuclide analyses were not conducted (e.g., Kress Creek surface water). Data were collected over different time frames for some portions of the study area and combined with data from earlier investigations.

5.10.1 Specific Limitations of the RAD-BCG Model

The evaluation of radionuclide effects on aquatic systems using the RAD-BCG model proceeds through an analysis of both sediment and surface water components. In the absence of one of the two parts, the model calculates these values based on established physicochemical characteristics for the radionuclides of concern. However, actual media concentrations of radionuclides in sediments or surface water may be higher or lower than those predicted by the model, and therefore the eventual screen may not be wholly indicative of actual site conditions.

Additionally, not all detected radionuclides could be screened using the RAD-BCG model because some constituents have not been adequately tested for toxicity in wildlife receptors. As a result, some constituents could create deleterious effects that may be unevaluated.

Conclusions

Radionuclides and chemical contaminants, at concentrations high enough to potentially adversely affect ecological receptors, have impacted sediments, surface water, and soils at the KCK and STP Sites. For each area of concern and each complete pathway identified, the analytical data were evaluated to determine the potential for ecological risk. Finally, a determination was made as to whether:

1. Risks are acceptable
2. Risks are unacceptable (i.e., calculated HQs were greater than 1) and require immediate mitigation
3. Risks are equivocal and require further investigation

In general, based on the SERA results, it cannot be concluded that there is acceptable risk, and therefore, further investigation would be required to determine the actual risk.

The following sections discuss the conclusions of the SERA for the KCK and STP sites.

6.1 Kress Creek

6.1.1 Radionuclides

The potential for adverse ecological effects in KCK sediments appears to be associated with maximum and mean detections of radionuclides, primarily Ra-226 and Ra-228, the daughter products of uranium and thorium decomposition, respectively. The potential for effects associated with radionuclides may be underestimated due to the unavailability of benchmarks for some radioisotopes that were detected in sediments but not evaluated with the RAD-BCG model.

6.1.2 Chemical Contaminants

Copper, lead, mercury, chrysene, and pyrene are the most important chemical COPCs. While the target HQ of 1.0 for wildlife receptors was exceeded for other metals, PAHs, and Aroclor 1260, these exceedences appear to be minor and may be mitigated by the conservatism inherent in the screening analysis.

The HQs estimated for surface water indicate that few analytes occur at levels sufficient to warrant their inclusion as COPCs, and that exceedences of target HQ values are very slight.

6.2 Sewage Treatment Plant River

6.2.1 Radionuclides

Radionuclide concentrations were markedly lower in sediments from the STP river as compared to KCK, with only Ra-228 demonstrating a HI greater than target risk of 1.0. Concentrations of the uranium isotopes and Ra-226 did not exceed BCGs. Mean concentrations of radionuclides appear to be protective of sensitive wildlife species.

6.2.2 Chemical Contaminants

In general, chemical constituent concentrations in sediments and surface water associated with the STP river were very similar to those in KCK media, except for mercury, which was almost twice that of KCK. Therefore, the list of inorganic and organic COPCs, and those with the potential to affect ecological risk to upper trophic level receptors, was also similar.

Barium appeared to be the dominant constituent of concern in STP river water samples; other inorganic constituents demonstrated slight exceedences, and radionuclide concentrations were low in general. Therefore, risks from sediments and surface water from the STP should be considered lower than those from KCK.

6.3 Sewage Treatment Plant Upland

6.3.1 Radionuclides

Although concentrations of Ra-228 and Th-232 were half those of KCK sediments, these two radionuclides have greater ecological effects on terrestrial mammals than on aquatic receptors, and as a result, have higher partial fractions relative to screening benchmarks. The mean concentrations of radionuclides are significantly lower and do not result in a total sum of fractions greater than 1.0. Therefore, the mean concentrations can be considered protective of sensitive receptors.

6.3.2 Chemical Contaminants

Concentrations of lead and mercury were significant in surface soils collected near the STP. As a result, these two constituents demonstrated high bioaccumulation (and high exceedences of benchmarks) in terrestrial receptors, primarily those that feed on invertebrate prey. It should be noted, however, that the inherent conservatism of the ERA paradigm requires an evaluation of the “worst case scenario” for the site.

6.4 Discussion

These quantitative results should also be considered in the context of the qualitative characterization of habitat quality and the occurrence of other stressors within the study area. As noted above, high quality aquatic and riparian habitat is generally limited to the lower reaches of the study area. Additional stressors related to residential and commercial development within and in close proximity to the study area in the upper reaches may contribute to the relatively poor habitat quality in those areas and may be responsible for

the chemical constituents seen in media samples collected in Kress Creek. These locations generally coincide with the occurrence of both radionuclide and chemical COPCs at levels that significantly exceed ecologically-based benchmarks, indicating the potential for adverse effects. The combination of effects potentially associated with the COPCs and those associated with radiological stressors may further increase the possibilities of adverse impacts to ecological receptors in the upper reaches of the Kress Creek system and as a result, remedial activities should focus on the mitigation of these sediments. The proposed cleanup standard of 7.2 pCi/g for combined radium-226 and radium-228 is protective of biota when compared to the toxicological thresholds used here to calculate risk (e.g., BCGs for uranium and Ra-226 and Ra-228 are 2000 pCi/g (U-238); 100 pCi/g and 90 pCi/g, respectively).

References

BBL. 2004. Remedial Investigation Report, Kress Creek/West Branch DuPage River and Sewage Treatment Plant Sites, DuPage County, Illinois..

Barendsen, G.W. 1990. :Mechanisms of cell reproductive death and shapes of radiation dose-survival curves of mammalian cells.: *Int J Rad Biol.*, 57: 885-896.

Beyer, W.N. 1990. Evaluating soil contamination. *U.S. Fish Wildl. Serv., Biol. Rep.* 90(2).

Blus, L.J., S.N. Wiemeyer, and C.J. Henny. 1996. Organochlorine Pesticides. Chapter 6 in *Non-infectious Diseases of Wildlife, Second Edition* (eds., A. Fairbrother, L.N. Locke, and G.L. Hoff). Ames, Iowa: Iowa State University Press.

CH2M HILL. 1993. Source Characterization and Hydrologic Assessment: Sewage Treatment Plant, West Chicago Illinois. Prepared for USEPA WA 51-5FQW /Contract No. 68-W8-0040. September 30, 1993.

CH2M HILL. 1994. Technical Memorandum dated February 17, 1994. Source Characterization and Hydrologic Assessment: Kerr-McGee Kress Creek/West Branch DuPage River Site, West Chicago, Illinois. Prepared for USEPA. February 17, 1994.

CH2M HILL. 1995. Source Characterization and Hydrologic Assessment: Kerr-McGee Sewage Treatment Plant/West Branch DuPage River Site, West Chicago Illinois. Prepared for USEPA WA 51-5FQW /Contract No. 68-W8-0040. March 2, 1995.

Efroymson, R.A., M.E. Will, G.W. Suter, II, and A. C. Wooten. 1997a. Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Terrestrial Plants: 1997 Revision. Prepared for US Department of Energy. ES/ER/TM-85/R3. November 1997.

Efroymson, R.A., M.E. Will, and G.W. Suter, II. 1997b. Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Soil and Litter Heterotrophic Process: 1997 Revision. Prepared for US Department of Energy. ES/ER/TM-126/R2. November 1997.

Eisler, R. 1987. Polycyclic aromatic hydrocarbon hazards to fish, wildlife, and invertebrates: A synoptic review. *U.S. Fish and Wildlife Service Biological Report 85(1.11)*, Contaminant Hazard Reviews Report No. 11.

Eisler, R. 1994. Radiation hazards to fish, wildlife, and invertebrates. A synoptic review. *U.S. Fish and Wildlife Service, Biological Report 26*, Contaminant Hazard Reviews, Report 29. National Biological Service, U.S. Department of the Interior.

Hall, R.J. and P.F.P. Henry. 1992. "Assessing effects of pesticides on amphibians and reptiles: status and needs." *Herpetological Journal*. 2: 65-71.

- Hobbs, C.H. and R.O. McClellan. 1986. Toxic effects of radiation and radioactive materials. In: Cassarett and Doull's Toxicology. eds. Klaassen, C.D., M.O. Amdur, and J. Doull, Third Edition. New York: MacMillan. 669-705.
- Jones, D. S., G. W. Suter II, and R. N. Hull. 1997. Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Sediment- Associated Biota: 1997 Revision. ES/ER/TM-95/R4. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Kiefer, J. 1990. Biological Radiation Effects. Berlin: Springer-Verlag.
- Languir, D., 1997. *Aqueous Environmental Geochemistry*: Prentice Hall, New Jersey.
- Long, E. R. et al. 1995. "Incidence of Adverse Biological Effects Within Ranges of Chemical Concentrations In Marine and Estuarine Sediments." *Environ. Manag.* 19: 81-97.
- Malmberg, P.K., and M.F. Willson. 1988. Foraging ecology of avian frugivores and some consequences for seed dispersal in an Illinois woodlot. *Condor.* 90: 173-186.
- McLean, A.S. 1973. "Early adverse effects of radiation." *Brit Med J.* 29: 69-73.
- Heinz, G.H. 1996. Mercury Poisoning in Wildlife. Chapter 11 in *Non-infectious Diseases of Wildlife*, Second Edition (eds., A. Fairbrother, L.N. Locke, and G.L. Hoff). Ames, Iowa: Iowa State University Press.
- Menzie, C.A., D.E. Burmaster, J.S. Freshman, and C.A. Callahan. 1992. "Assessment of methods for estimating ecological risk in the terrestrial component: a case study at the Baird & McGuire Superfund Site in Holbrook, Massachusetts." *Environmental Toxicology and Chemistry.* 11:245-260.
- NAS. 1979. Polychlorinated biphenyls. Rep Comm Assess PCBs in the Environ., *Environ Stud Bd., Comm Nat Resour., Natl Res Coun., Natl Acad Sci.* Washington, D.C.
- NOAA. 1999. Screening Quick Reference Tables. HAZMAT Report 99-1. Updated September 1999.
- Ozuntali, O. I., and G. W. Roles. 1984. *De Minimus Waste Impacts Analysis Methodology*. U. S. Nuclear Regulatory Commission. NUREG/CR-3585.
- Persaud, D., R. Jaagumagi, and A. Hayton. 1993. Guidelines for the Protection and Management of Aquatic Sediment Quality in Ontario. Ontario Ministry of the Environment, Water Resources Branch. Revised 1993.
- RESRAD, Table E.3, page 202. Manual for Implementing Residual Radioactive Material Guidelines Using RESRAD, Version 5.0; ANL/EAD/LD-2.
- Sample, B.E., M.S. Aplin, R.A. Efroymsen, G.W. Suter II, and C.J.E. Welsh. 1997. Methods and tools for estimation of the exposure of terrestrial wildlife to contaminants. Environmental Sciences Division, Oak Ridge National Laboratory. ORNL/TM-13391.
- Sample, B.E., J.J. Beauchamp, R.A. Efroymsen, G.W. Suter II, and T.L. Ashwood. 1998. Development and validation of bioaccumulation models for earthworms. Environmental Restoration Division, ORNL Environmental Restoration Program. ES/ER/TM-220.

- Simmons, G.J. and M.J. McKee. 1992. "Alkoxyresorufin metabolism in white-footed mice at relevant environmental concentrations of Aroclor 1254." *Fundamental and Applied Toxicology*. 19:446-452.
- Suter, G. W. and C. L. Tsao. 1996. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Aquatic Biota: 1996 Revision*. Lockheed Martin Energy Systems, Inc. Oak Ridge, Tennessee.
- U.S. Department of Energy. 2002. *A Graded Approach for Evaluating Radiation Doses to Aquatic and Terrestrial Biota*. DOE-STD-1153-2002.
- United Nations Scientific Committee on the Effects of Atomic Radiation. 1988. *Sources, effects and risks of ionizing radiation*. New York: United Nations.
- U.S. Environmental Protection Agency. 2000. *Bioaccumulation Testing and Interpretation for the Purpose of Sediment Quality Assessment*. EPA-823-R-00-001. February 2000.
- U.S. Environmental Protection Agency. 1997. *Ecological risk assessment guidance for Superfund: process for designing and conducting ecological risk assessments*. Interim Final. EPA/540/R-97/006.
- U. S. Environmental Protection Agency. 1996a. *Soil Screening Guidance*. Office of Emergency and Remedial Response. EPA/540/R-96/018. April 1996.
- U. S. Environmental Protection Agency. 1996b. *Ecotox Thresholds in ECO Update*. Vol. 3 No. 2. January 1996. U. S. Environmental Protection Agency, Office of Emergency and Remedial Response, Washington, DC.
- U.S. Environmental Protection Agency. 1996c. *Superfund chemical data matrix*. EPA/540/R-96/028.
- U.S. Environmental Protection Agency. 1995. *Internal report on summary of measured, calculated and recommended log k_{ow} values*. Environmental Research Laboratory, Athens, Georgia. April 10, 1995.
- U.S. Environmental Protection Agency. 1993. *Wildlife exposure factors handbook*. Volume I of II. EPA/600/R-93/187a.
- U.S. Environmental Protection Agency. *The Superfund Public Health Evaluation Manual (SPHEM)*. From the Office of Emergency and Remedial Response (OERR). EPA 540/1-86-060. October 1986.
- Travis, C.C. and A.D. Arms. 1988. "Bioconcentration of organics in beef, milk, and vegetation." *Environmental Science and Technology*. 22:271-274.
- Whicker, F.W. and V. Schultz. 1982. *Radioecology: Nuclear Energy and the Environment*. Volume 1. Boca Raton, Florida: CRC Press.

Tables

TABLE 3-1

Wildlife Species Potentially Occurring and Habitat Associations
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Species	Cover Types			
	Streams, Rivers & Small Ponds	Wetland Areas Including Edges of Ponds and Creeks	Meadows or Open Areas	Upland Woods
Birds				
Great Blue Heron/ <i>Ardea herodias</i> *	U	U		
Great Egret/ <i>Casmerodius albus</i>	U	U		
Green-Backed Heron/ <i>Butorides straitus</i>	F	F		
Black-Crowned Night Heron/ <i>Nycticorax nycticorax</i>	U	U		
Canada Goose/ <i>Branta canadensis</i> *	A	A	A	
Wood Duck/ <i>Aix sponsa</i> *	F	F		
Mallard/ <i>Anas platyrhynchos</i> *	A	A		
Killdeer/ <i>Charadrius vociferus</i> *	C	C	C	
Spotted Sandpiper/ <i>Actitis macularia</i>	U	U		
Chimney Swift/ <i>Chaetura pelagica</i>	F		F	
Belted Kingfisher/ <i>Ceryle alcyon</i>	U	U		
Purple Martin/ <i>Progne subis</i>		U	U	
Tree Swallow/ <i>Tachycineta bicolor</i>	F	F	F	
N. Rough-Winged Swallow/ <i>Stelgidopteryx serripennis</i>	U	U	U	
Bank Swallow/ <i>Riparia riparia</i>		U	U	
Barn Swallow/ <i>Hirundo rustica</i> *	C	C	C	
American Crow/ <i>Corvus rachyrhynchus</i> *			C	C
Cedar Waxwing/ <i>Bombycilla cedrorum</i> *				U
Blue Winged Teal/ <i>Anas discors</i>	U	U		
Red-Tailed Hawk/ <i>Buteo jamaicensis</i> *		F	F	F
American Kestrel/ <i>Falco sparverius</i>		F	F	
Ring-Necked Pheasant/ <i>Phasianus colchicus</i> *			F	
Sora/ <i>Porzana carolina</i>		U	U	
American Coot/ <i>Fulica americana</i>		U	U	
Mourning Dove/ <i>Zenaida macroura</i> *		C	C	C
Eastern Screech-Owl/ <i>Otus asio</i>				F
Great Horned Owl/ <i>Bubo virginianus</i>		F	F	F
Common Nighthawk/ <i>Chordeiles minor</i>		F	F	
Chimney Swift/ <i>Chaetura pelagica</i>		F	F	F
Red-Headed Woodpecker/ <i>Melanerpes erthrocephalus</i>			F	F
Willow Flycatcher/ <i>Empidonax traillii</i>		U		
Eastern Kingbird/ <i>Tyrannus tyrannus</i>		C	C	C
Common Yellowthroat/ <i>Geothlypis trichas</i> *		C	C	

TABLE 3-1
 Wildlife Species Potentially Occurring and Habitat Associations
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Species	Cover Types			
	Streams, Rivers & Small Ponds	Wetland Areas Including Edges of Ponds and Creeks	Meadows or Open Areas	Upland Woods
Birds (Cont'd)				
Song Sparrow/ <i>Melospiza melodia</i> *		C	C	
Northern Oriole/ <i>Icterus galbula</i> *	C	C		
Common Grackle/ <i>Quiscalus quiscula</i>		C	A	A
Brown-Headed Cowbird/ <i>Molothrus ater</i>		C	C	C
Rock Dove/ <i>Columba livia</i>			A	
Northern Flicker/ <i>Colaptes auratus</i> *			C	C
Eastern Bluebird/ <i>Sialia sialis</i>			R	R
American Robin/ <i>Turdus migratorius</i> *			A	A
Brown Thrasher/ <i>Toxostoma rufum</i>			C	C
Dickcissel/ <i>Spiza americana</i>			U	
Field Sparrow/ <i>Spizella pusilla</i>			F	
Savannah Sparrow/ <i>Passerculus sandwichensis</i>			U	
Grasshopper Sparrow/ <i>Ammodrammus savannarum</i>			U	
Bobolink/ <i>Dolichonyx oryzivorus</i>			F	
Red-Winged Blackbird/ <i>Agelaius phoeniceus</i> *	A	A	A	
Eastern Meadowlark/ <i>Sturnella magna</i> *			F	
American Goldfinch/ <i>Carduelis tristis</i> *			C	
House Sparrow/ <i>Passer domesticus</i> *			A	A
American Woodcock/ <i>Scolopax minor</i>		U		
Black-Billed Cuckoo/ <i>Coccyzus erythrophthalmus</i>				U
Yellow-Billed Cuckoo/ <i>Coccyzus americanus</i>				U
Red-Bellied Woodpecker/ <i>Melanerpes carolinus</i>				U
Downy Woodpecker/ <i>Picoides pubescens</i> *				F
Hairy Woodpecker/ <i>Picoides villosus</i>				U
Eastern Wood-Pewee/ <i>Contopus virens</i> *				U
Eastern Phoebe/ <i>Sayornis phoebe</i>				U
Great Crested Flycatcher/ <i>Myiarchus crinitus</i> *				F
Blue Jay/ <i>Cyanocitta cristata</i> *				C
Black-Capped Chickadee/ <i>Parus atricapillus</i> *				C
Tufted Titmouse/ <i>Parus bicolor</i>				R
White-Breasted Nuthatch/ <i>Sitta carolinensis</i>				U
House Wren/ <i>Troglodytes aedon</i> *				C
Blue-Gray Gnatcatcher/ <i>Poliophtila caerulea</i>				R
Veery/ <i>Catharus fuscescens</i>				R

TABLE 3-1
 Wildlife Species Potentially Occurring and Habitat Associations
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Species	Cover Types			
	Streams, Rivers & Small Ponds	Wetland Areas Including Edges of Ponds and Creeks	Meadows or Open Areas	Upland Woods
Birds (Cont'd)				
Wood Thrush/ <i>Hylocichla mustelina</i> *				U
Gray Catbird/ <i>Dumetella carolinensis</i> *				C
Indigo Bunting/ <i>Passerina cyanea</i> *			F	
Brown Thrasher/ <i>Toxostoma rufum</i>				C
Red-Eyed Vireo/ <i>Vireo olivaceus</i> *				F
Yellow Warbler/ <i>Dendroica petechia</i>		F		F
Common Yellowthroat/ <i>Geothlypis trichas</i> *		C		C
Scarlet Tanager/ <i>Piranga olivacea</i>				U
Northern Cardinal/ <i>Cardinalis cardinalis</i> *				C
Amphibians				
Bullfrog/ <i>Rana catesbeiana</i> *	C	C		
Eastern Tiger Salamander/ <i>Ambystoma tigrinum</i>		U	U	U
American Toad/ <i>Bufo americanus</i>		F	F	F
Western Chorus Frog/ <i>Pseudacris t. triseriata</i>	F	F	F	
Green Frog/ <i>Rana calmitans melanota</i> *	C	C		
Northern Leopard Frog/ <i>Rana pipiens pipiens</i>	U	U		
Reptiles				
Common Snapping Turtle/ <i>Chelydra serpentina</i>	C	C		
Midland Painted Turtle/ <i>Chrysemys picta marginata</i>	A	A		
Red-Eared Slider/ <i>Trachemys scripta elegans</i>	U	U		
Eastern Spiny Softshell/ <i>Aplouson s. spinifera</i>	R	R		
Northern Water Snake/ <i>Nerodia sipedon</i>	R	R		
Western Fox Snake/ <i>Elaphe vulpina vulpina</i>			F	F
E. Plains Garter Snake/ <i>Thamnophis radix radix</i>			F	F
Midland Brown Snake/ <i>Storeria dekayi wrightorum</i>			U	U
Eastern Garter Snake/ <i>Thamnophis s. semifaciata</i> *			F	F
Mammals				
Raccoon/ <i>Procyon lotor</i> *	C	C	C	C
Beaver/ <i>Castor canadensis</i> *	U	U		
Muskrat/ <i>Ondatra zibethicus</i>	F	F		
Virginia Opossum/ <i>Didelphis virginiana</i> *		F	F	F
Masked Shrew/ <i>Blarina brevicauda</i>		U	U	U
Mink/ <i>Mustela vison</i>		U		U

TABLE 3-1
 Wildlife Species Potentially Occurring and Habitat Associations
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Species	Cover Types			
	Streams, Rivers & Small Ponds	Wetland Areas Including Edges of Ponds and Creeks	Meadows or Open Areas	Upland Woods
Mammals (Cont'd)				
Striped Skunk/Mephitis mephitis*		F	F	F
Red Fox/Vulpes vulpes		U	U	U
Short-Tailed Shrew/Blarina brevicauda			U	U
Least Weasel/Mustela nivalis			R	R
Coyote/Canis latrans		R	R	R
Woodchuck/Marmota monax*			F	F
13-Lined Ground Squirrel/Spermophilus tridecemlineatus			F	
White -Footed Mouse/Peromyscus leucopus			C	C
Meadow Vole/Microtus pennsylvanicus*			C	
Eastern Cottontail/Sylvilagus floridanus*		F	F	F
White-Tailed Deer/Odocoileus virginianus		F	F	F
Eastern Mole/Scalopus aquaticus*			U	U
Little Brown Myotis/Nyotis lucifugus				U
Silver-Haired Bat/Lasionycteris noctivagans				R
Red Bat/Lasiurus borealis				U
Big Brown Bat/Eptesicus fuscus				U
Hoary Bat/Lasiurus cinereus				R
Eastern Chipmunk/Tamias striatus			C	C
Gray Squirrel/Sciurus carolinensis*				F
Fox Squirrel/Sciurus niger*			F	F
Southern Flying Squirrel/Glaucomys volans				U

Status Codes:

Introduced=I, Abundant=A, Common =C, Fairly Common=F, Uncommon=U, Unknown=N, Rare=R or Watch list=W, Threatened=T, and Endangered=E

Source: FPDDC 1991

TABLE 3-2
 Macroinvertebrate Inventory Results - WBDR
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Species	GBK-07 (Upstream of Kress Creek)	GBK-05 (Downstream of Kress Creek)
Chironomidae		
<i>Ablabesmyia</i> sp.		1
<i>Ablabesmyia mallochi</i>	18	
<i>Chironomus</i> sp.		2
<i>Cryptochisaromus fulves</i>	6	
<i>Dicrotendipes neounodestus</i>	28	
<i>Glyptotendipes</i> sp.		10
<i>Glyptotendipes lobiferus</i>	57	
<i>Nanocladius</i> sp.		1
<i>Orthocladius</i> sp.	14	
<i>Polypedilum</i> sp.		2
<i>Polypedilum illinoense</i>	21	
<i>Procladius</i> sp.	2	
<i>Thienemannimyia</i> gr.	28	2
Tabanidae		
<i>Chrysops</i> sp.		1
Mollusca		
<i>Ferrissia</i> sp.		1
<i>Physo</i> sp.	1	3
Hirudinea		
		21
Isopoda		
<i>Caecidotea</i> sp.		3
<i>Caecidotes intermedia</i>		1
Decapoda		
<i>Cambarida</i>		1
Ephemeroptera		
<i>Baetis</i> sp.	26	
<i>Baetis flavistriga</i>	4	
<i>Caenis</i> sp.	6	7
<i>Stenacron</i> sp.	1	
<i>Stenacron interpunctatum</i>		4
<i>Stenacron minnetonka</i>	2	
<i>Stenonema</i> sp.	2	
<i>Tricorythodes</i> sp.	1	
Odonata		
<i>Argia</i> sp.	2	
<i>Argia tibialis</i>		3
<i>Enallagma signatum</i>		1
<i>Ischnura</i> sp.		4
Trichoptera		
<i>Cheumatopsyche</i> sp.	194	
Coleoptera		
<i>Stenelmis</i> sp.	3	
<i>Stenelmis crenata</i>		5
<i>Stenelmis vittipennis</i>	46	
Total Number of Organisms	462	73
Total Number of Taxa	17	18

IEPAWPC/88-010, 1988

TABLE 3-3
 Fish Survey Results
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Species	Kress Creek	W. Branch Du Page River
Goldfish		—*
Carp x Goldfish		—*
Carp	X	X
Golden Shiner	X	X
Sand Shiner	—	—*
Bluntnose Minnow	X	X
Fathead Minnow	—	—*
Creek Chub	X	—*
Quillback		—*
White Sucker	X	X
Black Bullhead	X	X
Green Sunfish	X	X
Bluegill		—*
Hybrid Sunfish	X	—*
Largemouth Bass	X	—*
White Crappie		—*
Black Crappie	X	X
Gizzard Shad		X

*Reported by IEPA/WPC/88-010, 1988.

X = observed by CH2M HILL

— = not observed by CH2M HILL

Table 3-4
Identification of Constituents of Potential Concern
Process Summary - Sediment/Floodplain Soil - Kress Creek
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in Sediment/Floodplain Soil	Constituents preceded by * → * were removed because they are considered essential nutrients (Ca, Mg, K, Na) or daughter products (radiological)	Constituents preceded by * → * were removed due to low frequency of detection (<5% of samples)	Constituents preceded by * → * result of background comparison (Max < Bkgd)	Constituents preceded by * → * were removed because no applicable screening criteria exist	Constituents preceded by * → * were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in Sediment/Floodplain Soil
Surface Sediment/Floodplain Soil (< 24")						
INORGANICS	INORGANICS	INORGANICS	INORGANICS	INORGANICS	INORGANICS	INORGANICS
Aluminum	Aluminum	Aluminum	Aluminum	Aluminum	→ Aluminum	--
Antimony	Antimony	→ Antimony	--	--	--	--
Arsenic	Arsenic	Arsenic	Arsenic	Arsenic	Arsenic	Arsenic
Barium	Barium	Barium	Barium	→ Barium	--	--
Beryllium	Beryllium	Beryllium	Beryllium	→ Beryllium	--	--
Cadmium	Cadmium	Cadmium	Cadmium	Cadmium	Cadmium	Cadmium
Calcium	→ Calcium	--	--	--	--	--
Chromium, Total	Chromium, Total	Chromium, Total	Chromium, Total	Chromium, Total	Chromium, Total	Chromium, Total
Cobalt	Cobalt	Cobalt	Cobalt	→ Cobalt	--	--
Copper	Copper	Copper	Copper	Copper	Copper	Copper
Iron	Iron	Iron	Iron	Iron	→ Iron	--
Lead	Lead	Lead	Lead	Lead	Lead	Lead
Magnesium	→ Magnesium	--	--	--	--	--
Manganese	Manganese	Manganese	Manganese	Manganese	Manganese	Manganese
Mercury	Mercury	Mercury	Mercury	Mercury	Mercury	Mercury
Nickel	Nickel	Nickel	Nickel	Nickel	Nickel	Nickel
Potassium	→ Potassium	--	--	--	--	--
Selenium	Selenium	Selenium	Selenium	→ Selenium	--	--
Silver	Silver	Silver	Silver	Silver	→ Silver	--
Sodium	→ Sodium	--	--	--	--	--
Thallium	Thallium	Thallium	Thallium	→ Thallium	--	--
Vanadium	Vanadium	Vanadium	Vanadium	→ Vanadium	--	--
Zinc	Zinc	Zinc	Zinc	Zinc	Zinc	Zinc

Table 3-4
 Identification of Constituents of Potential Concern
 Process Summary - Sediment/Floodplain Soil - Kress Creek
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in Sediment/Floodplain Soil	Constituents preceded by * → * were removed because they are considered essential nutrients (Ca, Mg, K, Na) or daughter products (radiological)	Constituents preceded by * → * were removed due to low frequency of detection (<5% of samples)	Constituents preceded by * → * result of background comparison (Max < Bkgd)	Constituents preceded by * → * were removed because no applicable screening criteria exist	Constituents preceded by * → * were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in Sediment/Floodplain Soil
Surface Sediment/Floodplain Soil (< 24")						
PESTICIDES/PCBs p,p'-DDD Aroclor 1260 SEMIVOLATILES 1,2,4-Trichlorobenzene 2-Methylnaphthalene Acenaphthene Acenaphthylene Anthracene Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(K)Fluoranthene Bis(2-Ethylhexyl) Phthalate Carbazole Chrysene Dibenzofuran Di-N-Butyl Phthalate Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Naphthalene Phenanthrene Pyrene	PESTICIDES/PCBs p,p'-DDD Aroclor 1260 SEMIVOLATILES 1,2,4-Trichlorobenzene 2-Methylnaphthalene Acenaphthene Acenaphthylene Anthracene Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(K)Fluoranthene Bis(2-Ethylhexyl) Phthalate Carbazole Chrysene Dibenzofuran Di-N-Butyl Phthalate Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Naphthalene Phenanthrene Pyrene	PESTICIDES/PCBs p,p'-DDD Aroclor 1260 SEMIVOLATILES 1,2,4-Trichlorobenzene 2-Methylnaphthalene Acenaphthene Acenaphthylene Anthracene Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(K)Fluoranthene Bis(2-Ethylhexyl) Phthalate Carbazole Chrysene Dibenzofuran Di-N-Butyl Phthalate Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Naphthalene Phenanthrene Pyrene	PESTICIDES/PCBs p,p'-DDD Aroclor 1260 SEMIVOLATILES 1,2,4-Trichlorobenzene 2-Methylnaphthalene Acenaphthene Acenaphthylene Anthracene Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(K)Fluoranthene Bis(2-Ethylhexyl) Phthalate Carbazole Chrysene Dibenzofuran Di-N-Butyl Phthalate Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Naphthalene Phenanthrene Pyrene	PESTICIDES/PCBs p,p'-DDD Aroclor 1260 SEMIVOLATILES 1,2,4-Trichlorobenzene 2-Methylnaphthalene Acenaphthene Acenaphthylene Anthracene Benzo(A)Anthracene Benzo(A)Pyrene → Benzo(B)Fluoranthene → Benzo(K)Fluoranthene Bis(2-Ethylhexyl) Phthalate → Carbazole Chrysene Dibenzofuran Di-N-Butyl Phthalate Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Naphthalene Phenanthrene Pyrene	PESTICIDES/PCBs p,p'-DDD Aroclor 1260 SEMIVOLATILES → 1,2,4-Trichlorobenzene 2-Methylnaphthalene Acenaphthene Acenaphthylene Anthracene Benzo(A)Anthracene Benzo(A)Pyrene -- -- → Bis(2-Ethylhexyl) Phthalate -- Chrysene → Dibenzofuran → Di-N-Butyl Phthalate Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Naphthalene Phenanthrene Pyrene	PESTICIDES/PCBs p,p'-DDD Aroclor 1260 SEMIVOLATILES -- 2-Methylnaphthalene Acenaphthene Acenaphthylene Anthracene Benzo(A)Anthracene Benzo(A)Pyrene -- -- -- -- Chrysene -- -- Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Naphthalene Phenanthrene Pyrene

Table 3-4
Identification of Constituents of Potential Concern
Process Summary - Sediment/Floodplain Soil - Kress Creek
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in Sediment/Floodplain Soil	Constituents preceded by * → * were removed because they are considered essential nutrients (Ca, Mg, K, Na) or daughter products (radiological)	Constituents preceded by * → * were removed due to low frequency of detection (<5% of samples)	Constituents preceded by * → * result of background comparison (Max < Bkgd)	Constituents preceded by * → * were removed because no applicable screening criteria exist	Constituents preceded by * → * were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in Sediment/Floodplain Soil
Surface Sediment/Floodplain Soil (< 24")						
RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL
Actinium-228	Actinium-228	Actinium-228	Actinium-228	→ Actinium-228	--	--
Bismuth-212	Bismuth-212	Bismuth-212	Bismuth-212	→ Bismuth-212	Bismuth-212	--
Bismuth-214	Bismuth-214	Bismuth-214	Bismuth-214	→ Bismuth-214	--	--
Cesium-137	Cesium-137	Cesium-137	Cesium-137	Cesium-137	Cesium-137	Cesium-137
Lead-211	Lead-211	Lead-211	Lead-211	→ Lead-211	Lead-211	--
Lead-212	Lead-212	Lead-212	Lead-212	→ Lead-212	--	--
Lead-214	Lead-214	Lead-214	Lead-214	→ Lead-214	--	--
Potassium-40	Potassium-40	Potassium-40	Potassium-40	→ Potassium-40	--	--
Protactinium-234	Protactinium-234	Protactinium-234	Protactinium-234	→ Protactinium-234	--	--
Radium-223	Radium-223	Radium-223	Radium-223	→ Radium-223	--	--
Radium-224	Radium-224	Radium-224	Radium-224	→ Radium-224	--	--
Radium-226	Radium-226	Radium-226	Radium-226	Radium-226	Radium-226	Radium-226
Radium-228	Radium-228	Radium-228	Radium-228	Radium-228	Radium-228	Radium-228
Rhenium-226	Rhenium-226	Rhenium-226	Rhenium-226	→ Rhenium-226	--	--
Rhenium-228	Rhenium-228	Rhenium-228	Rhenium-228	→ Rhenium-228	--	--
Rhenium-total	Rhenium-total	Rhenium-total	Rhenium-total	→ Rhenium-total	--	--
Thallium-208	Thallium-208	Thallium-208	Thallium-208	→ Thallium-208	--	--
Thorium-227	Thorium-227	Thorium-227	Thorium-227	→ Thorium-227	--	--
Thorium-228	Thorium-228	Thorium-228	Thorium-228	→ Thorium-228	--	--
Thorium-230	Thorium-230	Thorium-230	Thorium-230	→ Thorium-230	--	--
Thorium-232	Thorium-232	Thorium-232	Thorium-232	Thorium-232	Thorium-232	Thorium-232
Thorium-234	Thorium-234	Thorium-234	Thorium-234	→ Thorium-234	--	--
Uranium-234	Uranium-234	Uranium-234	Uranium-234	Uranium-234	Uranium-234	Uranium-234
Uranium-235	Uranium-235	Uranium-235	Uranium-235	Uranium-235	Uranium-235	Uranium-235
Uranium-238	Uranium-238	Uranium-238	Uranium-238	Uranium-238	Uranium-238	Uranium-238

Table 3-5
 Identification of Constituents of Potential Concern
 Process Summary - Surface Water - Kress Creek
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in Surface Water	Constituents preceded by " → " were removed because they are considered essential nutrients (Ca, Mg, K, Na)	Constituents preceded by " → " were removed due to low frequency of detection (<5% of samples)	Constituents preceded by " → " result of background comparison (Max < Bkgd)	Constituents preceded by " → " were removed because no applicable screening criteria exist	Constituents preceded by " → " were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in Surface Water
INORGANICS Aluminum Arsenic Barium Calcium Chromium, Total Cobalt Copper Iron Lead Magnesium Manganese Nickel Potassium Selenium Sodium Vanadium Zinc	INORGANICS Aluminum Arsenic Barium → Calcium Chromium, Total Cobalt Copper Iron Lead → Magnesium Manganese Nickel → Potassium Selenium → Sodium Vanadium Zinc	INORGANICS Aluminum Arsenic Barium -- Chromium, Total Cobalt Copper Iron Lead -- Manganese Nickel -- Selenium -- Vanadium Zinc	INORGANICS → Aluminum → Arsenic Barium -- → Chromium, Total Cobalt Copper → Iron Lead -- → Manganese Nickel -- → Selenium -- Vanadium → Zinc	-- -- Barium -- -- Cobalt Copper -- Lead -- -- Nickel -- -- Vanadium --	INORGANICS -- -- Barium -- -- Cobalt Copper -- → Lead -- -- Nickel -- -- → Vanadium --	INORGANICS -- -- Barium -- -- Cobalt Copper -- -- -- Nickel -- -- -- -- --

Table 3-6
Identification of Constituents of Potential Concern
Process Summary - Sediment/Floodplain Soil - STP River
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in Sediment/Floodplain Soil	Constituents preceded by " → " were removed because they are considered essential nutrients (Ca, Mg, K, Na)	Constituents preceded by " → " were removed due to low frequency of detection (<5% of samples)	Constituents preceded by " → " result of background comparison (Max < Bkgd)	Constituents preceded by " → " were removed because no applicable screening criteria exist	Constituents preceded by " → " were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in Sediment/Floodplain Soil
Surface Sediment/Floodplain Soil (< 24")						
INORGANICS	INORGANICS	INORGANICS	INORGANICS	INORGANICS	INORGANICS	INORGANICS
Aluminum	Aluminum	Aluminum	Aluminum	Aluminum	→ Aluminum	--
Antimony	Antimony	Antimony	Antimony	Antimony	→ Antimony	--
Arsenic	Arsenic	Arsenic	Arsenic	Arsenic	Arsenic	Arsenic
Barium	Barium	Barium	Barium	Barium	→ Barium	--
Beryllium	Beryllium	Beryllium	Beryllium	→ Beryllium	--	--
Cadmium	Cadmium	Cadmium	Cadmium	Cadmium	Cadmium	Cadmium
Calcium	→ Calcium	--	--	--	--	--
Chromium, Total	Chromium, Total	Chromium, Total	Chromium, Total	Chromium, Total	Chromium, Total	Chromium, Total
Cobalt	Cobalt	Cobalt	Cobalt	→ Cobalt	--	--
Copper	Copper	Copper	Copper	Copper	Copper	Copper
Iron	Iron	Iron	Iron	Iron	→ Iron	--
Lead	Lead	Lead	→ Lead	--	--	--
Magnesium	→ Magnesium	--	--	--	--	--
Manganese	Manganese	Manganese	Manganese	Manganese	Manganese	Manganese
Mercury	Mercury	Mercury	Mercury	Mercury	Mercury	Mercury
Nickel	Nickel	Nickel	Nickel	Nickel	Nickel	Nickel
Potassium	→ Potassium	--	--	--	--	--
Selenium	Selenium	Selenium	→ Selenium	--	--	--
Silver	Silver	Silver	Silver	Silver	Silver	Silver
Sodium	→ Sodium	--	--	--	--	--
Thallium	Thallium	Thallium	Thallium	→ Thallium	→ Thallium	--
Vanadium	Vanadium	Vanadium	Vanadium	→ Vanadium	→ Vanadium	--
Zinc	Zinc	Zinc	Zinc	Zinc	Zinc	Zinc

Table 3-6
Identification of Constituents of Potential Concern
Process Summary - Sediment/Floodplain Soil - STP River
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in Sediment/Floodplain Soil	Constituents preceded by * → * were removed because they are considered essential nutrients (Ca, Mg, K, Na)	Constituents preceded by * → * were removed due to low frequency of detection (<5% of samples)	Constituents preceded by * → * result of background comparison (Max < Bkgd)	Constituents preceded by * → * were removed because no applicable screening criteria exist	Constituents preceded by * → * were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in Sediment/Floodplain Soil
Surface Sediment/Floodplain Soil (< 24")						
SEMIVOLATILES Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(G,H,I)Perylene Benzo(K)Fluoranthene Carbazole Chrysene Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Phenanthrene Pyrene VOLATILES 2-Butanone Acetone Carbon Disulfide Toluene RADIOLOGICAL Potassium-40 Radium-226 Radium-228 Thorium-227 Thorium-228 Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	SEMIVOLATILES Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(G,H,I)Perylene Benzo(K)Fluoranthene Carbazole Chrysene Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Phenanthrene Pyrene VOLATILES 2-Butanone Acetone Carbon Disulfide Toluene RADIOLOGICAL → Potassium-40 Radium-226 Radium-228 Thorium-227 Thorium-228 Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	SEMIVOLATILES Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(G,H,I)Perylene Benzo(K)Fluoranthene Carbazole Chrysene Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Phenanthrene Pyrene VOLATILES 2-Butanone Acetone Carbon Disulfide Toluene RADIOLOGICAL -- Radium-226 Radium-228 Thorium-227 Thorium-228 Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	SEMIVOLATILES Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(G,H,I)Perylene Benzo(K)Fluoranthene Carbazole Chrysene Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Phenanthrene Pyrene VOLATILES 2-Butanone Acetone Carbon Disulfide Toluene RADIOLOGICAL -- Radium-226 Radium-228 Thorium-227 Thorium-228 Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	SEMIVOLATILES Benzo(A)Anthracene Benzo(A)Pyrene Benzo(B)Fluoranthene Benzo(G,H,I)Perylene Benzo(K)Fluoranthene Carbazole Chrysene Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Phenanthrene Pyrene VOLATILES 2-Butanone Acetone Carbon Disulfide Toluene RADIOLOGICAL -- Radium-226 Radium-228 Thorium-228 Thorium-228 -- Thorium-232 Uranium-234 Uranium-235 Uranium-238	SEMIVOLATILES Benzo(A)Anthracene → Benzo(A)Pyrene -- -- -- -- → Chrysene → Fluoranthene Fluorene Indeno(1,2,3-C,D)Pyrene Phenanthrene → Pyrene VOLATILES -- -- -- → Toluene RADIOLOGICAL -- Radium-226 Radium-228 -- Thorium-228 -- Thorium-232 Uranium-234 Uranium-235 Uranium-238	SEMIVOLATILES Benzo(A)Anthracene -- -- -- -- -- -- -- Fluorene Indeno(1,2,3-cd)pyrene Phenanthrene -- VOLATILES -- -- -- -- RADIOLOGICAL -- Radium-226 Radium-228 -- Thorium-228 -- Thorium-232 Uranium-234 Uranium-235 Uranium-238

Table 3-7
Identification of Constituents of Potential Concern
Process Summary - Surface Water - STP River
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in Surface Water	Constituents preceded by " → " were removed because they are considered essential nutrients (Ca, Mg, K, Na)	Constituents preceded by " → " were removed due to low frequency of detection (<5% of samples)	Constituents preceded by " → " result of background comparison (Max < Bkgd)	Constituents preceded by " → " were removed because no applicable screening criteria exist	Constituents preceded by " → " were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in Surface Water
INORGANICS Aluminum Antimony Arsenic Barium Beryllium Calcium Chromium, Total Iron Lead Magnesium Manganese Nickel Potassium Selenium Sodium Thallium Vanadium Zinc	INORGANICS Aluminum Antimony Arsenic Barium Beryllium → Calcium Chromium, Total Iron Lead → Magnesium Manganese Nickel → Potassium Selenium → Sodium Thallium Vanadium Zinc	INORGANICS Aluminum Antimony Arsenic Barium Beryllium -- Chromium, Total Iron Lead -- Manganese Nickel -- Selenium -- Thallium Vanadium Zinc	INORGANICS Aluminum Antimony Arsenic Barium Beryllium -- → Chromium, Total Iron Lead -- Manganese Nickel -- → Selenium -- Thallium Vanadium Zinc	INORGANICS Aluminum Antimony Arsenic Barium Beryllium -- -- Iron Lead -- -- Manganese Nickel -- -- Thallium Vanadium Zinc	INORGANICS Aluminum → Antimony Arsenic Barium → Beryllium -- -- Iron Lead -- -- Manganese Nickel -- -- → Thallium → Vanadium Zinc	INORGANICS -- -- Arsenic -- -- -- -- Iron -- -- Manganese Nickel -- -- Thallium -- Zinc
RADIOLOGICAL Radium-226 Radium-228 Thorium-227 Thorium-228 Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	RADIOLOGICAL Radium-226 Radium-228 Thorium-227 Thorium-228 Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	RADIOLOGICAL Radium-226 Radium-228 Thorium-227 Thorium-228 Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	RADIOLOGICAL Radium-226 Radium-228 Thorium-227 Thorium-228 Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	RADIOLOGICAL Radium-226 Radium-228 → Thorium-227 → Thorium-228 → Thorium-230 Thorium-232 Uranium-234 Uranium-235 Uranium-238	RADIOLOGICAL Radium-226 Radium-228 -- -- -- Thorium-232 Uranium-234 Uranium-235 Uranium-238	RADIOLOGICAL Radium-226 Radium-228 -- -- -- Thorium-232 Uranium-234 Uranium-235 Uranium-238

Table 3-8
Identification of Constituents of Potential Concern
Process Summary - STP Upland Soil
Krass Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in On-site Soil	Constituents preceded by " → " were removed because they are considered essential nutrients (Ca, Mg, K, Na)	Constituents preceded by " → " were removed due to low frequency of detection (<5% of samples)	Constituents preceded by " → " result of background comparison (Max < Bkgd)	Constituents preceded by " → " were removed because no applicable screening criteria exist	Constituents preceded by " → " were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in On-site Soil
INORGANICS Aluminum Antimony Arsenic Barium Beryllium Cadmium Calcium Chromium, Total Cobalt Copper Iron Lead Magnesium Manganese Mercury Nickel Potassium Selenium Silver Sodium Thallium Vanadium Zinc	INORGANICS Aluminum Antimony Arsenic Barium Beryllium Cadmium → Calcium Chromium, Total Cobalt Copper Iron Lead → Magnesium Manganese Mercury Nickel → Potassium Selenium Silver → Sodium Thallium Vanadium Zinc	INORGANICS Aluminum Antimony Arsenic Barium Beryllium Cadmium -- Chromium, Total Cobalt Copper Iron Lead -- Manganese Mercury Nickel -- Selenium Silver -- Thallium Vanadium Zinc	INORGANICS → Aluminum Antimony Arsenic Barium Beryllium Cadmium -- Chromium, Total Cobalt Copper Iron Lead -- Manganese Mercury Nickel -- Selenium Silver -- Vanadium Zinc	INORGANICS -- Antimony Arsenic Barium Beryllium Cadmium -- Chromium, Total Cobalt Copper Iron Lead -- Manganese Mercury Nickel -- Selenium Silver -- Vanadium Zinc	INORGANICS -- → Antimony Arsenic Barium → Beryllium → Cadmium -- Chromium, Total Cobalt Copper Iron Lead -- Manganese Mercury Nickel -- Selenium Silver -- Vanadium Zinc	INORGANICS -- -- Arsenic Barium -- -- -- Chromium Cobalt Copper Iron Lead -- Manganese Mercury Nickel -- Selenium Silver -- Vanadium Zinc

Table 3-8
Identification of Constituents of Potential Concern
Process Summary - STP Upland Soil
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in On-site Soil	Constituents preceded by * → * were removed because they are considered essential nutrients (Ca, Mg, K, Na)	Constituents preceded by * → * were removed due to low frequency of detection (<5% of samples)	Constituents preceded by * → * result of background comparison (Max < Bkgd)	Constituents preceded by * → * were removed because no applicable screening criteria exist	Constituents preceded by * → * were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in On-site Soil
SEMIVOLATILES	SEMIVOLATILES	SEMIVOLATILES	SEMIVOLATILES	SEMIVOLATILES	SEMIVOLATILES	SEMIVOLATILES
4-Nitrophenol	4-Nitrophenol	4-Nitrophenol	4-Nitrophenol	--	--	--
Acenaphthylene	Acenaphthylene	→ Acenaphthylene	--	--	--	--
Anthracene	Anthracene	Anthracene	Anthracene	Anthracene	Anthracene	Anthracene
Benzo(A)Anthracene	Benzo(A)Anthracene	Benzo(A)Anthracene	Benzo(A)Anthracene	Benzo(A)Anthracene	Benzo(A)Anthracene	Benzo(A)Anthracene
Benzo(A)Pyrene	Benzo(A)Pyrene	Benzo(A)Pyrene	Benzo(A)Pyrene	Benzo(A)Pyrene	Benzo(A)Pyrene	Benzo(A)Pyrene
Benzo(B)Fluoranthene	Benzo(B)Fluoranthene	Benzo(B)Fluoranthene	Benzo(B)Fluoranthene	Benzo(B)Fluoranthene	Benzo(B)Fluoranthene	Benzo(B)Fluoranthene
Benzo(G,H,I)Perylene	Benzo(G,H,I)Perylene	Benzo(G,H,I)Perylene	Benzo(G,H,I)Perylene	Benzo(G,H,I)Perylene	Benzo(G,H,I)Perylene	Benzo(G,H,I)Perylene
Benzo(K)Fluoranthene	Benzo(K)Fluoranthene	Benzo(K)Fluoranthene	Benzo(K)Fluoranthene	Benzo(K)Fluoranthene	Benzo(K)Fluoranthene	Benzo(K)Fluoranthene
Butylbenzylphthalate	Butylbenzylphthalate	Butylbenzylphthalate	Butylbenzylphthalate	Butylbenzylphthalate	→ Butylbenzylphthalate	--
Bis(2-Ethylhexyl) Phthalate	Bis(2-Ethylhexyl) Phthalate	Bis(2-Ethylhexyl) Phthalate	Bis(2-Ethylhexyl) Phthalate	Bis(2-Ethylhexyl) Phthalate	→ Bis(2-Ethylhexyl)Phthalate	--
Chrysene	Chrysene	Chrysene	Chrysene	Chrysene	Chrysene	Chrysene
Dibenz(A,H)Anthracene	Dibenz(A,H)Anthracene	Dibenz(A,H)Anthracene	Dibenz(A,H)Anthracene	Dibenz(A,H)Anthracene	→ Dibenz(A,H)Anthracene	--
Di-N-Butyl Phthalate	Di-N-Butyl Phthalate	Di-N-Butyl Phthalate	Di-N-Butyl Phthalate	Di-N-Butyl Phthalate	→ Di-N-Butyl Phthalate	--
Di-N-Octylphthalate	Di-N-Octylphthalate	Di-N-Octylphthalate	Di-N-Octylphthalate	Di-N-Octylphthalate	→ Di-N-Octylphthalate	--
Fluoranthene	Fluoranthene	Fluoranthene	Fluoranthene	Fluoranthene	Fluoranthene	Fluoranthene
Fluorene	Fluorene	→ Fluorene	--	--	--	--
Indeno(1,2,3-C,D)Pyrene	Indeno(1,2,3-C,D)Pyrene	Indeno(1,2,3-C,D)Pyrene	Indeno(1,2,3-C,D)Pyrene	Indeno(1,2,3-C,D)Pyrene	Indeno(1,2,3-C,D)Pyrene	Indeno(1,2,3-CD)Pyrene
Phenanthrene	Phenanthrene	Phenanthrene	Phenanthrene	Phenanthrene	Phenanthrene	Phenanthrene
Pyrene	Pyrene	Pyrene	Pyrene	Pyrene	Pyrene	Pyrene
VOLATILES	VOLATILES	VOLATILES	VOLATILES	VOLATILES	VOLATILES	VOLATILES
Acetone	Acetone	→ Acetone	--	--	--	--
Chloroform	Chloroform	Chloroform	Chloroform	→ Chloroform	--	--
Toluene	Toluene	Toluene	Toluene	Toluene	Toluene	Toluene

Table 3-8
 Identification of Constituents of Potential Concern
 Process Summary - STP Upland Soil
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

List of Positively Detected Constituents in On-site Soil	Constituents preceded by " → " were removed because they are considered essential nutrients (Ca, Mg, K, Na)	Constituents preceded by " → " were removed due to low frequency of detection (<5% of samples)	Constituents preceded by " → " result of background comparison (Max < Bkgd)	Constituents preceded by " → " were removed because no applicable screening criteria exist	Constituents preceded by " → " were removed based on the result of screening level comparison (Max < SL)	Final Constituents of Potential Concern (COPCs) in On-site Soil
RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL	RADIOLOGICAL
Alpha, Gross	Alpha, Gross	Alpha, Gross	Alpha, Gross	Alpha, Gross	--	--
Bismuth-214	Bismuth-214	Bismuth-214	Bismuth-214	Bismuth-214	--	--
Cesium-137	Cesium-137	Cesium-137	Cesium-137	Cesium-137	--	--
Lead-212	Lead-212	Lead-212	Lead-212	Lead-212	--	--
Lead-214	Lead-214	Lead-214	Lead-214	Lead-214	--	--
Potassium-40	→ Potassium-40	--	--	--	--	--
Protactinium 234	Protactinium 234	Protactinium 234	Protactinium 234	Protactinium 234	--	--
Radium-226	Radium-226	Radium-226	Radium-226	Radium-226	Radium-226	Radium-226
Radium-228	Radium-228	Radium-228	Radium-228	Radium-228	Radium-228	Radium-228
Thallium-208	Thallium-208	Thallium-208	Thallium-208	Thallium-208	--	--
Thorium-227	Thorium-227	Thorium-227	Thorium-227	Thorium-227	--	--
Thorium-228	Thorium-228	Thorium-228	Thorium-228	Thorium-228	Thorium-228	Thorium-228
Thorium-230	Thorium-230	Thorium-230	Thorium-230	Thorium-230	--	--
Thorium-232	Thorium-232	Thorium-232	Thorium-232	Thorium-232	Thorium-232	Thorium-232
Uranium-234	Uranium-234	Uranium-234	Uranium-234	Uranium-234	--	--
Uranium-235	Uranium-235	Uranium-235	Uranium-235	Uranium-235	Uranium-235	Uranium-235
Uranium-238	Uranium-238	Uranium-238	Uranium-238	Uranium-238	Uranium-238	Uranium-238

TABLE 3-9
 Comparison of Maximum Detections with Background Concentrations, Kress Creek
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
Surface Sediment/Floodplain Soil (< 24")								
INORGANICS								
Aluminum	mg/kg	93	93	100%	5.98E+02	1.73E+04	1.32E+04	Yes
Antimony	mg/kg	3	85	4%	5.50E+00	3.03E+01		Yes
Arsenic	mg/kg	86	93	92%	9.50E-01	1.77E+02	2.06E+01	Yes
Barium	mg/kg	93	93	100%	1.34E+01	6.28E+02	1.53E+02	Yes
Beryllium	mg/kg	45	93	48%	2.60E-01	2.40E+00	9.10E-01	Yes
Cadmium	mg/kg	5	93	5%	3.60E-01	2.60E+00		Yes
Calcium	mg/kg	93	93	100%	1.89E+03	1.72E+05	8.79E+04	Yes
Chromium, Total	mg/kg	83	93	89%	1.20E+00	4.34E+01	2.95E+01	Yes
Cobalt	mg/kg	85	93	91%	1.80E+00	2.10E+01	1.40E+01	Yes
Copper	mg/kg	66	93	71%	2.30E+00	2.87E+03	2.46E+01	Yes
Iron	mg/kg	93	93	100%	3.90E+03	4.67E+04	2.64E+04	Yes
Lead	mg/kg	93	93	100%	1.50E+00	3.10E+03	1.23E+02	Yes
Magnesium	mg/kg	93	93	100%	2.00E+03	1.06E+05	5.04E+04	Yes
Manganese	mg/kg	93	93	100%	1.25E+02	2.69E+03	9.97E+02	Yes
Mercury	mg/kg	31	93	33%	6.00E-02	2.00E+00	1.20E-01	Yes
Nickel	mg/kg	74	93	80%	1.20E+00	5.42E+01	2.62E+01	Yes
Potassium	mg/kg	92	93	99%	1.93E+02	3.12E+03	1.72E+03	Yes
Selenium	mg/kg	19	93	20%	6.40E-01	2.40E+00	1.40E+00	Yes
Silver	mg/kg	10	82	12%	8.50E-01	3.00E+00	2.10E+00	Yes
Sodium	mg/kg	78	93	84%	1.12E+02	1.86E+03	6.18E+02	Yes
Thallium	mg/kg	5	93	5%	4.10E-01	8.80E-01		Yes
Vanadium	mg/kg	78	93	84%	4.60E+00	7.03E+01	2.87E+01	Yes
Zinc	mg/kg	89	93	96%	7.60E+00	2.48E+03	1.62E+02	Yes
PESTICIDES/PCBs								
p,p'-DDD	mg/kg	2	6	33%	1.50E-02	2.80E-01		NA
Aroclor 1260	mg/kg	1	6	17%	2.00E+00	2.00E+00		NA

TABLE 3-9

Comparison of Maximum Detections with Background Concentrations, Kress Creek
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
Surface Sediment/Floodplain Soil (< 24")								
SEMIVOLATILES								
1,2,4-Trichlorobenzene	mg/kg	1	6	17%	5.80E-02	5.80E-02		NA
2-Methylnaphthalene	mg/kg	1	6	17%	9.70E-02	9.70E-02		NA
Acenaphthene	mg/kg	1	6	17%	5.10E-02	5.10E-02		NA
Acenaphthylene	mg/kg	1	6	17%	4.80E-01	4.80E-01		NA
Anthracene	mg/kg	3	6	50%	1.10E-01	9.20E-01		NA
Benzo(A)Anthracene	mg/kg	2	6	33%	3.00E-01	1.80E+00		NA
Benzo(A)Pyrene	mg/kg	1	6	17%	6.90E-01	6.90E-01		NA
Benzo(B)Fluoranthene	mg/kg	1	6	17%	1.90E+00	1.90E+00		NA
Benzo(K)Fluoranthene	mg/kg	1	6	17%	4.90E-01	4.90E-01		NA
Bis(2-Ethylhexyl) Phthalate	mg/kg	1	6	17%	7.80E-02	7.80E-02		NA
Carbazole	mg/kg	2	6	33%	5.50E-02	3.20E-01		NA
Chrysene	mg/kg	3	6	50%	7.70E-02	1.60E+00		NA
Dibenzofuran	mg/kg	1	6	17%	2.90E-01	2.90E-01		NA
Di-N-Butyl Phthalate	mg/kg	1	6	17%	1.80E-01	1.80E-01		NA
Fluoranthene	mg/kg	3	6	50%	1.80E-01	2.70E+00		NA
Fluorene	mg/kg	2	6	33%	8.70E-02	7.40E-01		NA
Indeno(1,2,3-C,D)Pyrene	mg/kg	1	6	17%	7.70E-01	7.70E-01		NA
Naphthalene	mg/kg	1	6	17%	1.10E-01	1.10E-01		NA
Phenanthrene	mg/kg	3	6	50%	1.00E-01	2.90E+00		NA
Pyrene	mg/kg	3	6	50%	6.30E-02	1.90E+00		NA

TABLE 3-9
 Comparison of Maximum Detections with Background Concentrations, Kress Creek
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
Surface Sediment/Floodplain Soil (< 24")								
RADIOLOGICAL								
Actinium 228	pCi/g	27	27	100%	9.84E-01	8.40E+02	1.16E+00	Yes
Bismuth-212	pCi/g	28	28	100%	1.05E+00	7.96E+02	1.32E+00	Yes
Bismuth-214	pCi/g	28	28	100%	4.48E-01	1.35E+01	1.05E+00	Yes
Cesium-137	pCi/g	21	21	100%	1.35E-03	7.30E-01	4.99E-01	Yes
Lead 211	pCi/g	5	5	100%	2.64E-03	2.31E+00	9.47E-02	Yes
Lead-212	pCi/g	28	28	100%	8.49E-01	8.01E+02	9.65E-01	Yes
Lead-214	pCi/g	28	28	100%	4.34E-01	1.53E+01	9.80E-01	Yes
Potassium-40	pCi/g	63	63	100%	3.12E+00	4.31E+01	1.76E+01	Yes
Protactinium 234	pCi/g	25	25	100%	6.33E-02	7.32E+01	9.15E-01	Yes
Radium-223	pCi/g	28	28	100%	4.32E-01	3.37E+02	6.63E-01	Yes
Radium-224	pCi/g	28	28	100%	9.11E-01	3.37E+03	1.28E+00	Yes
Radium-226	pCi/g	116	153	76%	3.00E-01	5.36E+01	2.97E+00	Yes
Radium-228	pCi/g	114	114	100%	4.13E-01	6.53E+02	5.39E+00	Yes
Rhenium-226	pCi/g	28	28	100%	4.41E-01	1.33E+01	9.87E-01	Yes
Rhenium-228	pCi/g	28	28	100%	9.48E-01	8.32E+02	1.09E+00	Yes
Rhenium-total	pCi/g	28	28	100%	1.53E+00	8.44E+02	2.08E+00	Yes
Thallium-208	pCi/g	28	28	100%	8.82E-01	8.13E+02	9.51E-01	Yes
Thorium-227	pCi/g	59	89	66%	3.72E-02	3.27E+01	2.36E-01	Yes
Thorium-228	pCi/g	118	118	100%	1.55E-01	9.97E+02	2.68E+00	Yes
Thorium-230	pCi/g	82	90	91%	2.42E-01	1.06E+02	1.24E+00	Yes
Thorium-232	pCi/g	125	125	100%	2.57E-01	6.54E+02	2.30E+00	Yes
Thorium-234	pCi/g	20	20	100%	3.36E-01	8.03E+01	6.25E-01	Yes
Uranium-234	pCi/g	73	90	81%	3.08E-01	4.71E+01	1.32E+00	Yes
Uranium-235	pCi/g	115	115	100%	4.30E-03	4.38E+00	1.90E-01	Yes
Uranium-238	pCi/g	125	125	100%	2.00E-01	4.26E+01	2.10E+00	Yes

TABLE 3-9
 Comparison of Maximum Detections with Background Concentrations, Kress Creek
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
<i>Surface Water</i>								
INORGANICS								
Aluminum	ug/L	12	12	100%	5.19E+02	3.31E+03	3.98E+03	No
Arsenic	ug/L	5	12	42%	1.70E+00	2.70E+00	2.70E+00	No
Barium	ug/L	12	12	100%	6.05E+01	1.07E+02	8.77E+01	Yes
Calcium	ug/L	12	12	100%	4.38E+04	6.99E+04	6.62E+04	Yes
Chromium, Total	ug/L	3	12	25%	5.90E+00	6.70E+00	8.40E+00	No
Cobalt	ug/L	1	12	8%	4.30E+00	4.30E+00		Yes
Copper	ug/L	10	12	83%	7.10E+00	1.45E+01	7.40E+00	Yes
Iron	ug/L	12	12	100%	7.57E+02	4.50E+03	6.07E+03	No
Lead	ug/L	5	12	42%	5.60E+00	1.04E+01	1.03E+01	Yes
Magnesium	ug/L	12	12	100%	1.80E+04	3.27E+04	3.16E+04	Yes
Manganese	ug/L	12	12	100%	5.51E+01	1.51E+02	1.84E+02	No
Nickel	ug/L	1	12	8%	7.00E+00	7.00E+00	5.90E+00	Yes
Potassium	ug/L	12	12	100%	4.57E+03	6.72E+03	1.15E+04	Yes
Selenium	ug/L	1	12	8%	2.80E+00	2.80E+00	3.20E+00	No
Sodium	ug/L	12	12	100%	2.32E+04	7.46E+04	9.63E+04	Yes
Vanadium	ug/L	11	12	92%	2.60E+00	9.00E+00	7.20E+00	Yes
Zinc	ug/L	12	12	100%	4.20E+00	3.34E+01	3.72E+01	No

NA -- Not applicable; no background data for organic constituents

TABLE 3-10
 Comparison of Maximum Detections with Background Concentrations, STP River
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
<i>Surface Sediment/Floodplain Soil (< 24")</i>								
INORGANICS								
Aluminum	mg/kg	39	39	100%	7.34E+02	1.35E+04	1.32E+04	Yes
Antimony	mg/kg	4	31	13%	5.80E+00	1.21E+01		Yes
Arsenic	mg/kg	39	39	100%	1.20E+00	3.16E+01	2.06E+01	Yes
Barium	mg/kg	39	39	100%	1.49E+01	2.03E+02	1.53E+02	Yes
Beryllium	mg/kg	2	39	5%	7.80E-01	1.20E+00	9.10E-01	Yes
Cadmium	mg/kg	6	39	15%	1.90E-01	7.50E-01		
Calcium	mg/kg	39	39	100%	1.50E+03	1.39E+05	8.79E+04	Yes
Chromium, Total	mg/kg	30	39	77%	2.90E+00	3.69E+01	2.95E+01	Yes
Cobalt	mg/kg	32	39	82%	1.70E+00	1.95E+01	1.40E+01	Yes
Copper	mg/kg	32	39	82%	3.50E+00	7.93E+01	2.46E+01	Yes
Iron	mg/kg	39	39	100%	3.70E+03	5.17E+04	2.64E+04	Yes
Lead	mg/kg	39	39	100%	2.10E+00	6.40E+01	1.23E+02	No
Magnesium	mg/kg	39	39	100%	1.94E+03	8.79E+04	5.04E+04	Yes
Manganese	mg/kg	39	39	100%	1.70E+02	1.63E+03	9.97E+02	Yes
Mercury	mg/kg	8	39	21%	9.20E-02	3.70E+00	1.20E-01	Yes
Nickel	mg/kg	39	39	100%	1.90E+00	2.81E+01	2.62E+01	Yes
Potassium	mg/kg	39	39	100%	1.84E+02	1.30E+03	1.72E+03	No
Selenium	mg/kg	2	39	5%	8.30E-01	1.20E+00	1.40E+00	No
Silver	mg/kg	7	39	18%	1.10E+00	2.40E+00	2.10E+00	Yes
Sodium	mg/kg	30	39	77%	4.60E+01	7.94E+02	6.18E+02	Yes
Thallium	mg/kg	2	39	5%	4.00E-01	6.70E-01		NA
Vanadium	mg/kg	39	39	100%	3.90E+00	2.95E+01	2.87E+01	Yes
Zinc	mg/kg	36	39	92%	9.60E+00	2.07E+02	1.62E+02	Yes

TABLE 3-10
 Comparison of Maximum Detections with Background Concentrations, STP River
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
Surface Sediment/Floodplain Soil (< 24")								
SEMIVOLATILES								
Benzo(A)Anthracene	mg/kg	2	3	67%	1.50E-01	1.70E-01		NA
Benzo(A)Pyrene	mg/kg	2	3	67%	6.60E-02	9.90E-02		NA
Benzo(B)Fluoranthene	mg/kg	2	3	67%	1.40E-01	1.50E-01		NA
Benzo(G,H,I)Perylene	mg/kg	1	2	50%	7.60E-02	7.60E-02		NA
Benzo(K)Fluoranthene	mg/kg	2	3	67%	1.20E-01	1.50E-01		NA
Carbazole	mg/kg	1	2	50%	4.70E-02	4.70E-02		NA
Chrysene	mg/kg	2	3	67%	1.70E-01	1.80E-01		NA
Fluoranthene	mg/kg	2	3	67%	4.60E-01	5.60E-01		NA
Fluorene	mg/kg	1	2	50%	5.00E-02	5.00E-02		NA
Indeno(1,2,3-C,D)Pyrene	mg/kg	2	3	67%	7.80E-02	7.90E-02		NA
Phenanthrene	mg/kg	2	3	67%	3.70E-01	4.10E-01		NA
Pyrene	mg/kg	2	3	67%	4.10E-01	4.60E-01		NA
VOLATILES								
2-Butanone	mg/kg	1	3	33%	4.80E-02	4.80E-02		NA
Acetone	mg/kg	1	3	33%	2.60E-01	2.60E-01		NA
Carbon Disulfide	mg/kg	1	3	33%	4.00E-03	4.00E-03		NA
Toluene	mg/kg	3	3	100%	3.00E-03	1.70E-02		NA
RADIONUCLIDES								
Potassium-40	pCi/g	8	8	100%	5.80E+00	1.84E+01	1.76E+01	Yes
Radium-226	pCi/g	33	47	70%	6.00E-01	4.80E+00	2.97E+00	Yes
Radium-228	pCi/g	43	43	100%	2.43E-01	1.09E+02	5.39E+00	Yes
Thorium-227	pCi/g	38	38	100%	9.66E-03	1.31E+01	2.36E-01	Yes
Thorium-228	pCi/g	39	39	100%	2.11E-01	1.03E+02	2.68E+00	Yes
Thorium-230	pCi/g	39	39	100%	2.75E-01	1.46E+01	1.24E+00	Yes
Thorium-232	pCi/g	47	47	100%	1.85E-01	9.92E+01	2.30E+00	Yes
Uranium-234	pCi/g	38	39	97%	2.43E-01	9.60E+00	1.32E+00	Yes
Uranium-235	pCi/g	38	38	100%	5.16E-03	2.83E-01	1.90E-01	Yes
Uranium-238	pCi/g	47	47	100%	2.48E-01	6.00E+00	2.10E+00	Yes

TABLE 3-10
 Comparison of Maximum Detections with Background Concentrations, STP River
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
Surface Water								
INORGANICS								
Aluminum	ug/L	6	6	100%	8.83E+02	4.50E+03	3.98E+03	Yes
Antimony	ug/L	1	6	17%	1.13E+01	1.13E+01		NA
Arsenic	ug/L	5	6	83%	1.90E+00	3.90E+00	2.70E+00	Yes
Barium	ug/L	6	6	100%	6.78E+01	9.30E+01	8.77E+01	Yes
Beryllium	ug/L	3	6	50%	3.70E-01	4.50E-01	3.70E-01	Yes
Calcium	ug/L	6	6	100%	3.84E+04	6.71E+04	6.62E+04	Yes
Chromium, Total	ug/L	1	6	17%	4.90E+00	4.90E+00	8.40E+00	Yes
Iron	ug/L	6	6	100%	1.33E+03	6.72E+03	6.07E+03	Yes
Lead	ug/L	6	6	100%	3.60E+00	1.31E+01	1.03E+01	Yes
Magnesium	ug/L	6	6	100%	1.60E+04	2.77E+04	3.16E+04	Yes
Manganese	ug/L	6	6	100%	4.14E+01	1.52E+02	1.34E+02	Yes
Nickel	ug/L	3	6	50%	9.20E+00	1.18E+01	5.90E+00	Yes
Potassium	ug/L	6	6	100%	6.06E+03	1.25E+04	1.15E+04	Yes
Selenium	ug/L	1	6	17%	2.60E+00	2.60E+00	3.20E+00	Yes
Sodium	ug/L	6	6	100%	3.64E+04	1.38E+05	9.63E+04	Yes
Thallium	ug/L	1	6	17%	2.20E+00	2.20E+00		NA
Vanadium	ug/L	6	6	100%	3.20E+00	8.70E+00	7.20E+00	Yes
Zinc	ug/L	3	6	50%	4.12E+01	4.86E+01	3.72E+01	Yes
RADIONUCLIDES								
Radium-226	pCi/L	3	6	50%	6.46E-01	8.22E-01		NA
Radium-228	pCi/L	6	6	100%	2.57E-02	1.25E+00	5.90E-01	Yes
Thorium-227	pCi/L	6	6	100%	7.50E-03	5.56E-02	2.65E-02	Yes
Thorium-228	pCi/L	5	5	100%	4.35E-02	2.02E-01		NA
Thorium-230	pCi/L	6	6	100%	5.43E-02	3.22E-01	2.39E-02	Yes
Thorium-232	pCi/L	6	6	100%	2.10E-02	1.73E-01	8.05E-03	Yes
Uranium-234	pCi/L	6	6	100%	5.79E-02	5.72E-01	5.54E-01	Yes
Uranium-235	pCi/L	6	6	100%	3.26E-03	3.64E-02	3.46E-02	Yes
Uranium-238	pCi/L	6	6	100%	1.88E-02	5.82E-01	3.60E-01	Yes

NA -- Not applicable; no background data exist

TABLE 3-11

Comparison of Maximum Detections with Background Concentrations, STP Upland Soil
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
INORGANICS								
Aluminum	mg/kg	161	161	100%	1.35E+03	1.91E+04	2.00E+04	No
Antimony	mg/kg	8	121	7%	4.10E+00	1.20E+01	1.10E+01	Yes
Arsenic	mg/kg	161	161	100%	2.80E+00	9.07E+01	2.92E+01	Yes
Barium	mg/kg	161	161	100%	8.10E+00	5.14E+02	2.08E+02	Yes
Beryllium	mg/kg	28	161	17%	4.30E-01	2.20E+00		Yes
Cadmium	mg/kg	18	161	11%	5.30E-01	3.40E+00	1.10E+00	Yes
Calcium	mg/kg	161	161	100%	1.74E+03	1.59E+05	1.58E+05	Yes
Chromium, Total	mg/kg	161	161	100%	2.30E+00	6.89E+01	2.51E+01	Yes
Cobalt	mg/kg	155	161	96%	2.30E+00	3.50E+01	1.45E+01	Yes
Copper	mg/kg	161	161	100%	6.20E+00	5.91E+02	2.62E+01	Yes
Iron	mg/kg	161	161	100%	5.64E+03	6.75E+04	4.98E+04	Yes
Lead	mg/kg	161	161	100%	4.40E+00	1.16E+03	5.37E+01	Yes
Magnesium	mg/kg	161	161	100%	2.23E+03	9.52E+04	9.54E+04	Yes
Manganese	mg/kg	161	161	100%	1.80E+02	2.14E+03	1.26E+03	Yes
Mercury	mg/kg	66	161	41%	6.30E-02	2.70E+00		Yes
Nickel	mg/kg	151	161	94%	7.00E+00	4.71E+01	3.61E+01	Yes
Potassium	mg/kg	151	161	94%	5.09E+02	3.01E+03	2.02E+03	Yes
Selenium	mg/kg	11	161	7%	6.00E-01	1.50E+00		Yes
Silver	mg/kg	10	161	6%	9.40E-01	1.97E+01		Yes
Sodium	mg/kg	97	161	60%	7.54E+01	7.25E+02	3.84E+02	Yes
Thallium	mg/kg	2	161	1%	8.10E-01	2.20E+00		Yes
Vanadium	mg/kg	161	161	100%	4.80E+00	4.17E+01	4.03E+01	Yes
Zinc	mg/kg	161	161	100%	1.45E+01	1.40E+03	1.05E+02	Yes

TABLE 3-11

Comparison of Maximum Detections with Background Concentrations, STP Upland Soil
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
SEMIVOLATILES								
4-Nitrophenol	mg/kg	1	22	5%	6.40E-02	6.40E-02	4.70E-02	Yes
Acenaphthylene	mg/kg	1	22	5%	4.80E-02	4.80E-02		NA
Anthracene	mg/kg	2	22	9%	4.80E-02	1.30E-01		NA
Benzo(A)Anthracene	mg/kg	10	22	45%	5.60E-02	4.60E-01	5.30E-02	Yes
Benzo(A)Pyrene	mg/kg	5	22	23%	4.70E-02	3.10E-01		NA
Benzo(B)Fluoranthene	mg/kg	9	22	41%	5.70E-02	7.40E-01	9.40E-02	Yes
Benzo(G,H,I)Perylene	mg/kg	2	22	9%	2.00E-01	2.40E-01		NA
Benzo(K)Fluoranthene	mg/kg	7	22	32%	4.70E-02	7.40E-01	9.40E-02	Yes
Benzyl Butyl Phthalate	mg/kg	3	22	14%	4.30E-02	5.30E-02		NA
Bis(2-Ethylhexyl) Phthalate	mg/kg	7	22	32%	3.00E-01	1.50E+00	2.90E+00	No
Chrysene	mg/kg	8	22	36%	4.80E-02	3.60E-01	4.80E-02	Yes
Dibenz(A,H)Anthracene	mg/kg	2	22	9%	6.00E-02	7.20E-02		NA
Di-N-Butyl Phthalate	mg/kg	2	22	9%	4.50E-02	5.20E-02	7.20E-02	No
Di-N-Octylphthalate	mg/kg	3	22	14%	8.90E-02	2.00E-01	5.00E-02	Yes
Fluoranthene	mg/kg	10	22	45%	8.40E-02	9.20E-01	8.30E-02	Yes
Fluorene	mg/kg	1	22	5%	5.70E-02	5.70E-02		NA
Indeno(1,2,3-C,D)Pyrene	mg/kg	2	22	9%	1.80E-01	2.80E-01		NA
Phenanthrene	mg/kg	10	22	45%	4.50E-02	4.40E-01	4.20E-02	Yes
Pyrene	mg/kg	10	22	45%	6.50E-02	6.30E-01	7.10E-02	Yes
VOLATILES								
Acetone	mg/kg	1	22	5%	1.90E-02	1.90E-02		NA
Chloroform	mg/kg	3	22	14%	1.00E-03	2.00E-03		NA
Toluene	mg/kg	18	22	82%	1.00E-03	5.20E-01	1.00E-01	Yes

TABLE 3-11

Comparison of Maximum Detections with Background Concentrations, STP Upland Soil
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detections	Number of Analyses	Frequency of Detection	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
RADIOLOGICAL								
Alpha, Gross	pCi/g	10	10	100%	4.49E+00	1.88E+02		NA
Bismuth-214	pCi/g	10	10	100%	7.60E-01	1.50E+00		NA
Cesium-137	pCi/g	6	6	100%	2.92E-02	1.47E-01		NA
Lead-212	pCi/g	10	10	100%	9.73E-01	1.38E+01		NA
Lead-214	pCi/g	10	10	100%	8.13E-01	1.60E+00		NA
Potassium-40	pCi/g	10	10	100%	1.13E+01	1.84E+01		NA
Protactinium 234	pCi/g	2	2	100%	2.19E+00	2.26E+00		NA
Radium-226	pCi/g	144	156	92%	5.58E-01	9.86E+00	1.73E+00	Yes
Radium-228	pCi/g	149	156	96%	4.42E-01	4.46E+02	4.99E+00	Yes
Thallium-208	pCi/g	10	10	100%	3.07E-01	4.68E+00		NA
Thorium-227	pCi/g	118	142	83%	1.05E-02	3.83E+01	1.12E-01	Yes
Thorium-228	pCi/g	146	146	100%	2.27E-01	3.35E+02	1.33E+00	Yes
Thorium-230	pCi/g	144	146	99%	4.26E-01	1.06E+02	1.25E+00	Yes
Thorium-232	pCi/g	146	146	100%	2.25E-01	3.02E+02	1.73E+00	Yes
Uranium-234	pCi/g	136	146	93%	4.02E-01	1.28E+01	1.01E+00	Yes
Uranium-235	pCi/g	142	148	96%	5.32E-03	8.11E-01	1.26E-01	Yes
Uranium-238	pCi/g	143	146	98%	3.76E-01	1.20E+01	1.19E+00	Yes

NA -- Not applicable; no background data exist

TABLE 3-12
 Comparison of Maximum Detections with Background Concentrations, Fish Tissues
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Analyte	Units	Number of Detection	Number of Sample	FOD	Minimum Detected Concentration	Maximum Detected Concentration	Maximum Background Concentration	Exceed Maximum Background Level?
INORGANICS								
Arsenic	mg/kg	1	15	1 / 15	3.50E-01	3.50E-01	5.00E-02	Yes
Barium	mg/kg	1	15	1 / 15	1.80E+00	1.80E+00	4.00E+00	No
Calcium	mg/kg	7	15	7 / 15	9.13E+02	1.15E+04	2.09E+04	--
Chromium, Total	mg/kg	1	15	1 / 15	1.10E+00	1.10E+00	4.80E-01	Yes
Iron	mg/kg	1	15	1 / 15	4.70E+00	4.70E+00	1.38E+01	No
Lead	mg/kg	3	15	3 / 15	1.70E-01	3.80E-01	2.00E-01	Yes
Magnesium	mg/kg	15	15	15 / 15	2.46E+02	4.54E+02	5.46E+02	No
Mercury	mg/kg	10	15	10 / 15	8.00E-02	2.20E-01	1.00E-01	Yes
Nickel	mg/kg	1	15	1 / 15	1.64E+02	1.64E+02	1.20E-01	Yes
Potassium	mg/kg	15	15	15 / 15	2.77E+03	3.95E+03	3.89E+03	--
Selenium	mg/kg	7	15	7 / 15	5.90E-01	9.60E-01	2.10E-01	Yes
Sodium	mg/kg	15	15	15 / 15	4.58E+02	9.11E+02	1.26E+03	--
Thallium	mg/kg	1	15	1 / 15	2.60E-01	2.60E-01	2.90E-02	Yes
RADIOLOGICAL								
Radium-228	pCi/g	12	12	12 / 12	5.36E-03	1.34E-01	1.03E-01	Yes
Thorium-227	pCi/g	11	11	11 / 11	1.11E-04	2.41E-02	4.80E-03	Yes
Thorium-228	pCi/g	12	12	12 / 12	4.36E-04	2.69E-02	1.37E-03	Yes
Thorium-232	pCi/g	12	12	12 / 12	3.29E-04	4.91E-03	1.37E-03	Yes
Uranium-234	pCi/g	1	12	1 / 12	4.61E-03	4.61E-03	5.16E-03	No
Uranium-235	pCi/g	9	9	9 / 9	3.43E-05	1.37E-03	2.65E-03	No
Uranium-238	pCi/g	5	12	5 / 12	6.36E-04	6.51E-03	6.27E-03	Yes

Note:

-- Indicates an essential nutrient and therefore, background comparison is irrelevant

TABLE 3-13

Constituents Not Evaluated Quantitatively for Ecological Risk (No Benchmarks)
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Site Location	Medium	Units	Constituent	Detection Frequency	Maximum Detection	Arithmetic Mean
KCK	Sediment	mg/kg	Beryllium	45 / 93	2.40E+00	6.99E-01
		mg/kg	Cobalt	85 / 93	2.10E+01	7.59E+00
		mg/kg	Selenium	19 / 93	2.40E+00	1.13E+00
		mg/kg	Thallium	5 / 93	8.80E-01	6.20E-01
		mg/kg	Vanadium	78 / 93	7.03E+01	2.38E+01
		pCi/g	Actinium-228	27 / 27	8.40E+02	9.35E+01
		pCi/g	Bismuth-212	28 / 28	7.96E+02	8.66E+01
		pCi/g	Bismuth-214	28 / 28	1.35E+01	2.56E+00
		pCi/g	Lead-211	5 / 5	2.31E+00	5.30E-01
		pCi/g	Lead-212	28 / 28	8.01E+02	8.60E+01
		pCi/g	Lead-214	28 / 28	1.53E+01	2.79E+00
		pCi/g	Potassium-40	63 / 63	4.31E+01	1.30E+01
		pCi/g	Proactinium-234	25 / 25	7.32E+01	1.06E+01
		pCi/g	Radium-223	28 / 28	3.37E+02	3.46E+01
		pCi/g	Radium-224	28 / 28	3.37E+03	3.61E+02
		pCi/g	Rhenium-226	28 / 28	1.33E+01	2.68E+00
		pCi/g	Rhenium-228	28 / 28	8.32E+02	8.80E+01
		pCi/g	Thallium-208	28 / 28	8.13E+02	8.48E+01
		pCi/g	Thorium-227	59 / 89	3.27E+01	1.64E+00
		pCi/g	Thorium-228	118 / 118	9.97E+02	4.94E+01
pCi/g	Thorium-230	82 / 90	1.06E+02	4.03E+00		
pCi/g	Thorium-234	20 / 20	8.03E+01	1.23E+01		
mg/kg	Carbazole	2 / 6	3.20E-01	1.88E-01		
STP River	Sediment	mg/kg	Barium	39 / 39	2.03E+02	9.34E+01
		mg/kg	Beryllium	2 / 39	1.20E+00	9.90E-01
		mg/kg	Cobalt	39 / 39	1.95E+01	8.33E+00
		mg/kg	Selenium	2 / 39	1.20E+00	1.02E+00
		mg/kg	Thallium	2 / 39	6.70E-01	5.35E-01
		mg/kg	Vanadium	39 / 39	2.95E+01	1.45E+01
		pCi/g	Potassium-40	9 / 9	1.84E+01	1.41E+01
		pCi/g	Thorium-227	38 / 38	1.31E+01	5.13E-01
		pCi/g	Thorium-228	39 / 39	1.03E+02	5.16E+00
		pCi/g	Thorium-230	39 / 39	1.46E+01	1.49E+00
		mg/kg	2-Butanone	1 / 3	4.80E-02	NA
		mg/kg	Acetone	1 / 3	2.60E-01	NA
		mg/kg	Benzo(g,h,i)perylene	1 / 2	7.60E-02	NA
		mg/kg	Benzo(k)fluoranthene	2 / 3	1.50E-01	1.35E-01
		mg/kg	Carbazole	1 / 2	4.70E-02	NA
		mg/kg	Carbon disulfide	1 / 3	4.00E-03	NA

TABLE 3-13

Constituents Not Evaluated Quantitatively for Ecological Risk (No Benchmarks)

Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Site Location	Medium	Units	Constituent	Detection Frequency	Maximum Detection	Arithmetic Mean
STP River	Surface Water	pCi/L	Thorium-227	6 / 6	5.60E-02	3.14E-02
		pCi/L	Thorium-228	5 / 5	2.02E-01	1.19E-01
		pCi/L	Thorium-230	6 / 6	3.22E-01	1.80E-01
STP Upland	Soil	pCi/g	Bismuth-214	10 / 10	1.50E+00	1.00E+00
		pCi/g	Lead-212	10 / 10	1.38E+01	3.89E+00
		pCi/g	Lead-214	10 / 10	1.60E+00	1.09E+00
		pCi/g	Potassium-40	10 / 10	1.84E+01	1.54E+01
		pCi/g	Proactinium-234	2 / 2	2.26E+00	2.23E+00
		pCi/g	Thallium-208	10 / 10	4.68E+00	1.29E+00
		pCi/g	Thorium-227	118 / 142	3.83E+01	6.33E-01
		pCi/g	Thorium-228	146 / 146	3.35E+02	6.57E+00
		pCi/g	Thorium-230	144 / 146	1.06E+02	2.89E+00
		mg/kg	Chloroform	3 / 22	2.00E-03	1.33E-03

TABLE 3-14
 Summary of Radiological Parameters
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Constituent	Ra-226	Ra-228	Th-227	Th-228	Th-230	Th-232	U-234	U-235	U-238
Half Life (yr)	1,600	5.7	18.7	1.9	77,000	14 billion	244,000	7×10^8	4.5×10^9
Nuclide Parent	U-238	Th-232		Ac-228	U-234	NA	Pa-234	NA	NA
Decay Mode	α, γ	β, γ		α	α	α	α	α, γ	α, γ
Distribution Coefficient	250	250	60,000	60,000	60,000	60,000	45	45	45

From: Languir, D., Aqueous Environmental Geochemistry: Prentice Hall, New Jersey. 1997 and Ozuntali, O. I., and G. W. Roles. De Minimus Waste Impacts Analysis Methodology. U. S. Nuclear Regulatory Commission. NUREG/CR-3585. 1984.

TABLE 3-15
 Distribution Coefficients for Inorganic Constituents Detected in KCK/STP Media
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical Name	K_{oc}/K_d	Ref	Chemical Name	K_{oc}/K_d	Ref
Metals					
Aluminum			Lead	4.5-7640	2
Antimony	45-550	1	Manganese	0.2-10,000	2
Arsenic	1-18	2	Nickel	0.2-929	3
Barium	60-16,000	1	Selenium	1.2-8.6	2
Cobalt	0.2-3800	3	Thallium	2,000-510,000	3
Copper	1.4-333	3	Vanadium	50-1000	1
Iron	1.4-1000	2	Zinc	0.1-8000	2

- 1 USEPA. Soil Screening Guidance: Technical Background Document, Office of Solid Waste and Emergency Response. EPA/540/R-95/128. May 1996.
- 2 C. Baes and R. Sharp. "A Proposal for Estimation of Soil Leaching and Leaching Constants for Use in Assessment Models." *Journal of Environmental Quality*. Vol. 12, No. 1, 1983.
- 3 J. Dragun. *The Soil Chemistry of Hazardous Materials*. 2nd ed. Amhurst Scientific Publishers. 1998.

TABLE 4-1
 Bioaccumulative Chemicals List and Log K_{ow} Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Log K_{ow} Range	Selected log K_{ow}	Reference	Evaluate for Food Web Exposures?
Volatile Organics				
1,1,1-Trichloroethane	2.47 to 2.51	2.48	USEPA 1995b	NO
1,1,2,2-Tetrachloroethane	2.31 to 2.64	2.39	USEPA 1995b	NO
1,1,2-Trichloroethane	2.03 to 2.07	2.05	USEPA 1995b	NO
1,1-Dichloroethane	1.78 to 1.85	1.79	USEPA 1995b	NO
1,1-Dichloroethene	2.13 to 2.37	2.13	USEPA 1995b	NO
1,2-Dibromo-3-Chloropropane	2.26 to 2.41	2.34	USEPA 1995b	NO
1,2-Dibromoethane	Not reported	2.00	USEPA 1996a	NO
1,2-Dichloroethane	1.40 to 1.48	1.47	USEPA 1995b	NO
1,2-Dichloropropane	1.94 to 1.99	1.97	USEPA 1995b	NO
2-Butanone	0.26 to 0.69	0.28	USEPA 1995b	NO
2-Hexanone	Not reported	1.40	USEPA 1996a	NO
4-Methyl-2-Pentanone	1.17 to 1.25	1.19	USEPA 1995b	NO
Acetone	-0.21 to -0.24	-0.24	USEPA 1995b	NO
Benzene	1.83 to 2.50	2.13	USEPA 1995b	NO
Bromochloromethane	Not reported	1.41	SRC 1998	NO
Bromodichloromethane	1.88 to 2.14	2.10	USEPA 1995b	NO
Bromoform	2.30 to 2.38	2.35	USEPA 1995b	NO
Bromomethane	1.08 to 1.19	1.19	USEPA 1995b	NO
Carbon disulfide	1.84 to 2.16	2.00	USEPA 1995b	NO
Carbon tetrachloride	2.03 to 3.10	2.73	USEPA 1995b	YES
Chlorobenzene	2.46 to 3.79	2.86	USEPA 1995b	YES
Chloroethane	Not reported	1.43	USEPA 1996a	NO
Chloroform	1.81 to 3.04	1.92	USEPA 1995b	YES
Chloromethane	0.90 to 0.94	0.91	USEPA 1995b	NO
Cis-1,2-Dichloroethene	1.77 to 2.10	1.86	USEPA 1995b	NO
Cis-1,3-Dichloropropene	1.76 to 2.10	2.00	USEPA 1995b	NO
Dibromochloromethane	2.13 to 2.24	2.17	USEPA 1995b	NO
Ethylbenzene	3.07 to 3.57	3.14	USEPA 1995b	YES
Methylene chloride	1.22 to 1.40	1.25	USEPA 1995b	NO
Styrene	2.76 to 3.16	2.94	USEPA 1995b	YES
Tetrachloroethene	2.53 to 3.70	2.67	USEPA 1995b	YES
Toluene	2.21 to 3.13	2.75	USEPA 1995b	YES
Trans-1,2-Dichloroethene	1.77 to 2.10	2.07	USEPA 1995b	NO
Trans-1,3-Dichloropropene	1.76 to 2.10	2.00	USEPA 1995b	NO
Trichloroethene	2.53 to 3.14	2.71	USEPA 1995b	YES
Vinyl chloride	1.23 to 1.52	1.50	USEPA 1995b	NO
Xylenes (total)	2.77 to 3.68	3.20	USEPA 1995b	YES

US EPA ARCHIVE DOCUMENT

TABLE 4-1
 Bioaccumulative Chemicals List and Log K_{ow} Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Log K_{ow} Range	Selected log K_{ow}	Reference	Evaluate for Food Web Exposures?
Semivolatile Organics				
1,2,4-Trichlorobenzene	3.89 to 4.23	4.01	USEPA 1995b	YES
1,2-Dichlorobenzene	3.20 to 3.61	3.43	USEPA 1995b	YES
1,3-Dichlorobenzene	Not reported	3.50	USEPA 1996a	YES
1,4-Dichlorobenzene	3.26 to 3.78	3.42	USEPA 1995b	YES
2,2'-Oxybis(1-Chloropropane)	Not reported	2.50	USEPA 1996a	NO
2,4,5-Trichlorophenol	2.39 to 4.19	3.90	USEPA 1995b	YES
2,4,6-Trichlorophenol	3.29 to 4.05	3.70	USEPA 1995b	YES
2,4-Dichlorophenol	2.80 to 3.30	3.08	USEPA 1995b	YES
2,4-Dimethylphenol	1.99 to 2.49	2.36	USEPA 1995b	NO
2,4-Dinitrophenol	1.40 to 1.79	1.55	USEPA 1995b	NO
2,4-Dinitrotoluene	1.98 to 2.05	2.01	USEPA 1995b	NO
2,6-Dinitrotoluene	1.72 to 2.03	1.87	USEPA 1995b	NO
2-Chloronaphthalene	Not reported	4.10	USEPA 1996a	YES
2-Chlorophenol	0.83 to 2.32	2.15	USEPA 1995b	NO
2-Methylnaphthalene	Not reported	3.90	USEPA 1996a	YES
2-Methylphenol	1.90 to 2.04	1.99	USEPA 1995b	NO
2-Nitroaniline	Not reported	1.90	USEPA 1996a	NO
2-Nitrophenol	Not reported	1.80	USEPA 1996a	NO
3,3'-Dichlorobenzidine	3.51 to 3.95	3.51	USEPA 1995b	YES
3-Nitroaniline	Not reported	1.40	USEPA 1996a	NO
4,6-Dinitro-2-Methylphenol	Not reported	2.10	USEPA 1996a	NO
4-Bromophenyl-Phenylether	4.89 to 5.24	5.00	USEPA 1995b	YES
4-Chloro-3-Methylphenol	Not reported	3.10	USEPA 1996a	YES
4-Chloroaniline	1.57 to 2.02	1.85	USEPA 1995b	NO
4-Chlorophenyl-Phenylether	4.08 to 5.09	4.95	USEPA 1995b	YES
4-Methylphenol	1.38 to 2.04	1.95	USEPA 1995b	NO
4-Nitroaniline	Not reported	1.40	USEPA 1996a	NO
4-Nitrophenol	Not reported	1.90	USEPA 1996a	NO
Acenaphthene	3.77 to 4.49	3.92	USEPA 1995b	YES
Acenaphthylene	Not reported	4.10	USEPA 1996a	YES
Anthracene	3.45 to 4.80	4.55	USEPA 1995b	YES
Benzo(a)anthracene	4.00 to 5.79	5.70	USEPA 1995b	YES

TABLE 4-1
 Bioaccumulative Chemicals List and Log K_{ow} Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Log K_{ow} Range	Selected log K_{ow}	Reference	Evaluate for Food Web Exposures?
Semivolatile Organics				
Benzo(a)pyrene	5.98 to 6.42	6.11	USEPA 1995b	YES
Benzo(b)fluoranthene	5.79 to 6.40	6.20	USEPA 1995b	YES
Benzo(g,h,i)perylene	6.63 to 7.05	6.70	USEPA 1995b	YES
Benzo(k)fluoranthene	6.12 to 6.27	6.20	USEPA 1995b	YES
Bis-(2-Chloroethoxy)methane	Not reported	0.75	USEPA 1996a	NO
Bis-(2-Chloroethyl)ether	1.00 to 1.29	1.21	USEPA 1995b	NO
Bis-(2-Ethylhexyl)phthalate	4.20 to 8.61	7.30	USEPA 1995b	YES
Butylbenzylphthalate	3.57 to 5.02	4.84	USEPA 1995b	YES
Carbazole	3.01 to 3.76	3.59	USEPA 1995b	YES
Chrysene	5.41 to 5.79	5.70	USEPA 1995b	YES
Dibenz(a,h)anthracene	6.50 to 6.88	6.69	USEPA 1995b	YES
Dibenzofuran	Not reported	4.20	USEPA 1996a	YES
Diethylphthalate	1.40 to 3.00	2.50	USEPA 1995b	YES
Dimethylphthalate	1.34 to 1.90	1.57	USEPA 1995b	NO
Di-n-butylphthalate	3.74 to 4.79	4.61	USEPA 1995b	YES
Di-n-octylphthalate	8.03 to 9.49	8.06	USEPA 1995b	YES
Fluoranthene	4.31 to 5.39	5.12	USEPA 1995b	YES
Fluorene	4.04 to 4.40	4.21	USEPA 1995b	YES
Hexachloro-1,3-butadiene	4.74 to 5.16	4.81	USEPA 1995b	YES
Hexachlorobenzene	5.00 to 7.42	5.89	USEPA 1995b	YES
Hexachlorocyclopentadiene	5.04 to 5.51	5.39	USEPA 1995b	YES
Hexachloroethane	3.82 to 4.14	4.00	USEPA 1995b	YES
Indeno(1,2,3-cd)pyrene	6.58 to 6.72	6.65	USEPA 1995b	YES
Isophorone	1.67 to 1.90	1.70	USEPA 1995b	NO
Naphthalene	3.01 to 4.70	3.36	USEPA 1995b	YES
Nitrobenzene	1.70 to 2.93	1.84	USEPA 1995b	NO
N-Nitrosodi-n-propylamine	1.31 to 1.49	1.40	USEPA 1995b	NO
N-Nitrosodiphenylamine	3.13 to 3.45	3.16	USEPA 1995b	YES
Pentachlorophenol	3.29 to 5.24	5.09	USEPA 1995b	YES
Phenanthrene	4.28 to 4.57	4.55	USEPA 1995b	YES
Phenol	0.79 to 1.55	1.48	USEPA 1995b	NO
Pyrene	4.76 to 5.52	5.11	USEPA 1995b	YES

TABLE 4-1
 Bioaccumulative Chemicals List and Log K_{ow} Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Log K_{ow} Range	Selected log K_{ow}	Reference	Evaluate for Food Web Exposures?
Pesticides/PCBs				
4,4'-DDD	4.73 to 6.65	6.10	USEPA 1995b	YES
4,4'-DDE	5.63 to 6.96	6.76	USEPA 1995b	YES
4,4'-DDT	3.98 to 7.01	6.53	USEPA 1995b	YES
Aldrin	5.11 to 7.50	6.50	USEPA 1995b	YES
Alpha-BHC	3.75 to 3.81	3.80	USEPA 1995b	YES
Alpha-Chlordane	5.80 to 6.41	6.32	USEPA 1995b	YES
Aroclor-1016	Not reported	5.60	Sample et al. 1996	YES
Aroclor-1221	Not reported	4.70	Jones et al. 1997	YES
Aroclor-1232	Not reported	5.10	Jones et al. 1997	YES
Aroclor-1242	Not reported	5.60	Jones et al. 1997	YES
Aroclor-1248	Not reported	6.20	Jones et al. 1997	YES
Aroclor-1254	Not reported	6.50	Jones et al. 1997	YES
Aroclor-1260	Not reported	6.80	Jones et al. 1997	YES
Beta-BHC	3.75 to 3.84	3.81	USEPA 1995b	YES
Delta-BHC	Not reported	4.10	USEPA 1996a	YES
Dieldrin	3.63 to 6.20	5.37	USEPA 1995b	YES
Endosulfan I	3.83 to 3.85	3.83	USEPA 1995b	YES
Endosulfan II	4.45 to 4.52	4.52	USEPA 1995b	YES
Endosulfan sulfate	Not reported	3.70	USEPA 1996a	YES
Endrin	2.92 to 5.20	5.06	USEPA 1995b	YES
Endrin aldehyde	--	4.00	USEPA 1995b	YES
Endrin ketone	--	4.00	Endrin aldehyde	YES
Gamma-BHC (Lindane)	3.00 to 4.95	3.73	USEPA 1995b	YES
Gamma-Chlordane	5.80 to 6.41	6.32	USEPA 1995b	YES
Heptachlor	4.93 to 6.26	6.26	USEPA 1995b	YES
Heptachlor epoxide	3.50 to 5.40	5.00	USEPA 1995b	YES
Methoxychlor	3.31 to 5.60	5.08	USEPA 1995b	YES
Toxaphene	3.23 to 5.56	5.50	USEPA 1995b	YES
PCBs (total)	Not reported	6.00	USEPA 1996a	YES

TABLE 4-1
 Bioaccumulative Chemicals List and Log K_{ow} Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Log K_{ow} Range	Selected log K_{ow}	Reference	Evaluate for Food Web Exposures?
Inorganics				
Aluminum	--	--	--	YES
Antimony	--	--	--	YES
Arsenic	--	--	--	YES
Barium	--	--	--	YES
Beryllium	--	--	--	YES
Cadmium	--	--	--	YES
Calcium	--	--	--	NO
Chromium	--	--	--	YES
Cobalt	--	--	--	YES
Copper	--	--	--	YES
Cyanide	--	--	--	NO
Iron	--	--	--	YES
Lead	--	--	--	YES
Magnesium	--	--	--	NO
Manganese	--	--	--	YES
Mercury	--	--	--	YES
Nickel	--	--	--	YES
Potassium	--	--	--	NO
Selenium	--	--	--	YES
Silver	--	--	--	YES
Sodium	--	--	--	NO
Thallium	--	--	--	YES
Vanadium	--	--	--	YES
Zinc	--	--	--	YES

TABLE 4-2
Soil Bioconcentration Factors For Plants, Soil Invertebrates and Small Mammals
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Soil-Plant BCF (dry weight)		Soil-Invertebrate BAF (dry weight)		Soil-Mouse BAF (dry weight)		Soil-Shrew BAF (dry weight)	
	Value	Reference	Value	Reference	Value	Reference	Value	Reference
Inorganics								
Aluminum	0.004	Baes et al. 1984	0.118	Sample et al. 1998a	0.093	Sample et al. 1998b	0.073	Sample et al. 1998b
Antimony	0.200	Baes et al. 1984	0.063	Helmke et al. 1979	--	see text	--	see text
Arsenic	1.103	Bechtel Jacobs 1998a	0.523	Sample et al. 1998a	0.014	Sample et al. 1998b	0.015	Sample et al. 1998b
Barium	0.150	Baes et al. 1984	0.160	Sample et al. 1998a	0.069	Sample et al. 1998b	0.112	Sample et al. 1998b
Beryllium	0.010	Baes et al. 1984	1.182	Sample et al. 1998a	--	see text	--	see text
Cadmium	3.250	Bechtel Jacobs 1998a	40.69	Sample et al. 1998a	0.462	Sample et al. 1998b	7.017	Sample et al. 1998b
Chromium	0.084	Bechtel Jacobs 1998a	3.162	Sample et al. 1998a	0.349	Sample et al. 1998b	0.333	Sample et al. 1998b
Cobalt	0.020	Baes et al. 1984	0.291	Sample et al. 1998a	0.025	Sample et al. 1998b	0.100	Sample et al. 1998b
Copper	0.625	Bechtel Jacobs 1998a	1.531	Sample et al. 1998a	0.554	Sample et al. 1998b	1.117	Sample et al. 1998b
Iron	0.004	Baes et al. 1984	0.078	Sample et al. 1998a	0.015	Sample et al. 1998b	0.017	Sample et al. 1998b
Lead	0.468	Bechtel Jacobs 1998a	1.522	Sample et al. 1998a	0.286	Sample et al. 1998b	0.339	Sample et al. 1998b
Manganese	0.250	Baes et al. 1984	0.124	Sample et al. 1998a	0.037	Sample et al. 1998b	0.059	Sample et al. 1998b
Mercury	5.000	Bechtel Jacobs 1998a	20.63	Sample et al. 1998a	0.130	Sample et al. 1998b	0.192	Sample et al. 1998b
Nickel	1.411	Bechtel Jacobs 1998a	4.730	Sample et al. 1998a	0.589	Sample et al. 1998b	0.578	Sample et al. 1998b
Selenium	3.012	Bechtel Jacobs 1998a	1.340	Sample et al. 1998a	1.263	Sample et al. 1998b	1.187	Sample et al. 1998b
Silver	0.037	Bechtel Jacobs 1998a	15.34	Sample et al. 1998a	0.810	Sample et al. 1998b	0.501	Sample et al. 1998b
Thallium	0.004	Baes et al. 1984	1.000	--	0.123	Sample et al. 1998b	0.123	Sample et al. 1998b
Vanadium	0.006	Baes et al. 1984	0.088	Sample et al. 1998a	0.013	Sample et al. 1998b	0.018	Sample et al. 1998b
Zinc	1.820	Bechtel Jacobs 1998a	12.89	Sample et al. 1998a	2.782	Sample et al. 1998b	2.901	Sample et al. 1998b
Pesticides/PCBs								
4,4'-DDD	0.0151	Travis and Arms 1988	2.00	Menzie et al. 1992	--	see text	--	see text
Aroclor-1260	0.0045	Travis and Arms 1988	15.9	Sample et al. 1998a	--	see text	--	see text
Semivolatile Organics								
4-Nitrophenol	3.0889	Travis and Arms 1988	1.00	--	--	see text	--	see text
Acenaphthene	0.2564	Travis and Arms 1988	0.30	Beyer and Stafford 1993	--	see text	--	see text
Acenaphthylene	0.1653	Travis and Arms 1988	0.22	Beyer and Stafford 1993	--	see text	--	see text
Anthracene	0.1051	Travis and Arms 1988	0.32	Beyer and Stafford 1993	--	see text	--	see text
Benzo(a)anthracene	0.0222	Travis and Arms 1988	0.27	Beyer and Stafford 1993	--	see text	--	see text
Benzo(a)pyrene	0.0135	Travis and Arms 1988	0.34	Beyer and Stafford 1993	--	see text	--	see text
Benzo(b)fluoranthene	0.0174	Travis and Arms 1988	0.21	Beyer and Stafford 1993	--	see text	--	see text
Benzo(g,h,i)perylene	0.0061	Travis and Arms 1988	0.15	Beyer and Stafford 1993	--	see text	--	see text
Benzo(k)fluoranthene	0.0112	Travis and Arms 1988	0.21	Beyer and Stafford 1993	--	see text	--	see text
bis(2-Ethylhexyl)phthalate	0.0029	Travis and Arms 1988	1.00	--	--	see text	--	see text
Chrysene	0.0289	Travis and Arms 1988	0.44	Beyer and Stafford 1993	--	see text	--	see text
Dibenz(a,h)anthracene	0.0068	Travis and Arms 1988	0.49	Beyer and Stafford 1993	--	see text	--	see text
Di-n-butylphthalate	0.1124	Travis and Arms 1988	1.00	--	--	see text	--	see text
Di-n-octylphthalate	0.0009	Travis and Arms 1988	1.00	--	--	see text	--	see text
Fluoranthene	0.0617	Travis and Arms 1988	0.37	Beyer and Stafford 1993	--	see text	--	see text
Fluorene	0.1790	Travis and Arms 1988	0.20	Beyer and Stafford 1993	--	see text	--	see text
Indeno(1,2,3-cd)pyrene	0.0061	Travis and Arms 1988	0.41	Beyer and Stafford 1993	--	see text	--	see text
Naphthalene	0.5261	Travis and Arms 1988	0.21	Beyer and Stafford 1993	--	see text	--	see text
Phenanthrene	0.1154	Travis and Arms 1988	0.28	Beyer and Stafford 1993	--	see text	--	see text
Pyrene	0.0687	Travis and Arms 1988	0.39	Beyer and Stafford 1993	--	see text	--	see text
Volatile Organics								
1,2,4-Trichlorobenzene	0.2186	Travis and Arms 1988	0.56	Beyer 1996	--	see text	--	see text
Toluene	2.0447	Travis and Arms 1988	1.00	--	--	see text	--	see text

TABLE 4-3
 Exposure Parameters for Upper Trophic Level Ecological Receptors
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Receptor	Body Weight (kg)		Water Ingestion Rate (L/day)		Food Ingestion Rate (kg/day - dry)	
	Value	Reference	Value	Reference	Value	Reference
Birds						
American robin	0.064	USEPA 1993	0.0129	allometric equation	0.0057	Levey and Karasov 1989
Great blue heron	2.10	Butler 1992	0.1090	allometric equation	0.4389	allometric equation
Mallard	0.612	Bellrose 1980	0.0850	allometric equation	0.0830	allometric equation
Mammals						
Deer mouse	0.012	Silva and Downing 1995	0.0040	USEPA 1993a	0.0007	USEPA 1993a
Mink	0.726	Silva and Downing 1995	0.0286	USEPA 1993a	0.0317	USEPA 1993a
Raccoon	4.23	Silva and Downing 1995	0.6092	allometric equation	0.1245	Conover 1989
Least shrew	0.013	USEPA 1993	0.0048	USEPA 1993a	0.0019	USEPA 1993a

Receptor	Dietary Composition (percent)						Soil/ Sediment Ingestion (percent)	
	Terr. Plants	Soil Invert.	Small Mammals	Fish/ Frogs	Aquatic Plants	Benthic Invert.	Value	Reference
Birds								
American robin	12	78	0	0	0	0	10	Beyer et al. 1994
Great blue heron	0	0	0	90	0	5	5	Based on average minimum for birds in Beyer et al. 1994
Mallard	0	0	0	0	40.7	56.0	3.3	Beyer et al. 1994
Mammals								
Deer mouse	84.0	14.0	0	0	0	0	2.0	Based on similar species in Beyer et al, 1994
Mink	17	0	2.5	63.0	0.0	11.0	2.8	Based on value reported in Beyer et al. 1994 for raccoon
Raccoon	34.4	27.2	3.6	2.7	0.0	22.7	9.42	Beyer et al. 1994
Least shrew	0	82.3	0	0	0	0	2.0	Based on similar species in Beyer et al. 1994

TABLE 4-4
 Ingestion Screening Values for Mammals
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Test Organism	Body Weight (kg)	Duration	Exposure Route	Effect/Endpoint	LOAEL (mg/kg/d)	NOAEL (mg/kg/d)	Reference
Inorganics								
Aluminum	mouse	0.03	390 days	oral in water	reproduction	193	19.3	ATSDR 1990a
	dog	10	6 months	oral	reproduction	600	60	ATSDR 1990a
Antimony	mouse	0.03	lifetime	oral in water	lifespan/longevity	1.25	0.125	Sample et al. 1996
Arsenic	mouse	0.03	3 generations	oral in water	reproduction	1.26	0.126	Sample et al. 1996
Barium	rat	0.435	16 months	oral in water	growth/hypertension	19.8	5.1	Sample et al. 1996
Beryllium	rat	0.35	lifetime	oral in water	longevity/weight loss	6.6	0.66	Sample et al. 1996
Cadmium	rat	0.303	6 weeks	oral (gavage)	reproduction	10	1	Sample et al. 1996
	dog	10	3 months	oral	reproduction	7.5	0.75	ATSDR 1993a
Chromium	rat	0.35	3 months	oral in water	mortality	131.4	13.14	Sample et al. 1996
Cobalt	rat	0.35	69 days	oral in diet	reproduction	50	5	ATSDR 1992a
Copper	mink	1	357 days	oral in diet	reproduction	15.14	11.7	Sample et al. 1996
Iron	rabbit	3.8	?	oral in diet	tolerance level	500	50	NAS 1980
Lead	rat	0.35	3 generations	oral in diet	reproduction	80	8	Sample et al. 1996
Manganese	rat	0.35	224 days	oral in diet	reproduction	284	88	Sample et al. 1996
	rat	0.35	3 generations	oral in diet	reproduction	0.16	0.032	Sample et al. 1996
Mercury	mink	1	93 days	oral in diet	mortality/weight loss	0.25	0.15	Sample et al. 1996
	rat	0.35	3 generations	oral in diet	reproduction	80	40	Sample et al. 1996
Selenium	rat	0.35	1 year	oral in water	reproduction	0.33	0.2	Sample et al. 1996
Silver	rat	0.35	2 weeks	oral in water	mortality	181	18.1	ATSDR 1990b
Thallium	rat	0.35	60 days	oral in water	reproduction	0.74	0.074	Sample et al. 1996
Vanadium	rat	0.26	60 days +	oral intubation	reproduction	2.1	0.21	Sample et al. 1996
Zinc	rat	0.35	GD 1-16	oral in diet	reproduction	320	160	Sample et al. 1996
	mink	1	25 weeks	oral	reproduction	208	20.8	ATSDR 1992b

TABLE 4-4
 Ingestion Screening Values for Mammals
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Test Organism	Body Weight (kg)	Duration	Exposure Route	Effect/Endpoint	LOAEL (mg/kg/d)	NOAEL (mg/kg/d)	Reference
PCB/Pesticides								
4,4'-DDD	rat	0.35	2 years	oral in diet	reproduction	4	0.8	Sample et al. 1996
	dog	10	2 generations	oral	reproduction	5	1	ATSDR 1994
4,4'-DDE	rat	0.35	2 years	oral in diet	reproduction	4	0.8	Sample et al. 1996
	dog	10	2 generations	oral	reproduction	5	1	ATSDR 1994
4,4'-DDT	rat	0.35	2 years	oral in diet	reproduction	4	0.8	Sample et al. 1996
	dog	10	2 generations	oral	reproduction	5	1	ATSDR 1994
Aldrin	rat	0.35	3 generations	oral in diet	reproduction	1	0.2	Sample et al. 1996
alpha-BHC	rat	0.35	4 generations	oral in diet	reproduction	3.2	1.6	Sample et al. 1996
alpha-Chlordane	mouse	0.03	6 generations	oral in diet	reproduction	9.16	4.58	Sample et al. 1996
Aroclor-1016	mink	1	18 months	oral in diet	reproduction	3.43	1.37	Sample et al. 1996
Aroclor-1221	mink	1	7 months	oral in diet	reproduction	0.69	0.069	Sample et al. 1996
Aroclor-1232	mink	1	7 months	oral in diet	reproduction	0.69	0.069	Sample et al. 1996
Aroclor-1242	mink	1	7 months	oral in diet	reproduction	0.69	0.069	Sample et al. 1996
Aroclor-1248	mouse	0.03	5 weeks	oral in diet	immunological	13	1.3	ATSDR 1995a
Aroclor-1254	oldfield mouse	0.014	12 months	oral in diet	reproduction	0.68	0.068	Sample et al. 1996
	mink	1	4.5 months	oral in diet	reproduction	0.69	0.14	Sample et al. 1996
Aroclor-1260	oldfield mouse	0.014	12 months	oral in diet	reproduction	0.68	0.068	Sample et al. 1996
	mink	1	4.5 months	oral in diet	reproduction	0.69	0.14	Sample et al. 1996
beta-BHC	rat	0.35	13 weeks	oral in diet	growth/systemic	20	4	Sample et al. 1996
delta-BHC	rat	0.35	4 generations	oral in diet	reproduction	3.2	1.6	Sample et al. 1996
Dieldrin	rat	0.35	3 generations	oral in diet	reproduction	0.2	0.02	Sample et al. 1996
Endosulfan I	rat	0.35	30 days	oral (intubation)	reproduction	15	1.5	Sample et al. 1996
Endosulfan II	rat	0.35	30 days	oral (intubation)	reproduction	15	1.5	Sample et al. 1996
Endosulfan Sulfate	rat	0.35	30 days	oral (intubation)	reproduction	15	1.5	Sample et al. 1996
Endrin	mouse	0.03	120 days	oral in diet	reproduction	0.92	0.092	Sample et al. 1996
Endrin Aldehyde	mouse	0.03	120 days	oral in diet	reproduction	0.92	0.092	Sample et al. 1996
Endrin Ketone	mouse	0.03	120 days	oral in diet	reproduction	0.92	0.092	Sample et al. 1996
Gamma-BHC (Lindane)	rat	0.35	3 generations	oral in diet	reproduction	80	8	Sample et al. 1996
Gamma-Chlordane	mouse	0.03	6 generations	oral in diet	reproduction	9.16	4.58	Sample et al. 1996
Heptachlor	mink	1	181 days	oral in diet	reproduction	1	0.1	Sample et al. 1996
Heptachlor Epoxide	mink	1	181 days	oral in diet	reproduction	1	0.1	Sample et al. 1996
Methoxychlor	rat	0.35	11 months	oral in diet	reproduction	8	4	Sample et al. 1996
Toxaphene	rat	0.35	3 generations	oral in diet	reproduction	80	8	Sample et al. 1996

TABLE 4-4
Ingestion Screening Values for Mammals
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Test Organism	Body Weight (kg)	Duration	Exposure Route	Effect/Endpoint	LOAEL (mg/kg/d)	NOAEL (mg/kg/d)	Reference
Semivolatile Organics								
1,2,4-Trichlorobenzene	rat	0.35	3 generations	oral in water	reproduction	106	53	Coulston and Kolbye 1994
1,2-Dichlorobenzene	rat	0.35	chronic	oral (gavage)	liver/kidney	857	85.7	Coulston and Kolbye 1994
1,3-Dichlorobenzene	rat	0.35	chronic	oral (gavage)	liver/kidney	857	85.7	Coulston and Kolbye 1994
1,4-Dichlorobenzene	rat	0.35	GD 6-15	oral (gavage)	reproduction	500	250	Coulston and Kolbye 1994
2,4,5-Trichlorophenol	rat	0.35	98 days	oral in diet	hepatic/renal	800	80	McCollister et al. 1961
2,4,6-Trichlorophenol	rat	0.35	98 days	oral in diet	hepatic/renal	800	80	McCollister et al. 1961
2,4-Dichlorophenol	rat	0.35	103 weeks	oral in diet	reproduction	4400	440	NTP 1989
2-Chloronaphthalene	--	--	--	--	--	NA	NA	--
2-Methylnaphthalene	mouse	0.03	81 weeks	oral in diet	systemic	1437	143.7	ATSDR 1995b
3,3'-Dichlorobenzidine	--	--	--	--	--	NA	NA	--
4-Bromophenyl-Phenylether	--	--	--	--	--	NA	NA	--
4-Chloro-3-Methylphenol	--	--	--	--	--	NA	NA	--
4-Chlorophenyl-Phenylether	--	--	--	--	--	NA	NA	--
Acenaphthene	mouse	0.03	13 weeks	oral (gavage)	reproduction	3500	350	ATSDR 1995b
Acenaphthylene	mouse	0.03	13 weeks	oral (gavage)	reproduction	3500	350	ATSDR 1995b
Anthracene	mouse	0.03	13 weeks	oral (gavage)	reproduction	10000	1000	ATSDR 1995b
Benzo(a)anthracene	mouse	0.03	GD 7-16	oral (intubation)	reproduction	10	1	Sample et al. 1996
Benzo(a)pyrene	mouse	0.03	GD 7-16	oral (intubation)	reproduction	10	1	Sample et al. 1996
Benzo(b)fluoranthene	mouse	0.03	GD 7-16	oral (intubation)	reproduction	10	1	Sample et al. 1996
Benzo(g,h,i)perylene	mouse	0.03	19 to 29 days	oral in diet	reproduction	1330	133	ATSDR 1995b
Benzo(k)fluoranthene	mouse	0.03	GD 7-16	oral (intubation)	reproduction	10	1	Sample et al. 1996
Bis-(2-Ethylhexyl)phthalate	mouse	0.03	105 days	oral in diet	reproduction	183.3	18.3	Sample et al. 1996
Butylbenzylphthalate	rat	0.35	2 years	oral in diet	hepatic	2400	240	NTP 1997
Carbazole	mouse	0.03	19 to 29 days	oral in diet	reproduction	1330	133	ATSDR 1995b
Chrysene	mouse	0.03	GD 7-16	oral (intubation)	reproduction	10	1	Sample et al. 1996
Dibenz(a,h)anthracene	mouse	0.03	GD 7-16	oral (intubation)	reproduction	10	1	Sample et al. 1996
Dibenzofuran	mouse	0.03	19 to 29 days	oral in diet	reproduction	1330	133	ATSDR 1995b
Diethylphthalate	mouse	0.03	105 days	oral in diet	reproduction	45830	4583	Sample et al. 1996
Di-n-butylphthalate	mouse	0.03	105 days	oral in diet	reproduction	1833	550	Sample et al. 1996
Di-n-octylphthalate	mouse	0.03	105 days	oral in diet	reproduction	550	55	Sample et al. 1996
Fluoranthene	mouse	0.03	13 weeks	oral (gavage)	hepatic	1250	125	ATSDR 1995b
Fluorene	mouse	0.03	13 weeks	oral (gavage)	hematological	1250	125	ATSDR 1995b
Hexachloro-1,3-butadiene	rat	0.35	90 days +	oral	reproduction	20	2	IPCS 1994
Hexachlorobenzene	rat	0.35	2 years	oral	reproduction	16	1.6	ATSDR 1989
Hexachlorocyclopentadiene	rat	0.35	GD 6-15	oral	reproduction	30	10	USEPA 1984

TABLE 4-4
 Ingestion Screening Values for Mammals
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Test Organism	Body Weight (kg)	Duration	Exposure Route	Effect/Endpoint	LOAEL (mg/kg/d)	NOAEL (mg/kg/d)	Reference
Semivolatile Organics								
Hexachloroethane	--	--	--	--	--	NA	NA	--
Indeno(1,2,3-cd)pyrene	mouse	0.03	GD 7-16	oral (intubation)	reproduction	10	1	Sample et al. 1996
Naphthalene	mouse	0.03	13 weeks	oral (gavage)	reproduction	1400	140	ATSDR 1995c
N-Nitrosodiphenylamine	rat	0.35	8 to 11 weeks	oral in diet	systemic	1500	150	ATSDR 1993b
Pentachlorophenol	rat	0.35	up to 24 months	oral in diet	reproduction	30	3	Coulston and Kolbye 1994
Phenanthrene	mouse	0.03	19 to 29 days	oral in diet	reproduction	1330	133	ATSDR 1995b
Pyrene	mouse	0.03	19 to 29 days	oral in diet	reproduction	1330	133	ATSDR 1995b
Volatile Organics								
Carbon Tetrachloride	rat	0.35	2 years	oral in diet	reproduction	160	16	Sample et al. 1996
Chlorobenzene	dog	12.7	chronic	?	liver	273	27.3	IRIS 1998
Chloroform	rat	0.35	13 weeks	oral (intubation)	systemic	410	150	Sample et al. 1996
Ethylbenzene	rat	0.35	chronic	?	liver/kidney	971	97.1	Wolf et al. 1956
Styrene	rat	0.35	?	?	?	350	35	Beliles et al. 1985
	dog	12.7	chronic	?	blood/liver	400	200	IRIS 1998
Tetrachloroethene	mouse	0.03	6 weeks	oral (gavage)	hepatotoxicity	70	14	Sample et al. 1996
Toluene	mouse	0.03	GD 6-12	oral (gavage)	reproduction	260	26	Sample et al. 1996
Trichloroethene	rat	0.35	?	oral	reproduction	10000	1000	Coulston and Kolbye 1994
Xylenes (total)	mouse	0.03	GD 6-15	oral (gavage)	reproduction	2.6	2.1	Sample et al. 1996

TABLE 4-5
Ingestion Screening Values for Birds
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Test Organism	Body Weight (kg)	Duration	Exposure Route	Effect/Endpoint	LOAEL (mg/kg/d)	NOAEL (mg/kg/d)	Reference
Inorganics								
Aluminum	ringed dove	0.155	4 months	oral in diet	reproduction	1097	109.7	Sample et al. 1996
Antimony	northern bobwhite	0.19	6 weeks	oral	?	47400	4740	Opresko et al. 1993
Arsenic	brown-headed cowbird	0.049	7 months	oral in diet	mortality	7.38	2.46	Sample et al. 1996
	mallard	1	128 days	oral in diet	mortality	12.84	5.14	Sample et al. 1996
Barium	chicks	0.121	4 weeks	oral in diet	mortality	417	208	Sample et al. 1996
Beryllium	--	--	--	--	--	NA	NA	--
Cadmium	mallard	1.153	90 days	oral in diet	reproduction	20	1.45	Sample et al. 1996
Chromium	American black duck	1.25	10 months	oral in diet	reproduction	5	1	Sample et al. 1996
Cobalt	chicken	1.8	14 days	oral in diet	growth	14.7	1.47	Diaz et al. 1994
Copper	chicks	0.534	10 weeks	oral in diet	growth/mortality	61.7	47	Sample et al. 1996
Iron	chicken	1.6	?	oral	maximum tolerance level	1000	100	NAS 1980
Lead	Japanese quail	0.15	12 weeks	oral in diet	reproduction	11.3	1.13	Sample et al. 1996
	American kestrel	0.13	7 months	oral in diet	reproduction	38.5	3.85	Sample et al. 1996
Manganese	Japanese quail	0.072	75 days	oral in diet	growth/behavior	9770	977	Sample et al. 1996
Mercury	Japanese quail	0.15	1 year	oral in diet	reproduction	0.9	0.45	Sample et al. 1996
	mallard	1	3 generations	oral in diet	reproduction	0.064	0.0064	Sample et al. 1996
Nickel	mallard	0.782	90 days	oral in diet	growth/mortality	107	77.4	Sample et al. 1996
Selenium	mallard	1	100 days	oral in diet	reproduction	0.8	0.4	Sample et al. 1996
	screech owl	0.2	13.7 weeks	oral in diet	reproduction	1.5	0.44	Sample et al. 1996
Silver	--	--	--	--	--	NA	NA	--
Thallium	--	--	--	--	--	NA	NA	--
Vanadium	mallard	1.17	12 weeks	oral in diet	growth/mortality	114	11.4	Sample et al. 1996
Zinc	chicken	1.935	44 weeks	oral in diet	reproduction	131	14.5	Sample et al. 1996

TABLE 4-5
 Ingestion Screening Values for Birds
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Test Organism	Body Weight (kg)	Duration	Exposure Route	Effect/Endpoint	LOAEL (mg/kg/d)	NOAEL (mg/kg/d)	Reference
PCB/Pesticides								
4,4'-DDD	mallard	1.134	chronic	oral	reproduction	5.2	0.52	Stickel 1973
	American kestrel	0.115	2 years	oral	reproduction	0.5	0.05	McLane and Hall 1972
4,4'-DDE	brown pelican	3.5	chronic	oral	reproduction	1.31	0.131	Beyer et al. 1996
	American kestrel	0.115	2 years	oral	reproduction	0.5	0.05	McLane and Hall 1972
4,4'-DDT	mallard	1.134	chronic	oral	reproduction	1.04	0.104	Davison and Sell 1974
	American kestrel	0.115	2 years	oral	reproduction	0.5	0.05	McLane and Hall 1972
Aldrin	mallard	1.134	chronic	oral	mortality	5	0.5	Tucker and Crabtree 1970
alpha-BHC	Japanese quail	0.15	90 days	oral in diet	reproduction	2.25	0.56	Sample et al. 1996
alpha-Chlordane	red-winged blackbird	0.064	84 days	oral in diet	mortality	10.7	2.14	Sample et al. 1996
Aroclor-1016	screech owl	0.181	2 generations	oral in diet	reproduction	4.1	0.41	Sample et al. 1996
Aroclor-1221	screech owl	0.181	2 generations	oral in diet	reproduction	4.1	0.41	Sample et al. 1996
Aroclor-1232	screech owl	0.181	2 generations	oral in diet	reproduction	4.1	0.41	Sample et al. 1996
Aroclor-1242	screech owl	0.181	2 generations	oral in diet	reproduction	4.1	0.41	Sample et al. 1996
Aroclor-1248	ring-necked pheasant	1	17 weeks	oral	reproduction	1.8	0.18	Sample et al. 1996
Aroclor-1254	ring-necked pheasant	1	17 weeks	oral	reproduction	1.8	0.18	Sample et al. 1996
Aroclor-1260	ring-necked pheasant	1	17 weeks	oral	reproduction	1.8	0.18	Sample et al. 1996
beta-BHC	Japanese quail	0.15	90 days	oral in diet	reproduction	2.25	0.56	Sample et al. 1996
delta-BHC	Japanese quail	0.15	90 days	oral in diet	reproduction	2.25	0.56	Sample et al. 1996
Dieldrin	barn owl	0.466	2 years	oral in diet	reproduction	0.77	0.077	Sample et al. 1996
Endosulfan I	gray partridge	0.4	4 weeks	oral in diet	reproduction	100	10	Sample et al. 1996
Endosulfan II	gray partridge	0.4	4 weeks	oral in diet	reproduction	100	10	Sample et al. 1996
Endosulfan Sulfate	gray partridge	0.4	4 weeks	oral in diet	reproduction	100	10	Sample et al. 1996
Endrin	mallard	1.15	>200 days	oral in diet	reproduction	3	0.3	Sample et al. 1996
	screech owl	0.181	>83 days	oral in diet	reproduction	0.1	0.01	Sample et al. 1996
Endrin Aldehyde	mallard	1.15	>200 days	oral in diet	reproduction	3	0.3	Sample et al. 1996
	screech owl	0.181	>83 days	oral in diet	reproduction	0.1	0.01	Sample et al. 1996
Endrin Ketone	mallard	1.15	>200 days	oral in diet	reproduction	3	0.3	Sample et al. 1996
	screech owl	0.181	>83 days	oral in diet	reproduction	0.1	0.01	Sample et al. 1996
Gamma-BHC (Lindane)	mallard	1	8 weeks	oral (intubation)	reproduction	20	2	Sample et al. 1996
Gamma-Chlordane	red-winged blackbird	0.064	84 days	oral in diet	mortality	10.7	2.14	Sample et al. 1996
Heptachlor	quail	0.191	5 days	oral in diet	mortality	4.05	0.405	Hill et al. 1975
Heptachlor Epoxide	quail	0.191	5 days	oral in diet	mortality	4.05	0.405	Hill et al. 1975
Methoxychlor	quail	0.191	5 days	oral in diet	mortality	4050	405	Hill and Camardese 1986
Toxaphene	mallard	1.043	5 days	oral in diet	mortality	3.07	0.307	Hill and Camardese 1986

TABLE 4-5
Ingestion Screening Values for Birds
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Test Organism	Body Weight (kg)	Duration	Exposure Route	Effect/Endpoint	LOAEL (mg/kg/d)	NOAEL (mg/kg/d)	Reference
Semivolatile Organics								
1,2,4-Trichlorobenzene	--	--	--	--	--	NA	NA	--
1,2-Dichlorobenzene	northern bobwhite	0.157	14 days	oral (gavage)	growth/mortality	2500	250	Grimes and Jaber 1989
1,3-Dichlorobenzene	northern bobwhite	0.157	14 days	oral (gavage)	growth/mortality	2500	250	Grimes and Jaber 1989
1,4-Dichlorobenzene	northern bobwhite	0.157	14 days	oral (gavage)	growth/mortality	2500	250	Grimes and Jaber 1989
2,4,5-Trichlorophenol	--	--	--	--	--	NA	NA	--
2,4,6-Trichlorophenol	--	--	--	--	--	NA	NA	--
2,4-Dichlorophenol	--	--	--	--	--	NA	NA	--
2-Chloronaphthalene	--	--	--	--	--	NA	NA	--
2-Methylnaphthalene	--	--	--	--	--	NA	NA	--
3,3'-Dichlorobenzidine	--	--	--	--	--	NA	NA	--
4-Bromophenyl-Phenylether	--	--	--	--	--	NA	NA	--
4-Chloro-3-Methylphenol	--	--	--	--	--	NA	NA	--
4-Chlorophenyl-Phenylether	--	--	--	--	--	NA	NA	--
Acenaphthene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Acenaphthylene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Anthracene	mallard	1.043	7 months	oral in diet	hepatic	228	22.8	Patton and Dieter 1980
Benzo(a)anthracene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Benzo(a)pyrene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Benzo(b)fluoranthene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Benzo(g,h,i)perylene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Benzo(k)fluoranthene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Bis-(2-Ethylhexyl)phthalate	ringed dove	0.155	4 weeks	oral in diet	reproduction	11	1.1	Sample et al. 1996
Butylbenzylphthalate	--	--	--	--	--	NA	NA	--
Carbazole	--	--	--	--	--	NA	NA	--
Chrysene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Dibenz(a,h)anthracene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Dibenzofuran	--	--	--	--	--	NA	NA	--
Diethylphthalate	--	--	--	--	--	NA	NA	--
Di-n-butylphthalate	ringed dove	0.155	4 weeks	oral in diet	reproduction	1.1	0.11	Sample et al. 1996
Di-n-octylphthalate	ring-necked pheasant	1	?	?	mortality	500	50	TERRETOX 1998

TABLE 4-5
 Ingestion Screening Values for Birds
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Test Organism	Body Weight (kg)	Duration	Exposure Route	Effect/Endpoint	LOAEL (mg/kg/d)	NOAEL (mg/kg/d)	Reference
Semivolatile Organics								
Fluoranthene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Fluorene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Hexachloro-1,3-butadiene	Japanese quail	0.19	90 days	oral	reproduction	8	2.5	Coulston and Kolbye 1994; IPCS 1994
Hexachlorobenzene	Japanese quail	0.19	?	oral	reproduction	0.8	0.08	Coulston and Kolbye 1994
Hexachlorocyclopentadiene	--	--	--	--	--	NA	NA	--
Hexachloroethane	--	--	--	--	--	NA	NA	--
Indeno(1,2,3-cd)pyrene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Naphthalene	mallard	1.04	7 months	oral in diet	hepatic	228	22.8	Patton and Dieter 1980
N-Nitrosodiphenylamine	--	--	--	--	--	NA	NA	--
Pentachlorophenol	chicken	1.5	8 weeks	oral	growth	200	100	Eisler 1989
Phenanthrene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Pyrene	chicken	1.5	34 days	oral in diet	reproduction	395	39.5	Rigdon and Neal 1963
Volatile Organics								
Carbon Tetrachloride	--	--	--	--	--	NA	NA	--
Chlorobenzene	--	--	--	--	--	NA	NA	--
Chloroform	--	--	--	--	--	NA	NA	--
Ethylbenzene	--	--	--	--	--	NA	NA	--
Styrene	--	--	--	--	--	NA	NA	--
Tetrachloroethene	--	--	--	--	--	NA	NA	--
Toluene	--	--	--	--	--	NA	NA	--
Trichloroethene	--	--	--	--	--	NA	NA	--
Xylenes (total)	quail	0.191	subacute	?	"toxicity"	405	40.5	Hill and Camardese 1986

TABLE 4-6
 Results of Rad-BCG Screening, KCK Sediment Maximum Concentrations
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Nuclide	Water (pCi/L)		Sediment (pCi/g)	
	Partial Fraction	Source of Calculation	Partial Fraction	Source of Calculation
Am-241				
Ce-144				
Cs-135				
Cs-137	3.4E-02	RA-Lumped, Default	2.3E-04	RA-Lumped, Default
Co-60				
Eu-154				
Eu-155				
H-3				
I-129				
I-131				
Pu-239				
Ra-226	1.9E+02	RA-Lumped, Default	5.3E-01	RA-Lumped, Default
Ra-228	2.8E+03	RA-Lumped, Default	7.5E+00	RA-Lumped, Default
Sb-125				
Sr-90				
Tc-99				
Th-232	3.6E-02	AA Default BiV	5.0E-01	RA-Lumped, Default
U-233				
U-234	4.7E-01	AA Default BiV	8.9E-03	RA-Lumped, Default
U-235	4.0E-01	AA Default BiV	1.2E-03	RA-Lumped, Default
U-238	3.8E+00	AA Default BiV	1.7E-02	RA-Lumped, Default
Zn-65				
Zr-95				
Partial fractions	2.9E+03		8.5E+00	
Total sum of fractions (water and sediment):				3.0E+03
Result:	You have failed the site screen			

RA: Riparian Animal
 AA: Aquatic Animal
 BiV: Bioaccumulation value

TABLE 4-7

Results of Rad-BCG Screening, KCK Sediment Mean Concentrations
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Nuclide	Water (pCi/L)		Sediment (pCi/g)	
	Partial Fraction	Source of Calculation	Partial Fraction	Source of Calculation
Am-241				
Ce-144				
Cs-135				
Cs-137	6.5E-03	RA-Lumped, Default	4.4E-05	RA-Lumped, Default
Co-60				
Eu-154				
Eu-155				
H-3				
I-129				
I-131				
Pu-239				
Ra-226	1.4E+01	RA-Lumped, Default	3.9E-02	RA-Lumped, Default
Ra-228	1.2E+02	RA-Lumped, Default	3.2E-01	RA-Lumped, Default
Sb-125				
Sr-90				
Tc-99				
Th-232	1.5E-03	AA Default BiV	2.1E-02	RA-Lumped, Default
U-233				
U-234	3.5E-01	AA Default BiV	6.6E-04	RA-Lumped, Default
U-235	3.2E-02	AA Default BiV	9.4E-05	RA-Lumped, Default
U-238	3.1E-01	AA Default BiV	1.4E-03	RA-Lumped, Default
Zn-65				
Zr-95				
Partial fractions	1.3E+02		3.9E-01	
Total sum of fractions (water and sediment):				1.3E+02
Result:	You have failed the site screen			

RA: Riparian Animal
 AA: Aquatic Animal
 BiV: Bioaccumulation value

TABLE 4-8

Comparison of Concentrations of Detected Analytes in KCK Sediment to Ecological Benchmark Values
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

ParamName	Units	Maximum Detection	Arithmetic Mean Concentration (1/2 ND)	Screening Criteria	Number of Analyses	Number of Detections	Number of Detections Exceeding Criteria	Detection Frequency (%)	Maximum Hazard Quotient	Mean Hazard Quotient	COPC?	Criteria Source
Metals												
Aluminum	mg/kg	17300	7406.32	58030	93	93	0	100	0.30	0.13	No	PEC ^a
Arsenic	mg/kg	177	13.16	6	93	86	47	92	29.50	2.19	Yes	LEL ^a
Cadmium	mg/kg	2.6	0.24	0.59	93	5	2	5	4.41	0.41	Yes	TEC ^a
Chromium, Total	mg/kg	43.4	14.07	26	93	83	11	89	1.67	0.54	Yes	LEL ^a
Copper	mg/kg	2870	84.85	16	93	66	35	71	179.38	5.30	Yes	LEL ^a
Iron	mg/kg	46700	16139.35	188400	93	93	0	100	0.25	0.09	No	Lowest ARCs TEL (NOAA SQRT)
Lead	mg/kg	3096	80.96	31	93	93	32	100	99.87	2.61	Yes	LEL ^a
Manganese	mg/kg	2690	640.43	460	93	93	59	100	5.85	1.39	Yes	LEL ^a
Mercury	mg/kg	2	0.09	0.15	93	31	10	33	13.33	0.57	Yes	ER-L ^b
Nickel	mg/kg	54.2	12.22	16	93	74	24	80	3.39	0.76	Yes	LEL ^a
Silver	mg/kg	3	0.67	1	82	10	6	12	3.00	0.67	Yes	ER-L ^b
Zinc	mg/kg	2480	137.66	120	93	89	24	96	20.67	1.15	Yes	LEL ^a
Semivolatiles												
2-Methylnaphthalene	mg/kg	0.097	0.19	0.07	6	1	1	17	1.39	2.73	Yes	ER-L ^a
Acenaphthene	mg/kg	0.051	0.18	0.016	6	1	1	17	3.19	11.47	Yes	ER-L ^b
Acenaphthylene	mg/kg	0.48	0.26	0.044	6	1	1	17	10.91	5.80	Yes	ER-L ^a
Anthracene	mg/kg	0.92	0.30	0.027	6	3	3	50	34.07	11.05	Yes	LCV ^a
Benzo(a)Anthracene	mg/kg	1.8	0.49	0.11	6	2	2	33	16.36	4.42	Yes	SCV ^a
Benzo(a)Pyrene	mg/kg	0.69	0.29	0.14	6	1	1	17	4.93	2.07	Yes	SCV ^a
Bis(2-Ethylhexyl) Phthalate	mg/kg	0.078	0.19	890	6	1	0	17	0.00	0.00	No	ORNL
Chrysene	mg/kg	1.6	0.43	0.34	6	3	1	50	4.71	1.25	Yes	LEL ^a
Dibenzofuran	mg/kg	0.29	0.22	2	6	1	0	17	0.15	0.11	No	SQB ^b
Di-n-Butyl Phthalate	mg/kg	0.16	0.21	11	6	1	0	17	0.02	0.02	No	SCV ^a
Fluoranthene	mg/kg	2.7	0.69	0.6	6	3	2	50	4.50	1.15	Yes	ER-L ^b
Fluorene	mg/kg	0.74	0.27	0.019	6	2	2	33	38.95	14.40	Yes	ER-L ^b
Indeno(1,2,3-c,d)Pyrene	mg/kg	0.77	0.30	0.078	6	1	1	17	9.87	3.89	Yes	TEC ^a
Naphthalene	mg/kg	0.11	0.19	0.033	6	1	1	17	3.33	5.86	Yes	TEC ^a
Phenanthrene	mg/kg	2.9	0.71	0.24	6	3	2	50	12.08	2.95	Yes	ER-L ^b
Pyrene	mg/kg	1.9	0.48	0.49	6	3	1	50	3.88	0.99	Yes	LEL ^a

TABLE 4-8
Comparison of Concentrations of Detected Analytes in KCK Sediment to Ecological Benchmark Values
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

ParamName	Units	Maximum Detection	Arithmetic Mean Concentration (1/2 ND)	Screening Criteria	Number of Analyses	Number of Detections	Number of Detections Exceeding Criteria	Detection Frequency (%)	Maximum Hazard Quotient	Mean Hazard Quotient	COPC?	Criteria Source
Volatiles												
1,2,4-Trichlorobenzene	mg/kg	0.058	0.18	9.6	6	1	0	17	0.01	0.02	No	SCV *
Pesticides/PCBs												
p,p'-DDD	mg/kg	0.28	0.05	0.002	6	2	2	33	140.00	25.26	Yes	ER-L *
Aroclor 1260	mg/kg	2	0.35	0.0023	6	1	1	17	869.57	152.54	Yes	ER-L* - Total PCB
Radionuclides												
Cs-137	pCi/g	0.73	0.14	NA	21	21	NA	100	NA	NA	NA	NA
Ra-226	pCi/g	53.6	3.97	NA	153	116	NA	76	NA	NA	NA	NA
Ra-228	pCi/g	653	28.30	NA	114	114	NA	100	NA	NA	NA	NA
Th-232	pCi/g	654	27.40	NA	125	125	NA	100	NA	NA	NA	NA
U-234	pCi/g	47.1	3.50	NA	90	73	NA	81	NA	NA	NA	NA
U-235	pCi/g	4.38	0.35	NA	115	115	NA	100	NA	NA	NA	NA
U-238	pCi/g	42.6	3.51	NA	125	125	NA	100	NA	NA	NA	NA

NA: Not applicable; radionuclides were screened using DOE's RAD-BCG model

* As reported in Jones et al. 1997; SCV - Secondary Chronic Value, LCV - Lowest Chronic Value (Suter and Tsao 1996), LEL - Lowest Effect Level (Persaud et al. 1993), ER-L - Effects Range - Low (Long et al. 1995), TEC - Threshold Effect Concentration, PEC - Possible Effects Concentration (USEPA 1996)

† As reported in U.S. EPA Ecotox, 1998; SQC - Sediment Quality Criteria, SQB - Sediment Quality Benchmark (U.S. EPA, 1995), ER-L (Long et al., 1995)

NOAA SQRT -- Screening Quick Reference Tables NOAA 1999.

TABLE 4-9
Comparison of Concentrations of Detected Analytes in KCK Surface Water to Ecological Benchmark Values
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

ParamName	Units	Maximum Detection	Arithmetic Mean Concentration (1/2 ND)	Screening Criteria	Number of Analyses	Number of Detections	Number of Detections Exceeding Criteria	Detection Frequency	Maximum Hazard Quotient	Mean Hazard Quotient	COPC?	Criteria Source
Metals												
Barium	ug/L	107	80.57	3.9	12	12	12	100	27.44	20.66	Yes	Tier II ^a
Cobalt	ug/L	4.3	1.50	3	12	1	1	8	1.43	0.50	Yes	Tier II ^a
Copper	ug/L	14.5	7.85	0.23	12	10	10	83	63.04	34.13	Yes	LCV ^b
Lead	ug/L	10.4	5.00	3.2	12	5	5	42	3.25	1.56	Yes	AWQC ^a
Nickel	ug/L	7	3.21	5	12	1	1	8	1.40	0.64	Yes	LCV ^b
Vanadium	ug/L	9	5.68	19	12	11	0	92	0.47	0.30	No	Tier II ^a

^a USEPA 1996; AWQC = National ambient water quality criteria; Tier II = Values calculated using Great Lakes Water Quality Initiative Tier II methodology.

^b ORNL = Oak Ridge National Laboratory (Suter and Tsao 1996); LCV - Lowest Chronic Value; Tier II = Values calculated using GLWQI Tier II methodology.

TABLE 4-10
 SERA Food Web Model Results for KCK
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Raccoon			Mink			Great Blue heron			Mallard		
	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC
Inorganics												
Aluminum	0.74	0.15	0.33	10.26	2.05	4.59	31.43	6.29	14.06	1.57	0.31	0.70
Antimony	1.01	0.20	0.45	4.06	0.81	1.82	NA	NA	NA	NA	NA	NA
Arsenic	0.18	0.04	0.08	0.72	0.14	0.32	1.19	0.48	0.75	2.37	0.95	1.50
Barium	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cadmium	0.01	<0.01	<0.01	0.03	<0.01	0.01	0.08	<0.01	0.02	0.39	0.03	0.10
Chromium	0.03	<0.01	0.01	0.04	<0.01	0.02	0.85	0.17	0.38	1.01	0.20	0.45
Cobalt	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Copper	0.29	0.22	0.25	1.02	0.79	0.90	1.81	1.38	1.58	2.56	1.95	2.24
Iron	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Lead	1.19	0.12	0.37	1.82	0.18	0.58	21.73	4.35	9.72	150.81	15.08	47.69
Manganese	0.25	0.08	0.14	1.03	0.32	0.57	0.58	0.12	0.26	0.26	0.05	0.12
Mercury	0.32	0.19	0.24	1.88	1.13	1.46	69.39	23.13	40.06	38.31	12.77	22.12
Nickel	<0.01	<0.01	<0.01	0.06	0.03	0.04	0.14	0.10	0.12	0.07	0.05	0.06
Selenium	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Silver	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.02	<0.01	<0.01	<0.01	<0.01	<0.01
Vanadium	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Zinc	0.23	0.05	0.10	0.71	0.14	0.32	6.78	0.75	2.25	19.82	2.19	6.60
Pesticides/PCBs												
4,4'-DDD	<0.01	<0.01	<0.01	0.02	<0.01	<0.01	0.42	0.04	0.13	0.26	0.05	0.12
Aroclor-1260	2.25	0.46	1.01	6.60	1.34	2.97	3.56	0.71	1.59	2.22	0.44	0.99

TABLE 4-10

SERA Food Web Model Results for KCK
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Raccoon			Mink			Great Blue heron			Mallard		
	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC
Semivolatile Organics												
2-Methylnaphthalene	NA	NA	NA	NA	NA	NA	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Acenaphthene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Acenaphthylene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.01	<0.01	<0.01	0.01	<0.01	<0.01
Anthracene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.03	<0.01	0.01	<0.01	<0.01	<0.01
Benzo(a)anthracene	0.05	0.01	0.02	0.06	0.01	0.03	0.07	0.01	0.03	0.17	0.03	0.08
Benzo(a)pyrene	<0.01	<0.01	<0.01	0.01	<0.01	<0.01	0.02	<0.01	<0.01	0.02	<0.01	<0.01
bis(2-Ethylhexyl)phthalate	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Chrysene	109.12	21.82	48.80	78.47	15.69	35.09	48.13	9.63	21.52	349.31	69.86	156.22
Di-n-butylphthalate	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.17	0.03	0.08	0.07	0.01	0.03
Fluoranthene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.17	0.03	0.07	0.66	0.13	0.30
Fluorene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.02	<0.01	<0.01	0.01	<0.01	<0.01
Indeno(1,2,3-cd)pyrene	<0.01	<0.01	<0.01	0.01	<0.01	<0.01	0.02	<0.01	<0.01	<0.01	<0.01	<0.01
Naphthalene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Phenanthrene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.08	0.02	0.04	0.02	<0.01	0.01
Pyrene	283.50	56.70	126.78	203.84	40.77	91.16	124.98	25.00	55.89	907.49	181.50	405.84
Volatile Organics												
1,2,4-Trichlorobenzene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01

Bolded HQs indicate exceedance of 1.0

TABLE 4-11

Results of Rad-BCG Screening, STP River Sediments and Surface Water Maximum Concentrations
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Nuclide	Water (pCi/L)		Sediment (pCi/g)	
	Partial Fraction	Source of Calculation	Partial Fraction	Source of Calculation
Ce-144				
Cs-135				
Cs-137				
Co-60				
Eu-154				
Eu-155				
H-3				
I-129				
I-131				
Pu-239				
Ra-226	2.0E-01	RA-Lumped, Default	4.7E-02	RA-Lumped, Default
Ra-228	3.7E-01	RA-Lumped, Default	1.2E+00	RA-Lumped, Default
Sb-125				
Sr-90				
Tc-99				
Th-232	5.7E-04	AA Default BiV	7.6E-02	RA-Lumped, Default
U-233				
U-234	2.8E-03	AA Default BiV	1.8E-03	RA-Lumped, Default
U-235	1.7E-04	AA Default BiV	7.6E-05	RA-Lumped, Default
U-238	2.6E-03	AA Default BiV	2.4E-03	RA-Lumped, Default
Zn-65				
Zr-95				
Partial fractions	5.8E-01		1.4E+00	
Total sum of fractions (water and sediment):				2.0E+00
Result:	You have failed the site screen			

RA: Riparian Animal

AA: Aquatic Animal

BiV: Bioaccumulation value

TABLE 4-12

Results of Rad-BCG Screening, STP River Sediments and Surface Water Mean Concentrations
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Nuclide	Water (pCi/L)		Sediment (pCi/g)	
	Partial Fraction	Source of Calculation	Partial Fraction	Source of Calculation
Ce-144				
Cs-135				
Cs-137				
Co-60				
Eu-154				
Eu-155				
H-3				
I-129				
I-131				
Pu-239				
Ra-226	1.1E-01	RA-Lumped, Default	1.3E-02	RA-Lumped, Default
Ra-228	1.8E-01	RA-Lumped, Default	7.3E-02	RA-Lumped, Default
Sb-125				
Sr-90				
Tc-99				
Th-232	3.3E-04	AA Default BiV	5.1E-03	RA-Lumped, Default
U-233				
U-234	1.9E-03	AA Default BiV	2.0E-04	RA-Lumped, Default
U-235	9.2E-05	AA Default BiV	1.3E-05	RA-Lumped, Default
U-238	1.5E-03	AA Default BiV	4.6E-04	RA-Lumped, Default
Zn-65				
Zr-95				
Partial fractions	2.9E-01		9.2E-02	
Total sum of fractions (water and sediment):				3.8E-01
Result:	You have passed the site screen			

RA: Riparian Animal
 AA: Aquatic Animal
 BiV: Bioaccumulation value

TABLE 4-13
 Comparison of Concentrations of Detected Analytes in STP River Sediment to Ecological Benchmark Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

ParamName	Units	Maximum Detection	Arithmetic Mean Concentration	Screening Criteria	Number of Analyses	Number of Detections	Number of Detections Exceeding Criteria	Detection Frequency (%)	Maximum Hazard Quotient	Mean Hazard Quotient	COPC?	Criteria Source
Metals												
Aluminum	mg/kg	13500	5040.51	58030	39	39	0	100	0.23	0.09	No	PEC ^a
Antimony	mg/kg	12.1	2.93	3000	31	4	0	13	0.00	0.00	No	UET (NOAA SQRT)
Arsenic	mg/kg	31.6	8.60	6	39	39	24	100	5.27	1.43	Yes	LEL ^a
Cadmium	mg/kg	0.75	0.20	0.59	39	6	2	15	1.27	0.33	Yes	TEC ^a
Chromium, Total	mg/kg	36.9	11.29	26	39	30	2	77	1.42	0.43	Yes	LEL ^a
Copper	mg/kg	79.3	21.17	16	39	32	15	82	4.96	1.32	Yes	LEL ^a
Iron	mg/kg	51700	17368.21	188400	39	39	0	100	0.27	0.09	No	Lowest ARCs TEL (NOAA SQRT)
Manganese	mg/kg	1630	580.26	460	39	39	22	100	3.54	1.26	Yes	LEL ^a
Mercury	mg/kg	3.7	0.15	0.15	39	8	4	21	24.67	1.00	Yes	ER-L ^b
Nickel	mg/kg	28.1	12.15	16	39	39	14	100	1.76	0.76	Yes	LEL ^a
Silver	mg/kg	2.4	0.74	1	39	7	7	18	2.40	0.74	Yes	ER-L ^b
Zinc	mg/kg	207	70.11	120	39	36	11	92	1.73	0.58	Yes	LEL ^a
Semivolatile												
Benzo(a)Anthracene	mg/kg	0.17	0.17	0.11	3	2	2	67	1.55	1.56	Yes	SCV ^a
Benzo(a)Pyrene	mg/kg	0.099	0.12	0.14	3	2	0	67	0.71	0.86	No	SCV ^a
Chrysene	mg/kg	0.18	0.18	0.34	3	2	0	67	0.53	0.53	No	LEL ^a
Fluoranthene	mg/kg	0.56	0.41	0.6	3	2	0	67	0.93	0.68	No	ER-L ^b
Fluorene	mg/kg	0.05	0.12	0.019	2	1	1	50	2.63	6.45	Yes	ER-L ^b
Indeno(1,2,3-c,d)Pyrene	mg/kg	0.079	0.12	0.078	3	2	1	67	1.01	1.50	Yes	TEC ^a
Phenanthrene	mg/kg	0.41	0.33	0.24	3	2	2	67	1.71	1.35	Yes	ER-L ^b
Pyrene	mg/kg	0.46	0.36	0.49	3	2	0	67	0.94	0.72	No	LEL ^a
Volatile												
Toluene	mg/kg	0.017	0.01	0.67	3	3	0	100	0.03	0.01	No	EPA SQB
Radionuclides												
Ra-226	pCi/g	4.8	1.29	NA	48	34	NA	71	NA	NA	NA	
Ra-228	pCi/g	108.5	6.12	NA	44	44	NA	100	NA	NA	NA	
Th-232	pCi/g	99.2	6.41	NA	48	48	NA	100	NA	NA	NA	
U-234	pCi/g	9.6	1.04	NA	39	38	NA	97	NA	NA	NA	
U-235	pCi/g	0.28	0.05	NA	38	38	NA	100	NA	NA	NA	
U-238	pCi/g	6	1.14	NA	48	48	NA	100	NA	NA	NA	

NA: Not applicable; radionuclides were screened using DOE's RAD-BCG model

^a As reported in Jones et al. 1997; SCV - Secondary Chronic Value, LCV - Lowest Chronic Value (Suter and Tsao 1996), LEL - Lowest Effect Level (Persaud et al. 1993), ER-L - Effects Range - Low (Long et al. 1995), TEC - Threshold Effect Concentration, PEC - Possible Effects Concentration (USEPA 1996)

^b As reported in USEPA Ecotox 1996; SQC - Sediment Quality Criteria, SQB - Sediment Quality Benchmark (USEPA 1995), ER-L (Long et al. 1995)

NOAA SQRT: Screening Quick Reference Tables, NOAA 1999.

EPA SQB: Sediment quality benchmarks by equilibrium partitioning. Assumes 1 percent organic carbon (USEPA 1995)

TABLE 4-14

Comparison of Concentrations of Detected Analytes in STP River Surface Water to Ecological Benchmark Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

ParamName	Units	Maximum Detection	Arithmetic Mean Concentration	Screening Criteria	Number of Analyses	Number of Detections	Number of Detections Exceeding Criteria	Detection Frequency (%)	Maximum Hazard Quotient	Mean Hazard Quotient	COPC?	Criteria Source
Metals												
Barium	ug/L	93	78.30	3.9	6	6	6	100	23.85	20.08	Yes	Tier II ^a
Iron	ug/L	6720	3968.33	1000	6	6	6	100	6.72	3.97	Yes	AWQC ^a
Lead	ug/L	13.1	7.22	3.2	6	6	6	100	4.09	2.26	Yes	AWQC ^a
Manganese	ug/L	152	95.20	80	6	6	3	100	1.90	1.19	Yes	Tier II ^a
Nickel	ug/L	11.8	7.23	5	6	3	3	50	2.36	1.45	Yes	LCV ^b
Selenium	ug/L	2.6	1.48	5	6	1	0	17	0.52	0.30	No	AWQC ^a
Thallium	ug/L	2.2	1.10	40	6	1	0	17	0.06	0.03	No	NOAA SQRT
Vanadium	ug/L	8.7	5.85	19	6	6	0	100	0.46	0.31	No	Tier II ^a
Zinc	ug/L	48.6	30.04	30	6	3	3	50	1.62	1.00	Yes	LCV ^b
Radionuclides												
Ra-226	pCi/L	0.822	0.43	NA	6	6	NA	100	NA	NA	NA	
Ra-228	pCi/L	1.247	0.60	NA	5	5	NA	100	NA	NA	NA	
Th-232	pCi/L	0.17	0.10	NA	6	6	NA	100	NA	NA	NA	
U-234	pCi/L	0.572	0.39	NA	6	6	NA	100	NA	NA	NA	
U-235	pCi/L	0.036	0.02	NA	6	6	NA	100	NA	NA	NA	
U-238	pCi/L	0.582	0.34	NA	6	6	NA	100	NA	NA	NA	

NA: Not applicable; radionuclides were screened using DOE's RAD-BCG model

^a USEPA 1996; AWQC = National ambient water quality criteria; Tier II = Values calculated using Great Lakes Water Quality Initiative Tier II methodology.

^b ORNL = Oak Ridge National Laboratory (Suter and Tsao 1996); LCV - Lowest Chronic Value; Tier II = Values calculated using GLWQI Tier II methodology.

NOAA SQRT: Screening Quick Reference Tables, NOAA 1999.

TABLE 4-15
SERA Food Web Model Results for STP River
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Raccoon			Mink			Great blue heron			Mallard		
	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC
Inorganics												
Aluminum	0.57	0.11	0.26	8.01	1.60	3.58	24.53	4.91	10.97	1.22	0.24	0.55
Antimony	0.40	0.08	0.18	1.62	0.32	0.73	NA	NA	NA	NA	NA	NA
Arsenic	0.03	<0.01	0.01	0.13	0.03	0.06	0.21	0.09	0.13	0.42	0.17	0.27
Barium	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Beryllium	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cadmium	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.02	<0.01	<0.01	0.11	<0.01	0.03
Chromium	0.02	<0.01	0.01	0.04	<0.01	0.02	0.72	0.14	0.32	0.86	0.17	0.38
Copper	<0.01	<0.01	<0.01	0.03	0.02	0.02	0.05	0.04	0.04	0.07	0.05	0.06
Iron	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Lead	0.02	<0.01	<0.01	0.04	<0.01	0.01	0.45	0.09	0.20	3.12	0.31	0.99
Manganese	0.15	0.05	0.09	0.62	0.19	0.35	0.35	0.07	0.16	0.16	0.03	0.07
Mercury	0.58	0.35	0.45	3.48	2.09	2.69	128.36	42.79	74.11	70.88	23.63	40.92
Nickel	<0.01	<0.01	<0.01	0.03	0.01	0.02	0.07	0.05	0.06	0.04	0.03	0.03
Selenium	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Silver	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Thallium	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Vanadium	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Zinc	0.02	<0.01	<0.01	0.06	0.01	0.03	0.57	0.06	0.19	1.65	0.18	0.55
Semivolatile Organics												
Benzo(a)anthracene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.02	<0.01	<0.01
Benzo(a)pyrene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Chrysene	12.28	2.46	5.49	8.83	1.77	3.95	5.41	1.08	2.42	39.30	7.86	17.57
Fluoranthene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.03	<0.01	0.02	0.14	0.03	0.06
Fluorene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Indeno(1,2,3-cd)pyrene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Phenanthrene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Pyrene	68.64	13.73	30.70	49.35	9.87	22.07	30.26	6.05	13.53	219.71	43.94	98.26
Volatile Organics												
Toluene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	NA	NA	NA	NA	NA	NA

Bolded HQs indicate exceedance of 1.0

TABLE 4-16

Results of Rad-BCG Screening, STP Upland Soils Maximum Concentrations
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Nuclide	Water (pCi/L)		Sediment (pCi/g)	
	Partial Fraction	Source of Calculation	Partial Fraction	Source of Calculation
Ce-144				
Cs-135				
Cs-137			7.1E-03	TA-Lumped, Default
Co-60				
Eu-154				
Eu-155				
H-3				
I-129				
I-131				
Pu-239				
Ra-226			2.0E-01	TA-Lumped, Default
Ra-228			1.0E+01	TA-Lumped, Default
Sb-125				
Sr-90				
Tc-99				
Th-232			2.0E-01	TA-Lumped, Default
U-233				
U-234			2.5E-03	TA-Lumped, Default
U-235			2.9E-04	TA-Lumped, Default
U-238			7.6E-03	TA-Lumped, Default
Zn-65				
Zr-95				
Partial fractions			1.1E+01	
Result:	1.1E+01			
You have failed the terrestrial site screen				

TA: Terrestrial Animal

TABLE 4-17
 Results of Rad-BCG Screening, STP Upland Soils Mean Concentrations
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Nuclide	Water (pCi/L)		Sediment (pCi/g)	
	Partial Fraction	Source of Calculation	Partial Fraction	Source of Calculation
Ce-144				
Cs-135				
Cs-137			2.9E-03	TA-Lumped, Default
Co-60				
Eu-154				
Eu-155				
H-3				
I-129				
I-131				
Pu-239				
Ra-226			3.3E-02	TA-Lumped, Default
Ra-228			1.8E-01	TA-Lumped, Default
Sb-125				
Sr-90				
Tc-99				
Th-232			4.3E-03	TA-Lumped, Default
U-233				
U-234			2.4E-04	TA-Lumped, Default
U-235			2.8E-05	TA-Lumped, Default
U-238			8.3E-04	TA-Lumped, Default
Zn-65				
Zr-95				
Partial fractions			2.2E-01	
Total sum of fractions:	2.2E-01			
You have passed the terrestrial site screen				

TA: Terrestrial Animal

TABLE 4-18

Comparison of Concentrations of Detected Analytes in STP Upland Surface Soil to Ecological Benchmark Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

ParamName	Units	Maximum Detection	Arithmetic Mean Concentration (1/2 ND)	Screening Criteria	Number of Analyses	Number of Detections	Number of Detections Exceeding Criteria	Detection Frequency (%)	Maximum Hazard Quotient	Mean Hazard Quotient	COPC?	Criteria Source
Metals												
Aluminum	mg/kg	19100	9306.40	50	161	161	161	100	382.00	186.13	No*	ORNL
Antimony	mg/kg	12	5.27	5	121	8	7	7	2.40	1.05	No*	ORNL
Arsenic	mg/kg	90.7	16.84	9.9	161	161	103	100	9.16	1.70	Yes	ORNL
Barium	mg/kg	514	105.79	283	161	161	5	100	1.82	0.37	Yes	ORNL
Beryllium	mg/kg	2.2	0.47	10	161	28	0	17	0.22	0.05	No	ORNL
Cadmium	mg/kg	3.4	0.67	4	161	18	0	11	0.85	0.17	No	ORNL
Chromium, Total	mg/kg	68.9	16.00	0.4	161	161	161	100	172.25	40.01	Yes	ORNL
Cobalt	mg/kg	35	7.48	20	161	155	1	96	1.75	0.37	Yes	ORNL
Copper	mg/kg	591	48.18	60	161	161	29	100	9.85	0.80	Yes	ORNL
Iron	mg/kg	67500	22561.30	200	161	161	161	100	337.50	112.81	Yes	ORNL
Lead	mg/kg	1160	52.37	40.5	161	161	47	100	28.64	1.29	Yes	ORNL
Manganese	mg/kg	2140	540.66	100	161	161	161	100	21.40	5.41	Yes	ORNL
Mercury	mg/kg	2.7	0.17	0.0005	161	66	66	41	5400.00	349.12	Yes	ORNL
Nickel	mg/kg	47.1	18.51	30	161	151	6	94	1.57	0.62	Yes	ORNL
Selenium	mg/kg	1.5	0.42	0.21	161	11	11	7	7.14	1.99	Yes	ORNL
Silver	mg/kg	19.7	1.09	2	161	10	6	6	9.85	0.55	Yes	ORNL
Thallium	mg/kg	2.2	0.40	1	161	2	1	1	2.20	0.40	No	ORNL
Vanadium	mg/kg	41.7	22.48	2	161	161	161	100	20.85	11.24	Yes	ORNL
Zinc	mg/kg	1400	115.95	8.5	161	161	161	100	164.71	13.64	Yes	ORNL
Semivolatiles												
Anthracene	mg/kg	0.13	0.19	0.1	22	2	1	9	1.30	1.86	Yes	Beyer 1990
Benzo(a)Anthracene	mg/kg	0.46	0.18	0.1	22	10	3	45	4.60	1.79	Yes	Beyer 1990 (surrogate)
Benzo(a)Pyrene	mg/kg	0.31	0.18	0.1	22	5	2	23	3.10	1.83	Yes	Beyer 1990
Benzo(b)Fluoranthene	mg/kg	0.74	0.22	0.1	22	9	6	41	7.40	2.19	Yes	Beyer 1990 (surrogate)
Benzo(g,h,i)Perylene	mg/kg	0.24	0.20	0.1	22	2	2	9	2.40	1.98	Yes	Beyer 1990 (surrogate)
Benzo(k)Fluoranthene	mg/kg	0.74	0.22	0.1	22	7	6	32	7.40	2.19	Yes	Beyer 1990 (surrogate)
Butyl benzyl phthalate	mg/kg	0.053	0.18	100	22	3	0	14	0.00	0.00	No	ORNL
Bis(2-Ethylhexyl) Phthalate	mg/kg	1.5	0.44	100	22	7	0	32	0.02	0.00	No	ORNL
Chrysene	mg/kg	0.36	0.18	0.1	22	8	3	36	3.60	1.79	Yes	Beyer 1990 (surrogate)
Dibenz(a,h)Anthracene	mg/kg	0.072	0.18	0.1	22	2	0	9	0.72	1.84	No	Beyer 1990 (surrogate)
Di-N-Butyl Phthalate	mg/kg	0.052	0.18	200	22	2	0	9	0.000	0.001	No	ORNL
Di-N-Octylphthalate	mg/kg	0.2	0.19	200	22	3	0	14	0.001	0.001	No	ORNL (surrogate)
Fluoranthene	mg/kg	0.92	0.23	0.1	22	10	7	45	9.20	2.28	Yes	Beyer 1990 (surrogate)
Indeno(1,2,3-c,d)Pyrene	mg/kg	0.28	0.20	0.1	22	2	2	9	2.80	1.98	Yes	Beyer 1990 (surrogate)
Phenanthrene	mg/kg	0.44	0.16	0.1	22	10	2	45	4.40	1.58	Yes	Beyer 1990
Pyrene	mg/kg	0.63	0.19	0.1	22	10	5	45	6.30	1.94	Yes	Beyer 1990

TABLE 4-18
 Comparison of Concentrations of Detected Analytes in STP Upland Surface Soil to Ecological Benchmark Values
 Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

ParamName	Units	Maximum Detection	Arithmetic Mean Concentration (1/2 ND)	Screening Criteria	Number of Analyses	Number of Detections	Number of Detections Exceeding Criteria	Detection Frequency (%)	Maximum Hazard Quotient	Mean Hazard Quotient	COPC?	Criteria Source
Volatiles												
4-Nitrophenol	mg/kg	0.064	0.46	7	22	1	0	5	0.01	0.07	No	ORNL
Toluene	mg/kg	0.52	0.05	0.05	22	18	5	82	10.40	1.06	Yes	Beyer 1990
Radionuclides												
Cs-137	pCi/g	0.147	0.06	NA	6	6	NA	100	NA	NA	NA	
Ra-226	pCi/g	9.86	1.76	NA	156	144	NA	92	NA	NA	NA	
Ra-228	pCi/g	445.9	7.55	NA	156	149	NA	96	NA	NA	NA	
Th-232	pCi/g	302.16	6.47	NA	146	146	NA	100	NA	NA	NA	
U-234	pCi/g	12.8	1.24	NA	146	136	NA	93	NA	NA	NA	
U-235	pCi/g	0.81	0.08	NA	148	142	NA	96	NA	NA	NA	
U-238	pCi/g	12	1.30	NA	146	143	NA	98	NA	NA	NA	

No: constituent eliminated due to high background concentrations

NA: Not applicable; radionuclides were screened using DOE's RAD-BCG model

ORNL: Oak Ridge National Laboratory (Efroymson et al. 1997); Beyer 1990 -- from Friday, G.P., November, 1998. Ecological Screening Values for Surface Water, Sediment, and Soil. Westinghouse Savannah River Company, Savannah River Technology Center (WSRC-TR-98-001), Aiken, SC 29808
 (Beyer, W.N. 1990. Evaluating soil contamination. U.S. Fish Wildl. Serv., Biol. Rep. 90(2). 25 p.)

TABLE 4-19

SERA Food Web Model Results for STP Upland
Kress Creek and Sewage Treatment Plant Sites - West Chicago, IL

Chemical	Least shrew			Deer mouse			American robin		
	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC	NOAEL	LOAEL	MATC
Inorganics									
Aluminum	6.49	1.30	2.90	0.84	0.17	0.38	3.00	0.60	1.34
Antimony	0.49	0.10	0.22	0.51	0.10	0.23	NA	NA	NA
Arsenic	23.05	4.61	10.31	19.83	3.97	8.87	2.11	0.70	1.22
Barium	2.17	0.56	1.10	0.92	0.24	0.47	0.54	0.27	0.38
Beryllium	0.47	0.09	0.21	0.03	<0.01	0.02	NA	NA	NA
Cadmium	16.20	1.62	5.12	1.55	0.16	0.49	6.76	0.49	1.82
Chromium	7.83	1.57	3.50	0.61	0.12	0.27	15.87	3.17	7.10
Cobalt	0.26	0.06	0.13	0.03	<0.01	0.01	NA	NA	NA
Copper	1.38	1.03	1.19	0.31	0.23	0.27	1.54	1.17	1.34
Iron	NA	NA	NA	NA	NA	NA	NA	NA	NA
Lead	26.23	2.62	8.30	4.91	0.49	1.55	36.18	7.24	16.18
Manganese	0.42	0.13	0.23	0.33	0.10	0.18	0.04	<0.01	0.02
Mercury	203.86	40.77	91.17	32.41	6.48	14.49	8.27	3.38	5.28
Nickel	0.66	0.33	0.46	0.12	0.06	0.08	0.22	0.16	0.18
Selenium	1.20	0.73	0.93	1.11	0.67	0.86	0.46	0.13	0.25
Silver	3.91	0.78	1.75	0.26	0.05	0.12	3.04	0.61	1.36
Thallium	7.13	1.43	3.19	0.52	0.10	0.23	0.49	0.10	0.22
Vanadium	1.30	0.26	0.58	0.20	0.04	0.09	0.06	0.01	0.02
Zinc	13.22	6.61	9.35	1.59	0.79	1.12	89.49	9.91	29.77
Semivolatile Organics									
4-Nitrophenol	NA	NA	NA	NA	NA	NA	NA	NA	NA
Anthracene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Benzo(a)anthracene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Benzo(a)pyrene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Benzo(b)fluoranthene	0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Benzo(g,h,i)perylene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Benzo(k)fluoranthene	0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
bis(2-Ethylhexyl)phthalate	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.11	0.02	0.05
Chrysene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Dibenz(a,h)anthracene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Di-n-butylphthalate	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.02	<0.01	<0.01
Di-n-octylphthalate	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	NA	NA	NA
Fluoranthene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Indeno(1,2,3-cd)pyrene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Phenanthrene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Pyrene	0.02	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Volatile Organics									
Toluene	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	NA	NA	NA

Bolded HQs indicate exceedance of 1.0

Figures




Figure 1-1
 Site Location Map
 Kress Creek/West Branch DuPage River Site
 Sewage Treatment Plant Site




Figure 3-1
Wildlife Survey Sampling Locations
 Kress Creek/West Branch DuPage River to Warrenville Dam

Kress Creek - West Branch Dupage River Watershed
 Watershed Overview

Kress Creek RI
 West Chicago, Illinois


- Aquatic Invertebrate Survey Area
- Terrestrial Survey Area
- Fish Survey Area

Water
 EG&G 1989 flyover boundary

Scale 1:18000
 1 in = 1500 feet



CRS/LL
 EPA










FIGURE 3-2
Wildlife Survey Sampling Locations
West Branch DuPage River from STP to Confluence
Sewage Treatment Plant
West Chicago, Illinois


-  Aquatic Invertebrate Survey Area
-  Terrestrial Survey Area
-  Fish Survey Area



1 inch = 500 feet


CH2MHILL

Figure 3-3. Conceptual Site Model for Radionuclides
Kress Creek -- West Chicago, IL




Legend:

- X - Potentially complete exposure pathway (quantitatively evaluated)
- O - Potentially complete exposure pathway but not evaluated quantitatively
- (1) -- Detected in surface soils, sediment and surface water
- - Not applicable or incomplete

Figure 3-4. Conceptual Site Model for Chemical Contaminants

Kress Creek - West Chicago, IL




Legend:

- X - Potentially complete exposure pathway (quantitatively evaluated)
- O - Potentially complete exposure pathway but not evaluated quantitatively
- Not applicable or incomplete
- (1) -- Detected in sediments only (not analyzed for in soils and surface water)
- (2) -- Detected in surface soils, sediment and surface water
- Dashed lines indicate theoretically complete pathways but unanalyzed media

Figure 3-5. Conceptual Site Model for Radionuclides

STP River -- West Chicago, IL




Legend:

- X - Potentially complete exposure pathway (quantitatively evaluated)
- O - Potentially complete exposure pathway but not evaluated quantitatively
- (1) - Detected in surface soils, sediment and surface water
- - Not applicable or incomplete

Figure 3-6. Conceptual Site Model for Chemical Contaminants

STP River -- West Chicago, IL




Legend:

- X - Potentially complete exposure pathway (quantitatively evaluated)
- O - Potentially complete exposure pathway but not evaluated quantitatively
- Not applicable or incomplete
- (1) -- Detected in sediments only (not analyzed for in soils and surface water)
- (2) -- Detected in surface soils, sediment and surface water
- Dashed lines indicate theoretically complete pathways but unanalyzed media

Figure 3-7. Conceptual Site Model for Radionuclides

STP Upland -- West Chicago, IL




Legend:

- X - Potentially complete exposure pathway (quantitatively evaluated)
- O - Potentially complete exposure pathway but not evaluated quantitatively
- (1) -- Detected in surface soils, sediment and surface water
- - Not applicable or incomplete

Figure 3-8. Conceptual Site Model for Chemical Contaminants

STP Upland - West Chicago, IL



Legend:

- X - Potentially complete exposure pathway (quantitatively evaluated)
- O - Potentially complete exposure pathway but not evaluated quantitatively
- Not applicable or incomplete
- (1) -- Detected in sediments only (not analyzed for in soils and surface water)
- (2) -- Detected in surface soils, sediment and surface water
- Dashed lines indicate theoretically complete pathways but unanalyzed media