

US EPA ARCHIVE DOCUMENT

Table 1. Children's food consumption data used in this analysis.

Age	Children Included In CSFII Survey	Valid Eating Days
6-12 mo.	274	572
1	842	1,732
2	868	1,768
3	726	1,448
4	701	1,410
5	671	1,372
TOTAL	4,082	8,302

Source: USDA Continuing Survey of Food Intake by Individuals. Survey Years: 1989-91, 1994-95.

Table 2. Pesticide residue data used in this analysis.

Database	Source	Dates	Number of foods used in analysis	Number of samples used in analysis	Attributes
Total Diet Study	FDA	1991-96	90	1,459	cooked and prepared food washed and peeled * raw commodities/reduction factors applied (see table 5)
Pesticide Data Program	USDA	1992-95	14	25,886	
Pesticide Monitoring Database	FDA	1992-96	68	23,238	
TOTAL			172**	50,583	

*90% reduction factor was applied to green bean results from the PDP because they are tested raw but eaten cooked.

** These 172 foods were linked to 407 distinct foods in the USDA food consumption database.

Source: EWG, compiled from USDA food consumption data 1989-1995, USDA and FDA pesticide residue data 1991-1996 and reference doses (RfDs) obtained from EPA in January 1998.