

US EPA ARCHIVE DOCUMENT


About Issues in Ecology

Issues in Ecology is designed to report, in language understandable by non-scientists, the consensus of a panel of scientific experts on issues relevant to the environment. *Issues in Ecology* was initially supported by the Pew Scholars in Conservation Biology program and is currently supported by the Ecological Society of America. It is published at irregular intervals, as reports are completed. All reports undergo peer review and must be approved by the Editorial Board before publication. No responsibility for the views expressed by authors in ESA publications is assumed by the editors or the publisher, the Ecological Society of America.

Issues in Ecology is an official publication of the Ecological Society of America, the nation's leading professional society of ecologists. Founded in 1915, ESA seeks to promote the responsible application of ecological principles to the solution of environmental problems. For more information, contact the Ecological Society of America, 1707 H Street, NW, Suite 400, Washington, DC, 20006. ISSN 1092-8987



Impacts of Atmospheric Pollution on Aquatic Ecosystems



Impacts of Atmospheric Pollutants on Aquatic Ecosystems

SUMMARY

Considerable progress has been made in reducing the discharge of atmospheric pollutants from point sources such as effluent pipes. A more difficult challenge involves identifying and controlling environmental contaminants generated by dispersed or nonpoint sources such as automobile exhaust, pesticide applications, and myriad commercial and industrial processes. Nonpoint pollutants can travel far from their sources when they are discharged into rivers or enter the atmosphere. While waterborne contaminants have received growing attention, little recognition has so far been given to the far-ranging environmental consequences of toxic substances and nutrients that are transported via the air.

This report reviews three categories of airborne pollutants that we consider of greatest concern, both for their ecological effects and their impacts on the health of fish, wildlife, and humans:

- **Organic compounds:** These include long-recognized persistent organic pollutants and a vastly larger group of chemicals such as brominated flame retardants, water-repellent coatings, and synthetic fragrances that remain largely unmonitored and unregulated.
- **Mercury:** Oxidized forms of mercury readily rain from the air onto terrestrial and aquatic ecosystems. In sediments, they can be transformed into monomethyl mercury, the form most toxic to fish and the wildlife and humans that consume fish.
- **Nutrients:** Atmospheric transport is a significant and increasing source of plant nutrients to freshwater and marine ecosystems and can accelerate eutrophication of these waters.

A review of the available scientific information indicates that:

- The pollutants that are most likely to present ecological risks are those that are (1) highly bioaccumulative, building up to high levels in animal tissues even when concentrations in the water remain relatively low, and (2) highly toxic, so that they cause harm at comparatively low doses.
- Atmosphere-water interactions that control the input and outgassing of persistent organic pollutants in aquatic systems are critically important in determining the cycling and residence times of these compounds and the extent of contamination of food webs.
- Although the effects of various types of pollutants are usually evaluated independently, many regions are subject to multiple pollutants, and their fate and impacts are intertwined. The effects of nutrient deposition on coastal waters, for instance, can alter how various organic contaminants and mercury are processed and bioaccumulated, and ultimately, how they affect aquatic organisms.
- For many organic pollutants, even long-banned chemicals such as PCBs and other organochlorines, non-atmospheric sources have been well controlled while atmospheric sources have either been neglected or ignored.
- Ecological effects of airborne organochlorines are a particular concern at high latitudes and altitudes. Even though concentrations of organochlorines in air masses and snow from northern and alpine regions are generally low, the food web dynamics, physiologies, and life cycles of cold region animals allow these contaminants to be biomagnified to extraordinary degrees in food chains.

Atmospherically deposited contaminants are generated largely by human activities, and reducing the extent and impacts of this increasingly significant source of environmental pollution will require greater recognition, monitoring, and ultimately, regulation.

Cover Photo: Fog over Lonesome Point on Lake Superior, Grand Marais, MI (courtesy the U.S. Environmental Protection Agency and U.S. Fish and Wildlife Service).

Impacts of Atmospheric Pollutants on Aquatic Ecosystems

by

Deborah L. Swackhamer, Hans W. Paerl, Steven J. Eisenreich, James Hurley, Keri C. Hornbuckle, Michael McLachlan, David Mount, Derek Muir, and David Schindler

INTRODUCTION

Over the past several decades, the United States has made considerable progress in reducing the amount of pollutants discharged from identifiable point sources such as municipal effluent pipes. A more difficult challenge has been to identify and control environmental contaminants generated by dispersed or nonpoint sources such as automobile exhaust, livestock wastes, fertilizer and pesticide applications, and myriad commercial and industrial processes. These nonpoint pollutants can travel far from their sources when they seep or flow into rivers or enter the air. In particular, volatile chemicals – those that evaporate readily – can be carried through the atmosphere and fall on parts of the world far removed from their origins. They can either be deposited directly onto terrestrial and aquatic ecosystems (“direct” deposition) or deposited onto land surfaces and subsequently run off and be transferred into downstream waters (“indirect” deposition). Deposition of these pollutants can occur via wet or dry forms. Wet deposition includes rain, snow, sleet, hail, clouds, or fog, while dry deposition includes gases, dust, and minute particulate matter. Rates of wet deposition are most influenced by how readily the chemicals dissolve in water, while rates of dry deposition are very sensitive to the form (gas or particle) of the chemicals and the “stickiness” of the surface upon which they are being deposited. Chemicals deposited to aquatic ecosystems can re-volatilize and thus be redistributed via the atmosphere. During atmospheric transport, pollutants also can be transformed into other chemicals, some of which are of greater concern than those originally released to the atmosphere. Pollutants may also be transformed into other chemicals once they are deposited on and travel through watersheds. Until recently, however, little recognition has been given to the environmental consequences of toxic substances and nutrients that fall from the air as wet and dry deposition onto land-based and aquatic ecosystems.

Since air moves rapidly, atmospheric pollutants can travel long distances quickly and be deposited on distant watersheds. The “airshed” for a particular body of water can encompass hundreds of miles. An airshed defines the geographic area that contains the emissions sources that contribute 75 percent of the pollutants deposited in a particular watershed¹ (Figure 1). Airsheds differ for each form of every pollutant and are determined by modeling atmospheric deposition of each chemical. They are useful theoretical tools for explaining atmospheric transport and for illustrating the need to control emission sources far removed from the ecosystem of concern.

This report reviews three categories of atmospheric pollutants that we consider of greatest concern, both for their ecological effects and their impacts on the health of a wide range of biota, including lower levels of the food web (algae, macrophytes, and invertebrates), fish, wildlife, and humans. These categories include organic compounds, mercury, and inorganic nutrients.

First, semi-volatile organic contaminants often have properties that allow them to persist in the environment for very long periods, to bioaccumulate (that is, build up in animal tissues), and to be toxic to aquatic organisms at lower levels of the food web, as well as to fish and to the wildlife and humans that eat fish. These persistent organic pollutants include a wide range of chemicals from pesticides and polychlorinated biphenyls (PCBs) to brominated flame-retardants, water- and stain-repellent coatings, and synthetic fragrances.

Second, the metal mercury can be transported in the atmosphere and fall onto terrestrial and aquatic ecosystems as precipitation or dry deposition. In aquatic systems, mercury may eventually be transformed into monomethyl mercury, a form that is bioaccumulative and can harm fish, wildlife, and humans.

Finally, the significance of inorganic forms of nutrients as atmospheric pollutants has been gaining increased attention. Nutrient-laden runoff from the land has long been acknowledged as a culprit in the over-enrichment and eutrophication of coastal




Figure 1 – Principal nitrogen oxide airsheds and corresponding watersheds for Hudson/Raritan Bay, Chesapeake Bay, Pamlico Sound, and Altamaha Sound (listed from north to south).² These airsheds show the geographic area that contains the emissions sources that contribute 75 percent of the nitrogen oxide deposited in each watershed. Via atmospheric transport, pollutants such as nitrogen oxide can impact watersheds hundreds of miles away.

waters. Now, atmospheric nitrogen deposited in coastal and estuarine waters has been shown to be a major nutrient source in some coastal regions. The result can be excessive algal (phytoplankton) growth, oxygen depletion, degradation of marine habitats, and loss of both biodiversity and commercially valuable fish and shellfish species.

The properties that determine whether or not a chemical is likely to become a “problem” in aquatic ecosystems include its intrinsic toxicity, how long it can persist in air without decomposing (or without transforming to a chemical of greater concern), whether it bioaccumulates, how it interacts with other chemicals, whether it re-volatilizes, and how it is transformed once deposited in water.

Usually, the emission, airborne transport, fate, and ecological impacts of these three classes of pollutants are considered independently. However, while these contaminants may be generated by different sources, their impacts on the environment cannot be evaluated separately. Many coastal regions are subject to pollution from multiple sources, and the atmospheric deposition of nutrients often occurs in concert with deposition of mercury and one or more organic contaminants. Thus, the effects of nutrients on coastal ecosystems and their food webs can alter how various organic contaminants and mercury are processed, how they build up in the food web, and ultimately, how these toxic chemicals affect fish, wildlife, and humans.

The first section of this report examines these three classes of pollutants, their characteristics, and sources. The second section explores atmosphere-water interactions that determine the fate and persistence of airborne pollutants in freshwater and marine ecosystems. The third discusses the factors that determine whether atmospherically delivered pollutants present a risk to fish, wildlife, and humans. The fourth section looks at the relationship between nutrient deposition and the fate and impact of organic pollutants. The fifth and final section outlines priorities for regulation and monitoring of atmospheric pollutants.

POLLUTANTS OF CONCERN

Organic Compounds

The organic compounds that merit concern as atmospheric pollutants have diverse chemical structures, sources, and uses. They can generally be categorized either as deliberately produced substances such as pesticides, industrial compounds, and their persistent degradation products, or as byproducts of fossil fuel combustion or impurities in the synthesis of other chemicals. Although diverse structurally, the organic chemicals that are transported atmospherically, deposited into remote environments, and build up to levels that can affect wildlife and human health, have a relatively narrow range of physical and chemical properties (see Box 1). These are properties that (1) allow them to move in measurable quantities from land and water surfaces to the atmosphere, (2) give them sufficient stability (in the form of resistance to degradation by ultraviolet light and oxidation by hydroxyl radicals) to be transported long distances, and (3) impart a relatively high affinity for fatty tissues and resistance to breakdown in the body and thus allow them to accumulate in

organisms and biomagnify (increase in concentration as they move up) in food chains.

Most atmospherically-transported chemicals that also bioaccumulate, such as PCBs and chlorobenzenes, are known as “multimedia chemicals” because they can be distributed through air, water, and soil rather than a single medium. Virtually all of the persistent organic pollutants listed under the Stockholm Convention — aldrin, chlordane, dieldrin, dichlorodiphenyltrichloroethane (DDT), endrin, heptachlor, hexachlorobenzene, mirex, toxaphene, PCBs, polychlorinated dibenzop-dioxins and –dibenzofurans (PCDD/Fs) — are multimedia chemicals.⁶ A few highly chlorinated PCDD/F and PCB congeners are solid phase chemicals that concentrate solely in soils and sediments. (Congeners are members of a family of chemicals that have the same basic structure but have different amounts of chlorine.)

Persistent organic pollutants, as defined by the Stockholm Convention, are now scheduled for either global bans (chlorinated pesticides) or emission reductions (by-products such as PCDD/Fs). Nevertheless, the risk they present to the environment will persist because of their extraordinary resistance to degradation and because contaminated sources such as agricultural soils or PCB-containing building materials continue to re-supply the atmosphere. In addition, the Priority Substances List in the European Water Framework Directive includes many of these same chemicals, as well as polybrominated diphenyl ethers (PBDEs) used as fire retardants and chlorinated alkanes.

Chemicals that accumulate largely in one environmental medium (air, water, or soil) are generally not a concern for ecosystems impacted primarily by atmospheric pollution. For example, the herbicide atrazine is known to be very persistent in nutrient-poor waters, but little of it volatilizes to the atmosphere. Because of this, its impacts are largely of concern locally, for example in agricultural streams and wetlands near fields where atrazine is applied.⁷ Similarly, alkyl phenols and acid pharmaceuticals present an exposure risk to aquatic life in receiving waters near municipal waste treatment plants.⁸ Substantial concentrations of alkyl phenols are also observed in the atmosphere above estuaries receiving wastewater effluents, but these chemicals adhere efficiently to atmospheric aerosols and are soon removed by rainfall.⁹ Thus they travel only short distances in the atmosphere and are generally not a concern for remote aquatic environments where atmospheric deposition is the predominant source of pollution.

It is more difficult to classify the atmospheric pollution potential of the many semi-volatile chemicals that have multimedia characteristics but are rapidly degraded either in the atmosphere or in the biosphere. Examples of this group are the 2, 3 and 4-ring polyaromatic hydrocarbons (PAHs), organophosphorus pesticides, and mono-, di- and trichlorobenzenes. Under some circumstances, concentrations of these chemicals could build up even in remote environments if rates of atmospheric and water degradation are low – for example, in cold climate regions. This might lead to exposure of some aquatic or terrestrial organisms, but these compounds would likely be broken down during metabolism by vertebrates and thus would generally not be expected to build up in food webs. This generality needs to be

Box 1 — Physical and Chemical Properties of Atmospherically Transported Organic Chemicals

The combination of physical properties that give rise to environmentally mobile and bioaccumulative substances is best viewed by a two-dimensional plot of the key partition coefficients (Figure 2). Partition coefficients describe how much of a contaminant will be in one medium (e.g. air) compared to another medium (e.g. water) at equilibrium. For example, if a chemical has an air-water partition coefficient of 2, then there will be twice as much of the chemical in air than in water when expressed in equivalent concentrations. The octanol-water partition coefficient (K_{ow}) is commonly used as an index of toxicity because solubility in octanol mimics solubility in biological lipid tissues and indicates the potential for bioaccumulation. Van de Meent et al.³ proposed classifying chemicals as either (A) gas phase chemicals that partition into the gas phase regardless of their mode of entry into the environment, (B) aqueous phase chemicals that partition into the aqueous environment regardless of mode of entry, (C) solid phase chemicals that partition in to soils and sediments, and (D) multimedia chemicals that partition into more than one environmental medium. To visualize these categories, a global scale multimedia model (similar to GloboPOP⁴) was applied that assumed no degradation except in air (class A), water (class B), and soil (class C). The shaded areas in Figure 2 reflect substances with a wide range of air-water and octanol-water partition coefficients, which indicate their relative affinity for air vs. water or for the lipid tissues of organisms vs. water, respectively.




Figure 2 – Plot of the two key partition coefficients, air-water partition coefficients ($\log K_{aw}$) and octanol-water partition coefficients ($\log K_{ow}$), illustrating predicted environmental media (gas—air, aqueous—water, and solid—soil) where organic contaminants accumulate or are transported as a function of their physical chemical properties⁵. Many toxic chemicals are multimedia and partition into more than one medium.

assessed on a case-by-case basis, however, since our ability to predict such biotransformations in the food web is weak.¹⁰

New emerging organic contaminants of interest

Since the late 1990s, there has been a major increase in measurement and detection of organic chemicals that are not presently classified as persistent organic pollutants in waters affected by atmospheric contaminants. These chemicals include:

- polybrominated diphenyl ether flame retardants (PBDEs) widely used in polymers and textiles;
- fluorinated surfactants used to make hundreds of everyday products from non-stick cookware and water- and stain-repellent coatings for carpets and raincoats to cosmetics, paper products, and polymers for electronics;
- chlorinated naphthalenes (PCNs) used in cable insulation, wood preservation, electronics manufacturing, and dye production;
- chlorinated alkanes (also known as chlorinated paraffins) found in paints and adhesives as well as fluids used in cutting and machining metals; and
- pesticides currently in use such as endosulfan and lindane.

Even this expanded list, however, represents only a tiny fraction of the chemicals in commerce or even of the subset known as “high production volume chemicals” (HPVCs). The U.S. HPVC

list includes 2,863 organic chemicals produced or imported at levels greater than 450 tons per year.¹¹ In the European Union, the European Inventory of Existing Commercial Chemical Substances lists 100,195 “existing chemicals” – meaning chemicals in commerce as of 1981 — of which about 2,704 are considered HPVCs based on production levels greater than 1,000 tons per year and 7,842 are low production volume chemicals produced at rates of 10 to 1,000 tons per year.¹² The Organization for Economic Cooperation and Development maintains an HPVC list based on a compilation of the U.S., E.U., and other national inventories. In 2000, that list contained 5,235 substances produced at levels greater than 1,000 tons globally.

While the majority of these HPVCs are probably not a concern with regard to their environmental persistence, bioaccumulation, and toxicity, the chemical industry has recognized that data are lacking for many of these chemicals. In the absence of data, production volume is assumed to be a surrogate for occupational, consumer, and environmental exposure.¹³ The International Council of Chemical Associations has established a list of 1,000 HPVCs for which full data sets on toxicity and environmental fate are to be developed by 2004.¹⁴ However, this will leave more than 50 percent of high production volume chemicals without full data sets.

Among the new organic contaminants of greatest concern are synthetic musk fragrances, PBDE flame retardants, and fluorinated surfactants.

Synthetic Fragrances. Synthetic musk fragrances are semi-volatile and lipophilic (literally “fat-loving” because they are attracted to fatty tissues) compounds that are added to a wide range of personal care products, including perfumes, cosmetics, soaps, and shampoos as well as laundry detergents.¹⁵ These synthetic fragrances are on the U. S. HPVC list but have only recently been studied as contaminants in any natural system in this country. The most common synthetic fragrances used are two nitro musks called musk xylene and musk ketone and two polycyclic musks known as HHCB (hexahydro-hexamethylcyclopentabenzopyran) and AHTN (hexamethyl-tetraline). In Europe, approximately 6,500 metric tons of these four synthetic compounds were produced in 1999 for use as consumer product additives.¹⁶

In the early 1980s, concentrations of synthetic musk fragrances were discovered in animal tissues for the first time. Since then, there has been an increasing awareness of the ubiquitous distribution and possible toxicological effects of these compounds. Recent measurements of these compounds in wastewater effluent and in air and water in the Great Lakes region, for instance, have illustrated that ecological exposures are chronic and likely to be increasing.¹⁷ This is cause for concern because both HHCB and AHTN have been shown to exhibit hormonal disruption in fish.¹⁸ (Hormonally active substances are chemicals that mimic or interfere with hormone function and can distort normal reproductive development, alter behavior, and impair disease resistance in wildlife and humans.) Several studies with cell cultures indicate that musk xylene, musk ketone, p-amino-musk xylene (a major breakdown product of musk xylene), and the polycyclic musk fragrance AHTN all demonstrate estrogenic activity in laboratory tests. In Europe, musk ketone and musk xylene were effectively banned from use as fragrances in 2002 because of their reported toxicities.¹⁹

Although HHCB and AHTN are both on the U. S. HPVC list, their use in personal care and household products is privileged information in the United States, and companies that use them do not have to report how much they use or manufacture. They also do not have to report any estimates on how much synthetic fragrance may ultimately be discharged into the environment. Because of this, ecological impacts of these compounds can only be identified through field and toxicological studies conducted long after exposures have begun. Fortunately, thanks to the intense interest in the fate and impacts of these compounds in Europe, analytical methods have been developed and standards are available for these fragrances.

For the vast majority of high production volume chemicals identified as potentially bioaccumulative and persistent, however, there are no trace analytical methods available for tracking their fate and impacts.²⁰ Many of the recently initiated measurements of organic chemicals have been made using advances in analytical methodology, especially in the case of fluorinated organics.

Flame retardants. Among the newly emerging chemical contaminants of aquatic environments, the PBDE flame retardants and the perfluorinated surfactants discussed below have generated

the greatest concern. PBDEs are used in thousands of consumer products from fire-resistant textiles and upholstered furniture to computers and televisions. Global demand for these additives increased from 40,000 tons in 1992 to 67,125 tons in 1999.²¹ The tetra- and pentaBDEs (TeBDE and PeBDE) are of greatest concern, and their concentrations are increasing in humans and wildlife.²² TeBDE and PeBDE are multimedia chemicals with physical properties similar to those of some PCBs. A higher brominated product, decabromodiphenyl ether (DecaBDE), is a solid phase chemical, but it may degrade in sunlight and in the tissues of fish to these lower brominated multimedia forms.²³ Researchers measured a nine-fold increase in PBDEs in the tissues of ringed seals from the western Canadian Arctic over the period of 1981 to 2000.²⁴

Fluorinated surfactants. Scientists have recently documented widespread contamination of wildlife and the general human population with perfluorinated acids.²⁵ “Perfluorinated” is a term used to describe organic molecules that are fully fluorinated, meaning fluorine atoms have replaced all hydrogen atoms in the carbon-hydrogen bonds. The most widely known perfluorinated acids are perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA); however, similar compounds having longer or shorter perfluorinated chains are also produced or exist as impurities within manufactured formulations. These important industrial chemicals fall into the category of surfactants because they are surface-active agents that repel water and oil or resist heat or other chemicals. The major use of PFOS is in treating fabric surfaces for stain resistance. The existing database describing physical properties of perfluorinated acids, including PFOS and PFOA, is severely limited because of their anomalous physical and chemical behavior. The properties of PFOS and PFOA suggest that they are poor candidates for long-range airborne transport, yet they have been discovered throughout the global environment. Worldwide dissemination of perfluorinated acids must therefore occur by way of an airborne neutral derivative that yields the free acid when it degrades.²⁶ Widespread detection of precursors of PFOS and PFOA in the air in North America is providing increasing evidence that this is indeed the means by which these nonvolatile compounds have become such widespread contaminants.²⁷

Over the past decade, researchers have found PFOS in birds, fish, and marine and land mammals around the world. For example, PFOS has been detected in the blood of ringed seals from the northern Baltic Sea, the eastern Canadian arctic, and Svalbard; the blood and liver of northern fur seals from Alaska; and the livers of polar bears from northern Alaska.²⁸ PFOS concentrations in polar bear livers range from 1 to 5 micrograms per gram of tissue (wet weight), making it the most prominent organohalogen contaminant in these mammals.²⁹

Mercury

Mercury is a metallic element (Hg) that has been extracted for centuries from sulfide ore or cinnabar (HgS). It has become a global pollutant and can be mobilized into the atmosphere from many human activities, including municipal trash incineration, burning of high sulfur coal (which contains cinnabar) in coal-fired power plants, metal smelting, chlorine-alkali plants, cement

making, and gold extraction, as well as from use of mercury-based fungicides in latex paints and the paper and pulp industry. Mercury in its elemental state has low reactivity and a long atmospheric residence time, thus allowing it to be mixed in the atmosphere on a global scale, while the oxidized forms are removed by wet and dry deposition.³⁰ Oxidized reactive gaseous mercury (RGM), for example, is very soluble in water and is effectively deposited on land and water by snow and rainfall. Particulate forms of mercury fall as dry deposition.³¹

The total mass of mercury in the atmosphere has been estimated at 5,000 to 6,000 metric tons, and approximately half of that was generated by human activities.³² Atmospheric concentrations of mercury peaked in the 1960s and 1970s and have been declining since then.³³ It has been estimated that human activities contribute 70 to 80 percent of the total annual mercury emissions to the atmosphere and that more than 95 percent of mercury vapor in the atmosphere exists as elemental

mercury (Hg).³⁴ The remaining balance of the mercury exists as RGM, as particulate complexes of divalent mercury, and in the organic form as monomethyl mercury.³⁵

Although atmospheric concentrations have been declining for several decades, mass balance calculations that relate net mercury accumulation in the atmosphere with net loss indicate that human inputs of mercury to the atmosphere have increased threefold since the beginning of the industrial age.³⁶ This estimate has been supported by data from several field-based studies of dated sediment cores from lakes and wetlands.³⁷ The mass balance calculations also suggest that a legacy of mercury inputs is stored in terrestrial landscapes since only 5 percent of the atmospheric mercury deposited on the land is carried to the oceans via runoff. Refinement of mass balance calculations has led some researchers to conclude that dry deposition of RGM from the atmosphere can represent up to 35 percent of the total mercury input to the ocean.³⁸

Table 1 - Natural and anthropogenic sources of atmospheric nitrogen compounds (the major chemical forms of atmospheric nitrogen compounds are the reduced, oxidized and organic forms).

Chemical Form	Sources (in approximate order of importance)
<p>Reduced Nitrogen Ammonia/Ammonium (NH₃/NH₄⁺)</p>	<p>Agricultural Livestock waste (volatilized NH₃) Chemical fertilizers (volatilized NH₃) Biomass burning Dust from deforestation & land clearing</p> <p>Urban & Rural (non-agricultural) Wastewater treatment (volatilized NH₃) Fossil fuel combustion (from automobile catalytic converters)</p> <p>Natural Biomass burning (forest and grass fires) Decomposition of organic matter Dust and aerosols Volcanism</p>
<p>Oxidized Nitrogen Nitrogen Oxides (NO/NO₂/NO₃)</p>	<p>Urban & Rural (non agricultural) Fossil fuel combustion mobile & stationary engines powerplants & industrial</p> <p>Natural Biomass burning Lightning Photolysis of N₂O (air, land, water) Dust and aerosols generated by storms Microbially-mediated volatilization</p>
<p>Organic Nitrogen (Dissolved and Particulate)</p>	<p>Agricultural Dust and volatilization of wastes??</p> <p>Urban & Rural (non-agricultural) Dust/aerosols??</p> <p>Natural Atmospheric photochemical and lightning Biological production in oceans??</p>
<p>?? = possible, but little known about, sources</p>	

While the mass balance has identified the magnitude of the various fluxes and pools of mercury and possible pathways for contamination of land and water, it does not provide information on the true partitioning of various forms of mercury in the atmosphere. This information is vital for predictive modeling of global mercury cycling and the effectiveness of mercury reduction strategies, and it continues to be an active topic of research. The potential for atmospheric deposition of mercury, for example, depends upon the distribution of various forms of mercury in emissions and plumes. Both particulate mercury and RGM are likely be deposited closer to their local or regional sources, while gaseous mercury is expected to be transported long range and have a one to two year residence time in the atmosphere. Current instrumentation allows for real-time measurement of atmospheric mercury as RGM, particulate mercury, and gaseous mercury at the picogram or sub-picogram level. The simultaneous measurement of these various atmospheric forms has allowed for analysis of phase distribution of mercury near point sources, at offshore oceanic stations, and in remote areas.

Nutrients

A significant and increasing source of nutrients to freshwater and marine ecosystems is atmospheric deposition, either as rain or snow or as dry deposition of particles and gases. The nutrients that have received most attention are those that are essential for plant growth (primary production) because their concentrations control the growth of algae, which form the base of aquatic food webs. These nutrients include nitrogen, phosphorus, iron, and trace elements such as zinc, manganese, copper, cobalt, molybdenum, boron, and selenium. By far, the greatest attention has been focused on nitrogen because it is the most common limiting nutrient in marine, estuarine, and a few freshwater systems. Nitrogen is also a highly significant component of atmospheric deposition.³⁹ In the marine environment, iron has been the subject of increasing interest because recent studies have shown that this metal limits primary production in some open ocean waters.⁴⁰ Iron can also act synergistically with nitrogen to enhance algal production in coastal and ocean waters.⁴¹ Both nitrogen and phosphorus have received attention in freshwater ecosystems, which are most often phosphorus limited.

Early studies on human-generated contaminants delivered to ecosystems via the atmosphere identified nitrogen as a major nutrient constituent of both rain- and dry-fall.⁴² Atmospherically deposited nitrogen provides aquatic systems with a variety of biologically available nitrogen compounds, reflecting a diverse array of human activities and, to a lesser extent, natural processes (Table 1). These compounds include inorganic reduced forms (ammonia, ammonium), inorganic oxidized forms (nitrogen oxides, nitrate, nitrite), and organic forms (urea, amino acids, and unknown compounds). During the past century, atmospherically deposited nitrogen has increased tenfold, driven by trends in urbanization, industrial expansion, and agricultural intensification.⁴³ Nitrogen deposition ranges from 400 to more than 1,200 kilograms per hectare each year and represents from 10 to more than 40 percent of the “new” nitrogen coming into North American and European inland and coastal waters (Table 2).⁴⁴ On a larger scale, nitrogen

flux to the North Atlantic Ocean basin is approximately 11.2 teragrams (trillion grams) per year and accounts for 46 to 57 percent of its “new” nitrogen input.⁴⁵ This is comparable to the “new” nitrogen inputs delivered to the ocean by rivers.⁴⁶ Indeed, in the waters of the North American continental shelf, nitrogen inputs via the atmosphere exceed those arriving by rivers.⁴⁷

Table 2 - Estimated contributions of atmospheric deposition of nitrogen to “new” nitrogen inputs in diverse estuarine, coastal and open ocean waters. When identified, the sources (wet: W and/or dry deposition: D) and chemical forms (inorganic: I and/or organic: O) of atmospherically deposited nitrogen are indicated⁴⁸.

RECEIVING WATERS	PERCENT OF “NEW” NITROGEN THAT IS ATMOSPHERICALLY DEPOSITED
Baltic Sea (Proper) ⁴⁹	~ 30 W+D, I
Kiel Bight (Baltic) ⁵⁰	40% W, I
North Sea (Coastal) ⁵¹	20-40% W+D, I
Western Mediterranean Sea ⁵²	10 60% W, I
Waquoit Bay, MA, USA ⁵³	29% W, I+O
Narragansett Bay, USA ⁵⁴	12% W, I+O
Long Island Sound, USA ⁵⁵	20% W, I+O
New York Bight, USA ⁵⁶	38% W, I+O
Barnegat Bay, USA ⁵⁷	40% W, I+O
Chesapeake Bay, USA ⁵⁸	27% W, I+O
Rhode River, MD, USA ⁵⁹	40% W, I+O
Neuse River Estuary, NC, USA ⁶⁰	35% W, I+O
Pamlico Sound, NC, USA ⁶¹	~ 40% W+D, I
Sarasota/Tampa Bay, FL, USA ⁶²	30% W+D, I
Mississippi River Plume, USA ⁶³	2-5% W+D, I+O

Excessive nitrogen loading to estuarine and coastal waters is the key cause of accelerating eutrophication and the associated environmental consequences, including algal blooms, decreases in water clarity, toxicity, hypoxia or anoxia (oxygen-depleted or “dead zones”), fish kills, declines in submerged aquatic vegetation, and associated habitat loss.⁶⁴

As a significant source of “new” nitrogen, atmospheric deposition is both a local and regional issue because emission sources may be situated either within or far outside affected watersheds.⁶⁵ Nitrogen oxides, mostly generated by fossil fuel combustion, account for 50 to 75 percent of nitrogen pollution in the United States, with reduced nitrogen and organic nitrogen making up the rest. Rapidly expanding livestock (swine, cattle and poultry) operations in the Midwest and Mid-Atlantic regions have accelerated the generation of nitrogen-enriched wastes and manures, and 30 to 70 percent or more of this may be emitted as ammonia (NH₃) gas. This has led to local and regional increases in ammonium (NH₄⁺) deposition, which can be seen in a two-decade analysis of atmospheric nitrogen deposition at the National Acid Deposition Program network site in Duplin County, North Carolina, a location that has experienced a rapid rise in animal operations during this period (Figure 3).⁶⁶ In Western Europe, where animal operations have dominated agricultural production for the

better part of the past century, ammonium is the most abundant form of atmospherically deposited nitrogen.⁶⁷

Phosphorus is a component of atmospheric deposition, but it typically occurs at concentrations less than a few percent those of nitrogen.⁶⁸ This is especially true in regions where wet exceeds dry deposition, since phosphorus is usually bound to particles such as dust and windblown soils. Accordingly, in agricultural regions where phosphorus is applied as a fertilizer, or in arid regions where soils are readily transported by wind, atmospheric deposition tends to be most highly enriched with phosphorus.⁶⁹ Even in these situations, phosphorus inputs rarely exceed nitrogen inputs.

From an ecological perspective, however, phosphorus may be of considerable

importance since far less phosphorus than nitrogen is required for balanced plant growth.⁷⁰ Therefore, in phosphorus-limited lakes, rivers, reservoirs, and even some marine systems such as the eastern Mediterranean Sea, atmospheric phosphorus inputs can be a significant nutrient source. For example, in Mid-western lakes, including the Great Lakes, atmospheric deposition of phosphorus contributes from 5 to 15 percent of the externally supplied phosphorus.⁷¹ In a recent study of the Mid-Atlantic coastal region, concentrations of total dissolved phosphorus in rainfall ranged from 4 to 15 micrograms per liter at nine sites, and total wet deposition ranged from 3.9 to 14 milligrams per square meter per year across the region.⁷² Annual total phosphorus loading to Lake Michigan in 1976 was 1.7 million kilograms per year, representing about 16 percent of the whole lake's phosphorus budget.⁷³ In alpine Lake Tahoe on the California-Nevada border, atmospherically deposited phosphorus accounts for approximately 25 percent of annual phosphorus inputs, while in the phosphorus-limited eastern Mediterranean Sea, atmospheric deliveries represent about 10 percent of the "new" phosphorus. Overall, it appears that airborne phosphorus typically accounts for 10 to 20 percent of total phosphorus loadings to water bodies from all sources. It remains unknown whether phosphorus transported into aquatic systems by river or air differs in its availability for stimulating plant growth.

Atmospheric deposition of both nitrogen and phosphorus varies with the seasons. For example, during the dry summer

months when plant nutrient demands are highest, phosphorus inputs from surface runoff are minimal. At the same time, dry and windy conditions tend to favor transport of dust. Since phosphorus is often bound to dust particles, it is possible that atmospherically deposited phosphorus assumes a more important role as a source of "new" phosphorus during these crucial growth

periods. Further investigation and quantification are needed of absolute and seasonal atmospheric phosphorus deposition rates in various geographic regions relative to other phosphorus input sources. In the case of nitrogen, emissions from agricultural, urban, and industrial sources are generally highest in summer.

In the case of iron and other metals, atmospheric deposition, mainly in the form

of dust, is a major source of "new" supplies of these nutrients to coastal and open ocean waters.⁷⁴ Iron can be transported over great distances, as demonstrated by the iron-enriched Saharan dust storms that travel thousands of kilometers over the subtropical North Atlantic to "fertilize" iron-deficient and nutrient-poor waters as far away as the Caribbean Sea and the Eastern Seaboard of the United States.⁷⁵ Iron and trace metals are also generated by volcanic emissions and by various continental pollution sources, including power plant, automotive, and industrial emissions.⁷⁶ While there is uncertainty about the chemical forms and behavior of atmospherically deposited iron that enters the ocean, there is little doubt that it represents an important source of "new" iron in an environment that is otherwise free of external iron inputs.⁷⁷

EMISSION, DEPOSITION, AND FATE PROCESSES AND SCALES

The three major atmospheric pathways by which persistent organic pollutants enter water bodies such as the Great Lakes, Chesapeake Bay, other coastal estuaries, and the coastal and open sea are as (1) wet deposition via rain, snow, and fog, (2) dry deposition of particles, and (3) gaseous exchange between the air and water (Figure 4). Many urban industrial centers are located on or near coastal estuaries and the Great Lakes. Emissions of pollutants into the urban atmosphere are reflected in elevated local and regional pollutant concentrations and also in areas of intense localized atmospheric deposition that are over and above




Figure 3 – A 20 year National Acid Deposition Program Network (NADP) nitrogen depositional record for monitoring station NC35 in Duplin Co., NC, showing increases in NH₄⁺ deposition over time. This area has experienced an increase in livestock operations during this period.

the regional signal. For example, the southern basin of Lake Michigan and northern Chesapeake Bay are subject to contamination by air pollutants (PCBs, polyaromatic hydrocarbons (PAHs), mercury, and trace metals) because of their proximity to industrialized and urbanized Chicago and Baltimore, respectively. Concentrations of PCBs and PAHs are significantly elevated in Chicago and coastal Lake Michigan⁷⁸ and in the air over Chesapeake Bay near Baltimore⁷⁹ compared to the regional signal. Higher atmospheric concentrations of pollutants are ultimately reflected in increased precipitation and dry particle inputs of contaminants to the lake or to estuarine waters, as well as enhanced air-water exchange of organic compounds such as PCBs and PAHs.⁸⁰ Of course, the relative importance of these atmospheric pathways to overall water pollution must be evaluated in terms of other inputs, including discharges from wastewater treatment facilities, pollution from upstream river flow, and mobilization of pollutants from sediments.

All three atmospheric pathways deliver pollutants directly to the water surface. This is especially significant for water bodies that have large surface areas compared to the area of the watershed that supplies their runoff. The Great Lakes and coastal seas are two examples. In turn, polluted water bodies may become sources of contaminants to the local and regional atmosphere as gases are lost from the water column to the air. This has been demonstrated for PCBs in the Great Lakes regions of southern Lake Michigan and Green Bay;⁸¹ for PCBs, PAHs, polychlorinated dibenzo-p-dioxins and -dibenzofurans (PCDD/Fs), and nonylphenols in the New York-New Jersey Harbor Estuary;⁸² and for PCBs and PAHs in the Chesapeake Bay.⁸³

In contrast, many aquatic systems have large watershed-to-water area ratios. In these systems, deposits of atmospheric pollutants onto forests, grasslands, crops, paved areas, and other land surfaces in the watershed serve as important sources of runoff contamination to down-stream lakes and estuaries. This is true of most lakes and estuaries in the Mid-Atlantic States, for example.

The relative importance of atmospheric deposition versus other sources of contamination is best demonstrated by chemical mass balances (Figures 5 and 6). Lake Superior, the largest and most pristine of the Great Lakes, is a prime example of an aquatic system in which the atmosphere must play a dominant role in inputs and losses of persistent

organic pollutants. That is because the lake is cold, nutrient poor, has a large surface area that covers most of its watershed, and the urban and industrial density in the area is low. Cold water and a large surface area enhance the lake's sensitivity to atmospheric inputs and air-water exchange through outgassing or volatilization. During the 1980s, for instance, the PCB burden in Lake Superior decreased exponentially at about 20 percent a year, primarily because of outgassing losses to the air.⁸⁴ Although some PCBs bind to organic particles and sink to the lake bottom, this sedimentation process does not provide permanent removal of these contaminants from the water column. Thus, water-air exchange is the dominant loss mechanism. PCBs in the water column today are in approximate equilibrium with atmospheric concentrations.

The two-to-five fold higher concentration of toxaphene (an insecticide banned in the United State since 1990) than PCBs in Lake Superior has been attributed to a lower sedimentation rate and colder water temperatures relative to the other Great Lakes.⁸⁵ Outgassing is an important loss mechanism for toxaphene, just as it is for PCBs, but on a longer time scale. The half-life for PCB decline in Lake Superior waters is 3.5 years compared to 12 years for toxaphene.⁸⁶ Clearance of toxaphene by volatilization would be faster were it not for the higher atmospheric concentrations generated by continued outgassing of toxaphene from agricultural soils in the southern states upwind from Lake Superior.

The pesticide atrazine provides a counter example to PCBs and toxaphene since it is delivered to water bodies mainly by riverine transport of agricultural runoff, and the role of atmospheric delivery is believed to be minimal. Although atrazine has a 30- to 90-day half-life in soils, transport into rivers and lakes significantly extends its half-life. Lake Michigan and other large aquatic systems are most sensitive to tributary inputs of atrazine, but the long-term impacts on the lake environment are controlled by the long residence times in water and the slow rates at which the compound is transformed.⁸⁷ Only about 1 percent of the atrazine applied to crop fields is lost by runoff to rivers and lakes and another 1 percent to aerial transport. Nevertheless, the large quantities of this pesticide that are applied combine with efficient transport, slow transformation rates, and long residence times in water to cause significant accumulation of atrazine in aquatic systems. The atmosphere plays little or no role in the re-




Figure 4 – Schematic showing the pathways, distribution and food web interactions of persistent organic pollutants entering and leaving aquatic systems (modified from D. Muir). Pollutants can be bound to particles or in gaseous phase and can be deposited directly on aquatic ecosystems via both wet and dry deposition. They can also be deposited on terrestrial systems and then enter aquatic systems via snow melt and run-off. Pollutants can also re-enter the atmosphere, where they can be transported and begin the cycle again.




Figure 5 – Mass balance of persistent organic pollutants in Lake Superior. (a) PCBs; (b) Toxaphene.⁸⁸ Dry deposition includes the contaminant deposition associated with atmospheric particle fallout; wet deposition includes the removal of gas phase and particle-bound contaminant by precipitation such as rain and snowfall.

removal of atrazine from large water bodies, and in-lake losses are dominated by degradation and water outflow.

As in the case of PCBs noted above, however, air-water exchange has been shown to dominate contaminant deposition and loss processes in many aquatic systems for a wide range of persistent organic pollutants, including PCBs, PAHs, chlorinated hydrocarbons (HCHs), toxaphene, and PCDDs/Fs.⁸⁹ In aquatic environments, persistent organic pollutants sorb (adhere or bind) to particulate organic matter, and a fraction of this material sinks into deeper waters and sediments. Once organic pollutants are sequestered in the sediments, they are effectively removed from participating in dynamic air-water exchange. In marine waters, this process represents a major sink controlling the surface recycling and impact of persistent organic pollutants. However, the role of sinking particles and other biogeochemical processes, such as algal uptake, on the global dynamics of persistent organic pollutants has so far not been assessed. We now know that algal uptake and air-water exchange behave as coupled processes in aquatic environments.⁹⁰ That is, atmospheric deposition to surface waters supports the concentration of organic contaminants in algal biomass, and the nutrient status of the waters influences how much of the contaminant is available for volatilization to the air.⁹¹ For instance, eutrophic (nutrient enriched) conditions lead to faster algal uptake and removal of contaminants from the water column as algae die and sink to the bottom.

As shown in Figure 4, the processes of air-water and water-algal exchange may promote the introduction of persistent organic pollutants into the aquatic food chain. This is the dominant process for contamination of remote freshwater ecosystems and their food webs, as well as large lakes and the global oceans.⁹² Recent

data even suggest that estuarine food chains dominated by surface runoff of pollutants may also experience food chain contamination from air-water exchange, especially where local and regional emissions create high atmospheric concentrations of pollutants.⁹³ A key consideration is that air-water exchange delivers organic pollutants to the water column in the form of 100 percent bioavailable dissolved gases, whereas contaminants from riverine sources and in stirred and resuspended sediments may not be readily available for use by organisms.

In summary, atmosphere-water interactions are critically important in the cycling and residence times of persistent organic pollutants and the contamination of food webs in lakes, estuaries, coastal waters and the global ocean. In remote aquatic systems or those with large surface areas, atmospheric deposition in general, and air-water exchange specifically, dominates total inputs. Moreover, air-water exchange is the likely mode of contaminant entry into the food chain where

inputs from surface runoff are minimal, and even in some cases where local surface loadings are significant. The nutrient status of water bodies and the cycling of organic material through the food web also play critical roles in determining the fate and impact of persistent organic pollutants in aquatic ecosystems.

ECOLOGICAL RESPONSES

The environmental factors discussed above determine whether aquatic life will be exposed to atmospherically transported chemicals. Several additional factors, however, determine whether these contaminants will harm aquatic organisms or the animals and humans that consume them. Essentially, all chemicals can be toxic to aquatic life if the exposure concentrations are sufficiently high. Conversely, most chemicals also have threshold concentrations below which no appreciable adverse effects on aquatic life are expected. From the perspective of identifying




Figure 6 – Mass balance of Atrazine in Lake Michigan.⁹⁴ Dry deposition includes the contaminant deposition associated with atmospheric particle fallout; wet deposition includes the removal of gas phase and particle-bound contaminant by precipitation such as rain and snowfall.

BOX 2 – BIOACCUMULATION AND BIOMAGNIFICATION

Bioaccumulation refers to an increase in the concentration of a pollutant over time in a biological organism compared to the chemical's concentration in the environment. Compounds accumulate in living things any time they are taken up and stored faster than they are broken down or excreted. *Biomagnification* refers to the increasing accumulation of a pollutant in organisms as it moves up a food chain. Both are important in considering the ecological impacts of atmospheric pollutants. Through bioaccumulation, chemicals that are dilute in the aquatic environment become more concentrated in an organism's tissues. Even when pollutant concentrations in organisms at lower levels of the food chain are low, top predators can be exposed to high concentrations of a pollutant that has been magnified through the consumption of many prey items with low levels of contamination.

ecological risks, then, we are most interested in those chemicals that are not only atmospherically transported, but transported in quantities sufficient to cause ecological risks.

Toxic chemicals that are notoriously associated with atmospheric transport, such as mercury, DDT, and several other pesticides, are generally considered to be bioaccumulative (see Box 2). Chemicals do not have to be highly bioaccumulative to cause toxicity, however. Copper and zinc, for example, are definitely toxic to aquatic life, but they do not accumulate in animal tissues to a large degree compared to chemicals such as mercury or DDT. Nevertheless, chemicals that are highly bioaccumulative are the most likely to present ecological risks because these contaminants can build up to high doses in animal tissues even when concentrations in the water remain relatively low. This is an important factor because a great deal of dispersion and dilution occurs during atmospheric transport between the emission source and the point at which the chemical falls onto land or water. Highly bioaccumulative chemicals reverse this effect by reconcentrating these dilute chemicals in the tissues of exposed animals.

A second common feature of chemicals that tend to cause ecological effects following airborne transport is that they often have relatively high toxicity; that is, they cause toxic effects at comparatively low exposure levels. Combining this with bioaccumulation builds a typical scenario: Low environmental concentrations of atmospherically transported chemicals are transformed into much higher doses in animal tissues via bioaccumulation, and the accumulated dose creates toxic effects because of the high potency of the chemical. By comparison, airborne contaminants with lower bioaccumulation or lower toxicity would be much less likely to cause ecological risk, unless the mass of atmospherically deposited chemicals was much greater.

Several other factors also affect bioaccumulation. Some chemicals biomagnify as they pass up the food chain. Because the chemical concentration increases with each step up the food chain, organisms that are part of long food chains with multiple links can be more susceptible to bioaccumulative chemicals.⁹⁵ For example, a lake trout might feed on relatively large fish, which eat smaller fish, which in turn eat zooplankton, which feed on algae. Thus a lake trout would be expected to acquire greater doses of contaminants than a fish feeding primarily on zooplankton. Another factor is the structure of the food chain in relation to where a chemical is found in the ecosystem. For example, because the atmospheric transport of DDT began many decades ago and has been reduced in recent years, concentrations of DDT in sediments may far exceed those in the water column. Thus, organisms whose food chains are heavily connected to sediments

—systems where fish feed on bottom-dwelling insect larvae, for instance — may experience greater bioaccumulation than organisms in food chains based on algae living in the water column. Also, as mentioned previously, eutrophic systems with high levels and turnover of algal biomass have the capacity to remove greater contaminant loads from the water column and sequester it in deep sediments where it is less available to the rest of the aquatic food chain than oligotrophic systems.

With regard to toxicity, many problem chemicals have such a high degree of biological activity in the body that they can disrupt normal physiology at comparatively low concentrations. In this sense, it should not be surprising that many pesticides are highly toxic because they are specifically designed to interact with biological systems. In other cases, however, high toxicity appears to be an unfortunate coincidence. Such is the case with methyl mercury, the organic form of mercury that is readily absorbed by fish and for which fish and other animals have evolved no specific detoxification mechanism. Chlorinated dioxins/furans and PCBs interact in the body with a specific cell receptor called the aryl-hydrocarbon (Ah) receptor. Depending on their molecular geometry, some congeners bind very tightly to this receptor and thereby cause a high degree of toxicity. While the mechanism by which a chemical causes toxicity is generally similar across a wide range of animal species, the absolute sensitivity — that is, the dose required to cause an adverse effect — can vary substantially across species, often by factors of 100-fold or more. This obviously places some species at greater risk than others.

Considering past and present environmental problems associated with toxic chemicals, it becomes clear how various factors can interact to produce ecological risks. For example, the scenario of risks to lake trout in Lake Ontario from dioxin/furan and PCB exposure combines highly toxic multimedia chemicals with an organism that has a relatively long food chain and a high toxicological sensitivity to Ah-active chemicals.⁹⁶ Loons may be particularly sensitive to mercury because they have a diet high in fish, which indicates a long food chain, and live comparatively long lives. Marine mammals are also long lived, eat aquatic life, and are often used as an indicator of exposure to bioaccumulative chemicals.

Organochlorine Chemicals, a Legacy of the Past

In general, the management response to many organochlorine pollutants — including PCBs, dioxins, furans, and chlorinated pesticides such as DDT, DDE, PCBs, toxaphene, HCHs, dieldrin, mirex, and chlordane — has become an ecological success story in recent decades. Following the recognition in the 1960s and 1970s that widespread use and airborne transport of several

chlorinated compounds was causing reproductive failures, embryonic deformities, and behavioral problems in predatory birds, the chemicals were quickly phased out in North America. While these chemicals are very persistent, long-term records for ecosystems such as the St. Lawrence Great Lakes indicate that concentrations in the environment have decreased with half times of 6 to 10 years.⁹⁷ As a result, environmental concentrations have been reduced enough to eliminate the most severe effects of organochlorines in temperate regions of North America and Europe, accelerating the recovery of populations of peregrine falcons, bald eagles, ospreys, cormorants, gulls, and other affected species (Figure 7).

A few organochlorines continue to be used in agriculture. In particular, gamma-hexachlorocyclohexane (gamma HCH) is used in seed treatment. High concentrations have been measured in the Rocky Mountains of Alberta at times when crops are planted on the nearby prairies.⁹⁸ This compound is also among the most volatile of organochlorines, and is found in relatively high concentrations in the arctic atmosphere.⁹⁹ Hexachlorobenzene (HCB) was once used as a fungicide. Although this use has been banned for more than 20 years, HCB is still released as a byproduct of production of other chlorinated compounds, in the flue gases from municipal waste incineration, and in some metallurgical processes.¹⁰⁰ In glacier samples from the Columbia Icefields of the Canadian Rockies obtained in 1995, concentrations of HCB continued to increase from depth to the very surface, indicating that deposition of HCB has increased even while deposition of other organochlorines has generally decreased in recent years.¹⁰¹

For many organic pollutants that are persistent, bioaccumulative, and toxic, non-atmospheric sources have been well controlled while atmospheric sources have been either neglected or ignored. PCBs are a good example of this phenomenon. PCBs were used in a wide variety of industrial applications and contaminated many industrial and municipal effluent discharges during the period of their use from 1930 to the 1970s. Significant inputs to ecological systems occurred through direct discharge to surface waters or to wastewater treatment plants. Currently, these sources have been diverted, eliminated, or significantly reduced. The entire state of Illinois, for example, discharges no municipal effluent to Lake Michigan, preventing significant input of residual PCBs to the lake. However, atmospheric sources have not received the same attention, probably because few measurements of atmospheric concentrations of PCBs in any urban area were available prior to the mid-1990s. Even now, Chicago is one of the very few urban areas for which substantial atmospheric measurements are available, and concentrations of PCBs are very high in the Chicago area.¹⁰² As a result, deposition of gas-phase

PCBs from Chicago to Lake Michigan is probably the most important current source of these chemicals to the lake. In fact, if the prevailing winds drive the Chicago source away from the lake, Lake Michigan will degas PCBs at a faster rate than it is absorbing them from the air. When the Chicago plume is pushed over the lake by southwest winds, the area of deposition can range from several kilometers to three-quarters of the entire lake (Figure 8). Other urban areas with similar industrial histories are also likely to remain major sources of long-banned organic pollutants, at least until the importance of atmospheric sources is recognized and decisions are made for their control.

Concerns remain about ecological effects of airborne organochlorines at high latitudes and altitudes. The semi-volatile nature of many organochlorines allows them to be re-emitted from contaminated ecosystems during warm weather, and to be carried in air masses and re-deposited where temperatures are cooler. This so-called "cold condensation" effect¹⁰³ has allowed the gradual atmospheric migration of persistent compounds from tropical and temperate regions to arctic and alpine sites.¹⁰⁴ Circumpolar movement in air masses has also allowed some of the chemicals to migrate from areas of Asia where they are still in use to be deposited in arctic and alpine regions of North America and Europe. As a result, deposition of many chemicals in northern and alpine regions continued to increase for 10 to 30 years after the chemicals were banned in North America, as shown from profiles in dated sediments and glaciers.¹⁰⁵ Concentrations in some areas of the arctic are still high enough to cause reproductive failure and eggshell thinning in predatory birds such as peregrine falcons and sea eagles. Liver enzyme induction, which is strongly affected by PCBs and other organochlorines, has been observed in beluga whales, seals, and polar bears.¹⁰⁶

While concentrations of organochlorines in air masses and precipitation from northern and alpine regions are generally low, the food web dynamics, physiologies, and life cycles of cold region organisms allow these chemicals to be biomagnified to extraordinary degrees in food chains. Most organochlorines are lipophilic, and animals in cold regions generally store large amounts of lipids to allow their survival during long winters. Many of the organisms also grow slowly and have long life spans, characteristics that promote biomagnification. As a result, many pollutants are biomagnified by a million-fold and more in arctic food chains.¹⁰⁷ Concentrations are high enough to be of considerable concern for predators of marine mammals, including native people and polar bears.¹⁰⁸ For example, Inuit women in northern Quebec who rely heavily on fish and marine mammals for food have concentrations of many pollutants in their breast milk and notocord blood that are many times higher than in urban Caucasian women



Figure 7 – The reproductive success of bald eagles and other birds of prey was impacted by the use and transport of chlorinated compounds, such as DDT. Since these compounds have been phased out in North America, environmental concentrations have been reduced, accelerating the recovery of these bird populations.

from that province. Concentrations of many of the compounds greatly exceed World Health Organization recommendations, and they also exceed concentrations that have caused detectable reproductive effects in women from contaminated temperate regions.¹⁰⁹ So far, no published studies have documented health effects in northern human populations, but such studies are only in preliminary stages and epidemiologically significant results are difficult to obtain from small populations.

Some northern inland areas where predatory freshwater fish grow slowly and live long can also have concentrations of organochlorines high enough to be of concern. One example is Lake Laberge, Yukon, where lake trout were found to have elevated concentrations of toxaphene, DDT, and PCBs. Toxaphene concentrations were high enough to cause Health Canada to issue a consumption advisory to aboriginal populations who used fish from the lake as an important part of their diets. The main reason for high values in Laberge appears to be that its food chain is effectively one step longer than in other area lakes, thus enhancing biomagnification of these pollutants.¹¹⁰

Unusually high concentrations of these same organo-chlorines were noted in some lake trout populations in the national parks of the Canadian Rocky Mountains, a region generally regarded as near pristine.¹¹² A detailed study of Bow Lake, where some of the highest concentrations were found, indicated that in this case the contamination levels resulted not from biomagnification but largely from increased deposition at high altitude as well as from the melting of glacial strata from the mid twentieth century when high concentrations of the substances were deposited in glaciers before their use was banned in North America.¹¹³ Biomagnification in food chains is low in the lake, and high concentrations of organochlorines in fish appear to result from high inputs and direct absorption of the compounds by small crustaceans on which the fish feed.¹¹⁴ No consumption advisories were issued for Bow Lake because there are no human populations who rely on the lake for subsistence. Similarly, high concentrations of organochlorines have been measured in the sediments and fishes of alpine lakes in Europe, the result of high atmospheric inputs. Unlike in western Canada, however, the source areas for the compounds appear to be regional rather than trans-Pacific.¹¹⁵

Another example of northern concerns involves polar bears (Figure 9). A study in the Svalbard region of Norway found no

reproductive effects in polar bears that were attributable to high PCBs. Yet researchers did observe high mortalities of young bears, and two of the bears had deformed genitalia.¹¹⁶ One hypothesis is that these effects might be the result of high body burdens of organochlorines. For polar bears and many other arctic species, concentrations of organochlorines exceed the No-Observed-Adverse-Effect Level (NOAEL) and Low-Observed-Adverse-Effect Level (LOAEL) that have been determined for southern species.¹¹⁷

There is also concern about additive or synergistic effects of many organochlorines. For example, it is now known that dioxins, furans, and some mono-ortho PCBs all bind to the Ah receptor. Effects appear to be additive, but different compounds have different potencies. In recognition of this, a combined toxic equivalency factor (TEF) scheme has been developed which allows concentrations of many dioxins, furans, and PCBs to be converted to an equivalent concentration of the most potent of these, the dioxin 2,3,7,8-TCDD.¹¹⁸ The TEF scheme is now widely used, whereas

10 years ago, the effects of various organo-chlorines were considered in isolation.

New Chemicals of Concern

While older organo-chlorines appear to have been controlled before they caused extinctions of predatory bird species or severe global reproductive effects, there are concerns about new organic pollutants that are persistent, toxic, and biomagnify in food chains. Atrazine, melalachlor, and endosulfan pesticides continue to be used, although they are reasonably persistent and toxic. Complex chemicals containing other halogens than chlorine (such as bromine and fluorine) have also emerged as potential environmental toxins. For example, recent studies have shown that levels of flame retardants (PBDEs) are increasing rapidly in the Great Lakes, with doubling times on the order of 3.5 years.¹¹⁹ These contaminants are also found in northern and arctic sites.¹²⁰ PBDEs are persistent and biomagnify in aquatic food chains.¹²¹

Reviews of chemicals detected in high-latitude snow indicate that dozens or even hundreds of industrial organic chemicals that are not used in the region are present, indicating long-range atmospheric transport. In most cases, little is known of the biogeochemistry or toxicity of these contaminants.




Figure 8 – The impact of PCB coastal sources is a strong function of meteorology. This series of plots shows the predicted effect of wind direction (indicated by arrows) and temperature (not shown) on the spatial distribution of PCB net deposition to the surface of Lake Michigan. The gray regions are regions that are degassing PCBs from the water. The black regions are regions that are absorbing PCBs from the air¹¹¹.

Mercury

Once removed from the atmosphere, much deposited mercury ends up in aquatic systems, either by direct deposit or in surface runoff.¹²² The effects of mercury on fish, as well as birds and mammals that eat contaminated fish, are of significant concern, and those concerns stem from exposure to the methylated (organic) form, monomethyl mercury (MMHg).¹²³ In anaerobic environments such as lake or wetland sediments, mercury is transformed to MMHg by microbial action, most notably by sulfur-reducing bacteria. The MMHg diffuses into the water column where it can be taken up by fish, accumulating in their muscle tissue by binding to organic compounds known as thiol groups or mercaptans. MMHg is the most toxic form of mercury. Reproductive effects have been documented in fish, fish-eating wildlife, and humans.¹²⁴ MMHg causes neurological, liver, and kidney damage, as well as neurodevelopmental effects in children.¹²⁵

In the United States, 48 states advise the public against unlimited consumption of freshwater fish due to their MMHg levels. In addition, the U.S. Environmental Protection Agency (EPA) has issued a mercury-based national fish consumption advisory for five species of oceanic fish. The EPA considers the maximum allowable no-effects dose of mercury to be 0.1 micrograms per kilogram of body weight per day, and this guideline was recently supported in an independent study by the National Academy of Sciences.¹²⁶

Arctic ecosystems and their associated human communities are particularly susceptible to mercury contamination. When mercury reaches the Arctic, it is transformed such that it is deposited on the snow at the start of the arctic sunrise (the first appearance of the sun after the long arctic winter). A significant amount of this mercury enters the ecosystem and the rest re-enters the atmosphere where it can be again transported and deposited in other locations.

Nutrients

Nitrogen over-enrichment has been blamed for a wide array of impacts on aquatic ecosystems, including changes in the function and composition of the algal community, changes in the food web, and declines in water quality and fisheries habitat. Increases in nitrogen and changes in nitrogen sources can influence competitive interactions and succession among algal groups, as well as dominance by certain undesirable groups such as red tide dinoflagellates and toxic cyanobacteria (formerly blue-green algae).¹²⁷ The ability of some algal groups to utilize organic




Figure 9 – The food web dynamics, physiologies, and long lives of arctic animals such as polar bears allow contaminants to be biomagnified to extraordinary degrees in food chains, resulting in deformities and, potentially, death.

nitrogen, which can be plentiful in atmospheric deposition, can also provide a competitive advantage.¹²⁸ Iron, which is largely atmospherically derived, may interact synergistically with nitrogen to enhance coastal and oceanic primary production.¹²⁹ These effects of atmospherically transported nitrogen and other nutrient sources may be promoting the major biological changes that are now apparent in coastal and oceanic waters, including the proliferation of harmful algal blooms and declines in water quality and fisheries. Because of its potentially large contribution to total “new” nitrogen loading to nitrogen-sensitive waters, atmospheric

deposition requires attention from those responsible for local and regional air and watershed nutrient management.

In estuarine and coastal marine settings, sediments represent relatively rich sources of phosphorus that are readily cycled between the sediments and the water column. In these waters, atmospherically deposited phosphorus has not been shown to be widely significant, although it may be significant in specific locales. An example is in Florida coastal waters where local airborne phosphorus inputs may be high at times. A few researchers have suggested that primary production in some oceanic regions is limited by low phosphorus levels;¹³⁰ however, this is a controversial notion and the overwhelming evidence continues to point to nitrogen and/or iron as the limiting nutrients in these waters.¹³¹ Although only a few measurements have been made, oceanic regions tend to exhibit extremely low levels of atmospherically deposited phosphorus.¹³² This is not surprising since the open oceans are far removed from continental sources of dust and windblown soils. Volcanic eruptions and large-scale dust storms may be a source of “new” airborne phosphorus inputs in some parts of the world, but the biogeochemical and ecological importance of these inputs remains unknown.

Iron deposited from the atmosphere appears to play a critical role in sustaining and stimulating productivity in iron-deprived coastal and oceanic waters.¹³³ Atmospheric sources of iron have been proposed as key stimulants of marine algal and red tide blooms.¹³⁴

The ecological roles and effects of airborne deposits of trace elements — copper, zinc, manganese, cobalt, molybdenum, and boron — are far less clear.¹³⁵ The impacts could include stimulation or inhibition of primary and secondary production or synergistic or antagonistic interactions with other nutrient effects. Far more mechanistic research (using response bioassays with natural microbial and plant communities as well as test species) is needed to help elucidate basic ecological and biogeochemical roles of

these trace nutrients. It is also conceivable that complex co-limitations exist between trace elements and the major nutrients such as nitrogen, phosphorus, and iron in freshwater and marine environments.¹³⁶ And it is possible that atmospheric deposition provides unique nutrient combinations that together have a more potent effect than individual nutrient constituents.

NITROGEN DEPOSITION AND THE FATE OF ORGANIC POLLUTANTS

Atmospheric deposition contains a mixture of biologically available nitrogen compounds, both organic and inorganic. Bioassay studies in nitrogen-limited waters have demonstrated that organisms respond differently to various nitrogen sources. This provides a mechanism whereby nitrogen deposition can influence the structure and makeup of aquatic communities.¹³⁷ Differential uptake and growth in response to ammonium versus nitrate have been attributed to contrasts in energy required to assimilate these compounds.¹³⁸ Under light-limited conditions encountered in turbid waters, organisms may prefer ammonium because the energy requirements for using this reduced nitrogen source are less than those for using nitrate.¹³⁹ In light-limited waters, motile algal groups such as dinoflagellates and cryptomonads are capable of migrating to near-surface depths to ensure access to the light energy needed to reduce nitrate to ammonium, which is a critical step for incorporating this oxidized form of nitrogen into biosynthetic pathways and growth. In contrast, non-motile organisms must cope with deeper, lower-irradiance waters, possibly limiting their nitrate uptake and thus ammonium uptake. Intrinsic physiological differences in nitrogen uptake among different algal taxa also exist, and these may lead to contrasting responses to different nitrogen sources.¹⁴⁰ Under conditions of restricted nitrogen availability, which are characteristic of many estuaries, such differences can lead to intense competition for ammonium, nitrate, or organic nitrogen.

Bioassay experiments on the Neuse River Estuary in North Carolina have shown that major algal taxonomic groups — diatoms, dinoflagellates, cryptomonads, cyanobacteria, and chlorophytes — may exhibit different growth responses to varying nitrogen sources and mixtures of sources.¹⁴¹ However, these differential responses are not consistent in time and space. Other complex environmental factors, including light availability, water column mixing depth, water residence time, salinity, and temperature, also exert control over the dynamics of nitrogen uptake and growth rates among the algal community.¹⁴² Atmospherically derived dissolved organic nitrogen has also been shown to stimulate bacterial and algal growth.¹⁴³ This organic nitrogen may selectively stimulate growth of facultative heterotrophic algae such as dinoflagellates and cyanobacteria.¹⁴⁴ (Facultative heterotrophic organisms can make their own food or derive their nutrition by consuming organic molecules.)

Responses to specific nitrogen inputs that alter the algal community may in turn spur changes all the way up the food chain, at the zooplankton, herbivorous fish, invertebrate, and higher consumer levels. Shifts in algal community composition may also alter the flux of carbon, nitrogen, phosphorus, and other nutrients and impact oxygen dynamics in the estuary.

The fact that different sources of atmospherically derived nitrogen may influence the structure and growth dynamics of the algal community is also important to the fate of persistent organic pollutants in aquatic food webs. Algae, with their high lipid content, readily bioaccumulate these pollutants, but the rate of uptake of these compounds is heavily influenced by the growth rate of the community.¹⁴⁵ The more productive the system, the less bioaccumulation occurs due to the difference in the rate of uptake compared to the rate of growth. In addition, the more productive the system, the more the fate of these contaminants is dominated by sedimentation of senescent and dead algae. In oligotrophic systems, algal biomass is usually low, but uptake of nutrients and other contaminants is quite efficient. In these systems, effective grazing by zooplankton ensures that these contaminants are readily passed to higher levels of the food web.

A second reason that the source of atmospherically derived nitrogen can influence the fate of persistent organic pollutants is related to the changes in community structure that can occur. It has recently been shown that the microbial loop may be an important vector in organic pollutant transfer in food webs, and its importance increases in systems that are less productive.¹⁴⁶ The microbial loop is an aquatic micro-food chain in which microscopic bacteria and pico-plankton feed on dissolved organic material, then these organisms are grazed on by flagellates and ciliates, and these are consumed by tiny crustaceans called copepods. Because the microbial loop consists of ciliates preying on flagellates preying on bacteria, and the flagellate/ciliate component can be influenced by atmospherically derived nitrogen, this new nitrogen source may indirectly influence the food web dynamics controlling pollutant bioaccumulation.

PRIORITIES FOR REGULATON AND MONITORING

Above we have identified a number of challenges that atmospheric pollutants present to aquatic ecosystems and to the human communities that rely on them. In this section, we suggest some priorities for regulation and monitoring of atmospherically transported chemicals. Effective regulations must be based on a solid understanding of the problem developed through research and monitoring. Actions should be focused on chemicals that are toxic, bioaccumulate or biomagnify, move in measurable quantities into and via the atmosphere, are sufficiently stable to be transported long distances, and interact with other chemicals to have negative impacts on organisms and ecosystems. The actions suggested below allow researchers and managers to identify with some certainty how atmospheric deposition affects aquatic ecosystems and to pinpoint likely sources and consequences.

- **Develop a comprehensive strategy for reducing impacts and problems caused by airborne organic contaminants.**
 - Use quantitative structure activity relationships (QSAR) to predict whether new chemicals that are proposed for commercial use will likely be persistent, bioaccumulative, and/or toxic to organisms, and use this information in decision-making and regulation.

- o Expand testing of HPVCs and carefully scrutinize synthetic fragrances and flame retardants as potential atmospheric pollutants.
- o Establish monitoring programs for mercury and persistent bioaccumulative, toxic organic compounds in critically vulnerable ecosystems, such as alpine and arctic systems.
- o Include the interactions of nutrient and toxic chemical deposition in models and regulatory considerations.
- **Develop a comprehensive strategy for reducing the nitrogen problem.**
 - o Develop more effective and broadly applied controls on nitrogen oxide emissions from internal combustion engines in automobiles, boats (i.e. outboard motors), personal watercraft and all terrain vehicles, lawnmowers, chain saws, and other fossil fuel-powered tools and machines.
 - o Develop more effective and broadly applied nitrogen oxide controls on industrial and power plant emissions (i.e. stack emission controls).
 - o Minimize open-air storage of animal wastes and other reduced nitrogen products and sources and improve treatment of animal wastes using on-site “treatment plants” and engineered wetlands. Use recycled water in animal operations to minimize the generation and storage of liquid animal waste.
 - o Recycle accumulated solid waste into commercial fertilizers. Apply nitrate, ammonium and urea-based fertilizers at agronomic rates.
 - o Use “controlled” burns to minimize atmospheric “fertilization” of downwind nitrogen-sensitive waters with either nitrogen oxides or ammonia/ammonium.
- **Develop less expensive and more accurate methods to measure atmospheric deposition and to monitor and model how atmospherically deposited pollutants travel through a watershed.** This should include establishing monitoring programs in critically vulnerable ecosystems, such as monitoring for organic pollutants in alpine and arctic ecosystems and for nutrients in estuaries.
- **Strengthen understanding of the linkages between atmospheric deposition and ecological effects, particularly for new chemicals of interest and interactions of nutrients and toxic compounds.** Exposure risk must be analyzed for sensitive habitats and sensitive life stages as well as general populations.
- **Develop quantitative estimates of atmospheric pollutant deposition loads through monitoring and measurements.** This should be done in conjunction with identifying sources of atmospheric contaminants and how much comes from local, regional, and long-range sources. Quantitative measurements will also help in identifying the role that atmospheric deposition plays in a watershed’s overall pollutant load.
- **Build proactive cooperation between air pollution agencies, water pollution agencies, and the public.** Agencies monitoring and regulating air quality and those monitoring and regulating water quality will need to cooperate to address this issue. Now that the public is beginning to embrace the

concept of watersheds, it is important to increase their understanding of the links between airsheds and watersheds.

CONCLUSION

In recent decades, much progress has been made in reducing the input of toxic chemicals and nutrients into the environment from point sources. However, atmospheric sources of toxic substances and nutrients are just now starting to be recognized for the role they play in contaminating the environment.

Organic chemicals that volatilize easily can be transported and deposited in other regions of the world, exposing aquatic ecosystems to chemicals not used in those regions. In some cases, little is known of their biogeochemistry or toxicity. Often, however, these chemicals have properties that make them environmentally mobile, persistent in the ecosystem, and bioaccumulative in living tissue. Many are toxic to aquatic organisms, fish, and fish-eating wildlife and humans.

Mercury is also a global pollutant that is mobile in the atmosphere, and its effects on fish, and on the birds and mammals that eat contaminated fish, are of significant concern. In some aquatic systems, mercury is transformed by microbial action into its methylated form, which can cause neurological, liver, and kidney damage as well as reproductive effects and neurodevelopmental problems.

Atmospheric deposition is also a significant and potent source of nutrients that can accelerate eutrophication and its associated environmental consequences in freshwater, estuarine, and coastal ecosystems. Aquatic ecosystems are often impacted by atmospheric deposition of both nutrients and toxic chemicals. The effects of nutrient deposition on food web structure and ecological function influence how other toxic substances are processed by the ecosystem, how they bioaccumulate, and ultimately how they impact fish, wildlife, and humans.

Considering that these atmospherically deposited contaminants are generated largely by human activities, it is clear that solutions must involve greater recognition, monitoring, and ultimately, regulation of this increasingly significant source of environmental pollution.

ACKNOWLEDGMENTS

We greatly appreciate the U.S. Environmental Protection Agency’s Office of Wetlands, Oceans and Watersheds for its support of the production of this report. We are particularly grateful to Deborah Martin and John Wilson. We would also like to thank David Whittall, NOAA, and Jules Blais, University of Ottawa, for their comments on a draft of this manuscript.

REFERENCES AND SUGGESTED READINGS

- ¹ USEPA (U.S. Environmental Protection Agency). 2001. Frequently Asked Questions About Atmospheric Deposition: A Handbook for Watershed Managers. EPA 453/R-01-009. http://www.epa.gov/owow/oceans/airdep/airdep_sept_final.pdf.
- ² Airsheds developed by R. Dennis, Atmospheric Science Modeling Division, NOAA Air Resources Laboratory and USEPA National

- Exposure Research Laboratory. Presented in USEPA 2001 (see 1); courtesy of the U.S. Environmental Protection Agency.
- 3 Van de Meent, T.E. McKone, T. Parkerton, M. Matthies, M. Scheringer, F. Wania, R. Perudy, D.H. Bennett. 2000. Persistence and transport potential of chemicals in a multimedia environment. In: Klečka, G. et al. (Eds). Evaluation of Persistence and Long Range Transport of Organic Chemicals in the Environment: SETAC Special Publ. Series, SETAC Books, Pensacola FL, pp. 169-204.
 - 4 Wania, F. and D. Mackay 2000. The global distribution model. A non-steady state multi-compartmental mass balance model of the fate of persistent organic pollutants in the global environment. Computer Program and Technical Report, 21 pages. (www.utscc.utoronto.ca/~wania).
 - 5 Figure reprinted with permission from Van de Meent, T.E. McKone, T. Parkerton, M. Matthies, M. Scheringer, F. Wania, R. Perudy, D.H. Bennett. 2000. Persistence and transport potential of chemicals in a multimedia environment. In: Klečka, G. et al. (Eds). Evaluation of Persistence and Long Range Transport of Organic Chemicals in the Environment: SETAC Special Publ. Series, SETAC Books, Pensacola FL, pp. 169-204. Copyright SETAC, Pensacola, FL
 - 6 UNEP (United Nations Environment Program), 2001. Final Act of the Conference of Plenipotentiaries on The Stockholm Convention On Persistent Organic Pollutants. UNEP, Geneva, Switzerland. 44 pp.
Detailed discussions of physical-chemical properties of the chemicals listed in the Stockholm Convention can be found in the Handbooks on Physical-Chemical Properties and Environmental Fate of Organic Chemicals by Mackay et al. (1991, 1992, 1997) and the Evaluation of Persistence and Long Range Transport of Organic Chemicals in the Environment resulting from a workshop of the Society of Environmental Toxicology and Chemistry (SETAC; Klečka, G., R.S. Boethling, J. Franklin, C.P.L. Grady Jr., D.G. Graham, P.H. Howard, K. Kannan, R.J. Larson, D. Mackay, D. Muir and D. van de Meent, 2000. Evaluation of persistence and long range transport of organic chemicals in the environment. SETAC Special Publications Series, SETAC Books, Pensacola FL, USA. 362 pp.)
 - 7 Solomon, K.R., Baker, D. B., Richards, R. P., Dixon, K.R., Klaine, S. J., La Point, T. W., Kendall, R. J., Weisskopf, C. P., Giddings, J.M., Giesy, J.P., Hall, L. W., Williams, W. M. 1996. Ecological risk assessment of atrazine in north american surface waters. *Environ. Toxicol. Chem.* 15: 31-76.
 - 8 Metcalfe, C.D., X.-S. Miao, B.G. Koenig and J. Struger. 2003. Distribution of acidic and neutral drugs in surface waters near sewage treatment plants in the lower Great Lakes, Canada. *Environ Toxicol Chem* 22 (12):2881-2889.
Keith, T.L., S.A. Snyder, C.G. Naylor, C.A. Staples, C. Summer, K. Kannan and J.P. Geisy. 2001. Identification and quantitation of nonylphenol ethoxylates and nonylphenol in fish tissues from Michigan. *Environ Sci Technol* 36: 10-13.
White R., Jobling, S., Hoare, S.A., Sumpter, J.P., and Parker, M.G. 1994. Environmentally persistent alkylphenolic compounds are estrogenic. *Endocrinology* 135:175-182.
 - 9 Dachs, J., T.R. Glenn, C.L. Gigliotti, P. Brunciak, E.D. Nelson, T.P. Franz, and S.J. Eisenreich. 2002. Occurrence and diurnal variability of PAHs in the Baltimore and adjacent Chesapeake Bay atmosphere. *Atmos Environ* 36(14):2281-2295.
 - 10 Sijm, D., J. de Bruijn, P. de Voogt and W. de Wolf (Eds.) 1997. Biotransformation in Environmental Risk Assessment. Society of Environmental Toxicology and Chemistry. (SETAC) - Europe, Brussels, Belgium.
 - 11 USEPA (US Environmental Protection Agency). 1998. Chemical Hazard Data Availability Study. High Production Volume (HPV) Chemicals and SIDS Testing. <http://www.epa.gov/opptintr/chemtest/hazchem.htm>.
 - 12 ECB (European Chemicals Bureau). 2002. Existing Chemicals. <http://ecb.jrc.it/existing-chemicals/>
 - 13 OECD (Organization of Economic Cooperation and Development). 2000. OECD integrated HPV database. <http://cs3-hq.oecd.org/scripts/hpv/>
 - 14 ICCA (International Council of Chemical Associations). 2003. <http://www.cefic.org/activities/hse/mgt/hpv/hpvinit.htm>
 - 15 Rimkus, G. G. 1999. Polycyclic musk fragrances in the aquatic environment. *Toxicology Letters* 111:37-56.
Rimkus, G. G., R. Gatermann, and H. Huhnerfuss. 1999. Musk xylene and musk ketone amino metabolites in the aquatic environment. *Toxicology Letters* 111:5-15.
 - 16 Editorial. 1999. The Toxicology of Musk Fragrances. *Toxicology Letters* 111:1-4.
Franke, S., C. Meyer, N. Heinzel, R. Gatermann, H. Huhnerfuss, G. Rimkus, W. A. König, and W. Francke. 1999. Enantiomeric composition of the polycyclic musks HCHB and AHTN in different aquatic species. *Chirality* 11:795-801.
Rimkus, G. G. 1999. Polycyclic musk fragrances in the aquatic environment. *Toxicology Letters* 111:37-56.
 - 17 Peck, A. M., and K. C. Hornbuckle. 2004. Synthetic musk fragrances in Lake Michigan. *Environ Sci Technol* 38:367-372.
 - 18 Schreurs, R. H. M. M., J. Legler, E. Artola-Garicano, T. L. Sinnige, P. H. Lanser, W. Seinen, and B. van der Burg. 2004. In vitro and in vivo antiestrogenic effects of polycyclic musks in zebrafish. *Environ Sci Technol* 38:997-1002.
 - 19 SCCNFP (Scientific Committee on Cosmetic Products and Non-Food Products). 2002. On the basis of the International Fragrance Association (IFRA) Code of Practice, the SCCNFP has listed 36 chemical substances, including musk ketone and musk xylene that must not form part of fragrance compounds used in cosmetic products. *European Community Legislation Official Journal L* 102 45:19 (18 Apr 2002).
 - 20 Walker, J.D. and L. Carlsen. 2002. QSARs for identifying and prioritizing substances with persistence and bioconcentration potential. *SAR QSAR Environ Res* 13: 7131-725.
 - 21 World Health Organization. 1994. Environmental Health Criteria 162: Brominated Diphenyl Ethers. Geneva.
Renner, R. 2000. *Environ Sci Technol* 34:452A-453A.
 - 22 Ikonomou, M.G., S. Rayne and R.F. Addison. 2002. Exponential increases of the brominated flame retardants, polybrominated diphenyl ethers, in the Canadian Arctic from 1981-2000. *Environ Sci Technol* 36:1886-1892.
 - 23 Watanabe, I. and R. Tatsukawa. 1987. Formation of brominated dibenzofurans from the photolysis of flame retardant decabromobiphenyl ether in hexane solution by UV and sunlight. *Bull Environ Contam Toxicol* 39:953-959.
Tysklind, M., U. Sellström, G. Söderström and C. de Wit, 2001. Abiotic transformation of polybrominated diphenylethers (PBDEs): Photolytic debromination of decabromo diphenyl ether. Extended abstract, pp 51-54, Second International Workshop on Brominated Flame Retardants BFR 2001, 14-16 May, Stockholm, Sweden.
Stapleton et al. 2003. Biotransformation of decabromodiphenyl ether in fish. *Environ Sci Technol* Submitted.
 - 24 Ikonomou et al. 2002 (see 22).

- ²⁵ Giesy, J.P. and K. Kannan, 2001. Global distribution of perfluorooctane sulfonate in wildlife. *Environ Sci Technol* 35: 1339-1342.
- Kannan K., J.C. Franson, W.W. Bowerman, K.J. Hansen, P.D. Jones, J.P. and Giesy. 2001a. Perfluorooctane sulfonate in fish-eating birds including bald eagles and albatrosses. *Environ Sci Technol* 35: 3065-3070.
- Kannan K., J. Koistinen, K. Beckman, T. Evans, J.F. Gorzelany, et al. 2001b. Accumulation of perfluorooctane sulfonate in marine mammals. *Environ Sci Technol* 35: 1593-1598.
- Kannan K., J.-W. Choi, N. Iseki, K. Senthilkumar, D.H. Kim, et al. 2002a. Concentrations of perfluorinated acids in livers of birds from Japan and Korea. *Chemosphere* 49: 225-231.
- Kannan K., Corsolini S, Falandysz J, Oehme G, Focardi S, Giesy JP. 2002b. Perfluorooctanesulfonate and related fluorinated hydrocarbons in marine mammals, fishes, and birds from coasts of the Baltic and the Mediterranean seas. *Environ Sci Technol* 36: 3210-3216.
- Kannan K., K.J. Hansen, T.L. Wade, and J.P. Giesy. 2002c. Perfluorooctane sulfonate in oysters, *Crassostrea virginica*, from the Gulf of Mexico and the Chesapeake Bay, USA. *Arch Environ Contam Toxicol* 42: 313-318.
- Kannan K., J. Newsted, R.S. Halbrook, and J.P. Giesy. 2002d. Perfluorooctanesulfonate and related fluorinated hydrocarbons in mink and river otters from the United States. *Environ Sci Technol* 36: 2566-2571.
- Hansen K.J., L.A. Clemen, M.E. Ellefson, and H.O. Johnson. 2001. Compound specific quantitative characterization of organic fluorochemicals in biological matrices. *Environ. Sci. Technol.* 35: 766-770.
- ²⁶ Renner, R. 2001. Growing concern over perfluorinated chemicals. *Environmental Science and Technology*, 35: 154A-160A.
- ²⁷ Stock, N.L., F.K. Lau, D.A. Ellis, J.W. Martin, D.C.G. Muir and S.A. Mabury. 2003. Polyfluorinated telomer alcohols and sulfonamides in the North American troposphere. *Environ Sci Technol* Submitted.
- ²⁸ Giesy and Kannan 2001 (see 25).
Kannan et al. 2001b (see 25).
- ²⁹ Smithwick, M.M., J.W. Martin, C. Sonne-Hansen, K.R. Solomon, S.A. Mabury, and D.C.G. Muir. 2003. Perfluorinated organics in polar bears (*Ursus maritimus*). Canadian Arctic and Greenland. Annual Meeting of Society of Environmental Toxicology and Chemistry – Europe, Hamburg Germany, May 2003.
- ³⁰ Fogg, T.R. and W.F. Fitzgerald. 1979. Mercury in southern New England rains. *J. Geophys. Res. – Oceans and Atmospheres* 84:6987-6988.
- Mason, R.P., W.F. Fitzgerald and F.M.M. Morel. 1994. The biogeochemical cycling of elemental mercury: anthropogenic influences. *Geochimica Et Cosmochimica Acta* 58: 3191-3198.
- ³¹ Keeler, G., G. Glinsorn, and N. Pirrone. 1995. Particulate mercury in the atmosphere: Its significance, transport, transformation and sources. *Water Air and Soil Pollution* 80:159-168.
- ³² Fitzgerald, W.F. and C.J. Watras. 1989. Mercury in surficial waters of rural Wisconsin lakes: *Total Environment* 87:8:223-232.
- Lindqvist, O., K. Johansson, M. Aastrup, A. Andersson, L. Bringmark, G. Hovsenius, L. Hakanson, A. Iverfeldt, M. Meili, and B. Timm. 1991. Mercury in the Swedish Environment - Recent Research on Causes, Consequences and Corrective Methods. *Water Air and Soil Pollution* 55.
- ³³ Engstrom, D. R., and E. B. Swain. 1997. Recent declines in atmospheric mercury deposition in the upper Midwest. *Environ Sci Technol* 312:960-967.
- ³⁴ Fitzgerald, W.F. 1995. Is mercury increasing in the atmosphere? The need for an Atmospheric Mercury Network (AMNET). *Water, Air and Soil Pollution* 80: 245-254.
- ³⁵ Stratton, W. J., and S. E. Lindberg. 1995a. Measurement of gas-phase ionic mercury(ii) species in ambient air. *Abstracts of Papers of the American Chemical Society* 210:60-Geoc.
- Stratton, W. J., and S. E. Lindberg. 1995b. Use of a refluxing mist chamber for measurement of gas-phase mercury(ii) species in the atmosphere. *Water Air and Soil Pollution* 80:1269-1278.
- ³⁶ Mason, R.P. and W.F. Fitzgerald. 1996. Sources, sinks and biochemical cycling of mercury in the ocean. In: Baeyens, W., Ebinghaus, R. and Valiliev, O. (eds.): *Global and regional mercury cycles: Sources, fluxes and mass balances*. NATO ASI Series, 2. Environment - Vol. 21. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- ³⁷ Rada, R.G., Winfrey, M.R., Wiener, J.G., and Powell, D.F. 1987. A comparison of mercury distribution in sediment cores and mercury volatilization from surface waters of selected northern Wisconsin lakes. Wisconsin Department of Natural Resources, Madison, Wis.
- Engstrom, D.R., E.B. Swain, T.A. Henning, M.E. Brigham and P.L. Brezonik. 1994. Atmospheric mercury deposition to lakes and watersheds: A quantitative reconstruction from multiple sediment cores. In Baker, L.A., (Ed.) *Environmental Chemistry of Lakes and Reservoirs*. Washington, D.C., American Chemical Society. pp. 33-66.
- Benoit G., E.X. Wang, W.C. Nieder, M. Levandowsky, and V. Breslin. 1999. Sources and history of heavy metal contamination and sediment deposition in Tivoli South Bay, Hudson River, NY. *Estuaries* 22:221-244.
- ³⁸ Mason, R.P. and G.-R. Sheu. 2002. Role of the ocean in the global mercury cycle. *Global Biogeochemical Cycles* 16(4) 1093.
- ³⁹ Ryther, J. and W. Dunstan. 1971. Nitrogen, phosphorus and eutrophication in the coastal marine environment. *Science* 171:1008-1112.
- Likens, G.E. [Ed.]. 1972. *Nutrients and Eutrophication*. American Society of Limnology and Oceanography Special Symposium 1.
- Likens, G., F. Borman, and M. Johnson. 1974. Acid rain. *Environment* 14:33-40.
- Paerl, H. W. 1985. Enhancement of marine primary production by nitrogen enriched acid rain. *Nature* 316:747-749.
- Duce, R. 1986. The impact of atmospheric nitrogen, phosphorus, and iron species on marine biological productivity, pp. 497-529. In P. Buat-Menard (Ed.), *The Role of Air-Sea Exchange in Geochemical Cycling*, D. Reidel, Norwell, MA.
- Paerl, H.W, R.L. Dennis and D.R. Whittall. 2002. Atmospheric deposition of nitrogen: Implications for nutrient over-enrichment of coastal waters. *Estuaries* 25:677-693.
- ⁴⁰ Martin, J.H., K. H. Coale, K. S. Johnson, S. E. Fitzwater, et al. 1994. Testing the iron hypothesis in ecosystems of the equatorial Pacific Ocean. *Nature* 371:123-129.
- Takeda, S., A. Kamatani, and K. Kawanobe. 1995. Effects of nitrogen and iron enrichments on phytoplankton communities in the Northwestern Indian Ocean. *Mar Chem* 50:229-241.
- ⁴¹ Takeda et al. 1995 (see 40).
- Paerl, H.W., J.D. Willey, M. Go, B.L. Peierls, J.L. Pinckney and M.L. Fogel. 1999. Rainfall stimulation of primary production in

- Western Atlantic Ocean waters: Roles of different nitrogen sources and co-limiting nutrients. *Mar Ecol Progr Ser* 176:205-214.
- ⁴² Likens et al. 1974 (see 39).
- Bribblecombe, P. and D.H. Stedman. 1982. Historical evidence for dramatic increase in the nitrate component of acid rain. *Nature* 298:460-462.
- ⁴³ Howarth, R.W., G. Billen, D. Swaney, A. Townsend, N. Jaworski, K. Lajtha, J.A. Downing, R. Elmgren, N. Caraco, T. Jordan, F. Berendse, J. Freney, V. Kudeyarov, P. Murdoch and Z. Zhao-Liang. 1996. Regional nitrogen budgets and riverine N and P fluxes for the drainages to the North Atlantic Ocean: Natural and human influences. *Biogeochemistry* 35:75-139.
- ⁴⁴ Paerl, H.W. 1995. Coastal eutrophication in relation to atmospheric nitrogen deposition: current perspectives. *Ophelia* 41:237-259.
- Jaworski, N., R. Howarth and L. Hetling. 1997. Atmospheric deposition of nitrogen oxides onto the landscape contributes to coastal eutrophication in the Northeast United States. *Environ Sci Technol* 31:1995-2004.
- Castro, M.S., C.T. Driscoll, T.E. Jordan, W.R. Reay, and W.R. Boynton, S.P. Seitzinger, R.V. Styles, and J.E. Cable. 2000. Contribution of atmospheric deposition to the total nitrogen loads of thirty-four estuaries on the Atlantic and Gulf Coast of the United States. Pp., 77-106, In: R. Valigura (ed.), *Atmospheric Nitrogen Deposition in Coastal Waters*. No. 57, Coastal Estuarine Science Series. American Geophysical Union Press, Washington, DC.
- Valigura, R.A., R.B. Alexander, M.S. Castro, T.P. Meyers, H.W. Paerl, P.E. Stacey, and R.E. Turner (eds.). 2000. Nitrogen Loading in Coastal Water Bodies: An Atmospheric Perspective. *Coastal and Estuarine Studies* No. 57. American Geophysical Union, Washington, D.C. 254 p.
- Elmgren, R. and U. Larsson. 2001. Eutrophication of the Baltic Sea area. *Integrated Coastal Management Issues*. pp. 15-35, In: B. von Bodungen, and R. K. Turner (eds.), *Science and Integrated Coastal Management*. Dahlem Press, Berlin.
- ⁴⁵ Paerl, H.W. and M.L. Fogel. 1994. Isotopic characterization of atmospheric nitrogen inputs as sources of enhanced primary production in coastal Atlantic Ocean waters. *Mar Biol* 119:635-645.
- Paerl, H.W. and D.R. Whitall. 1999. Anthropogenically-derived atmospheric nitrogen deposition, marine eutrophication and harmful algal bloom expansion: Is there a link? *Ambio* 28:307-311.
- ⁴⁶ Howarth et al. 1996 (see 43).
- ⁴⁷ Paerl et al. 2002 (see 39).
- ⁴⁸ Table adapted from Paerl et al. 2002 (see 39).
- ⁴⁹ Elmgren R. 1989. Man's impact on the ecosystem of the Baltic Sea: Energy flows today and at the turn of the century. *Ambio* 18:326-332.
- Ambio*. 1990. Marine eutrophication. *Ambio* 19:101-176.
- ⁵⁰ Prado-Fielder, R.R. 1990. Atmospheric input of inorganic nitrogen species to the Kiel Bight. *Helgoland Meeresuntersuchungen* 44:21-30.
- ⁵¹ GESAMP. 1989. The Atmospheric Input of Trace Species to the World Ocean: Report and Studies No. 38. World Meteorological Association, Geneva.
- ⁵² Martin, J.M., F. Elbaz-Poulichet, C. Gwue, M.D. Loye-Pilot and G. Han. 1989. River versus atmospheric input of material to the Mediterranean Sea: An overview. *Mar Chem* 28:159-182.
- ⁵³ Valiela, I., K. Foreman, M. LaMontagne, D. Hersch, J. Costa, P. Peckol, B. DeMeo-Anderson, C. D'Avanzo, M. Babione, C.-H. Sham, J. Brawley and K. Lajtha. 1992. Couplings of watersheds and coastal waters: Sources and consequences of nutrient enrichment in Waquoit Bay, Massachusetts. *Estuaries* 15:443-457.
- ⁵⁴ Nixon, S. 1995. Coastal marine eutrophication: A definition, social causes, and future concerns. *Ophelia* 41:199-220.
- ⁵⁵ Long Island Sound Study. 1996. Report on nitrogen and organic carbon loads to Long Island Sound. Prepared by P. Stacey, Connecticut Department of Environmental Protection, Bureaus of Water Management, Hartford, CT.
- ⁵⁶ Valigura, R.A., W. Luke, R. Artz, and B. Hicks. 1996. Atmospheric Nutrient Inputs to Coastal Areas: Reducing the Uncertainties. U.S. National Oceanic and Atmospheric Administration Coastal Ocean Program Decision Analysis Series No. 9. Washington, DC.
- ⁵⁷ Moser, F.C. 1997. Sources and sinks of nitrogen and trace metals, and benthic macrofauna assemblages in Barnegat Bay, New Jersey. Ph.D. Dissertation, Rutgers, The State University of New Jersey, New Brunswick, New Jersey.
- ⁵⁸ Chesapeake Bay Program. 1994. Annual Report #2 on atmospheric N deposition: 1990-1994. Chesapeake Bay Program, U.S. Environmental Protection Agency, Annapolis, MD.
- ⁵⁹ Correll, D. L., and D. Ford. 1982. Comparison of precipitation and land runoff as sources of estuarine nitrogen. *Estuarine, Coastal and Shelf Science* 15:45-56.
- ⁶⁰ Whitall, D.R. and H.W. Paerl. 2001. Spatiotemporal variability of wet atmospheric nitrogen deposition to the Neuse River Estuary, North Carolina. *J Environ Qual* 30: 1508-1515.
- ⁶¹ Paerl and Fogel 1994 (see 45).
- ⁶² Tampa Bay NEP (National Estuary Program). 1996. Charting the Course for Tampa Bay: Final Comprehensive Conservation and Management Plan. Tampa Bay National Estuary Program, St. Petersburg, Florida.
- ⁶³ Goolsby, D. A., W. A. Battaglin, G. B. Lowrance, R. S. Artz, B. J. Aulenbach and R. P. Hooper. 2000. Gulf of Mexico Hypoxia Assessment, Topic #3, Flux and Sources of Nutrients in the Mississippi-Atchafalaya River Basin. Draft Report to White House Office of Science and Technology Policy, Committee on Environment and Natural Resources, Hypoxia Work Group, Mississippi River/Gulf of Mexico Watershed Nutrient Task Force, 90 pp.
- ⁶⁴ Nixon 1995 (see 54).
- Paerl, H.W. 1997. Coastal eutrophication and harmful algal blooms: Importance of atmospheric deposition and groundwater as "new" nitrogen and other nutrient sources. *Limnol Oceanogr* 42:1154-1165.
- ⁶⁵ Paerl 1997 (see 64).
- ⁶⁶ Whitall and Paerl 2001 (see 60).
- ⁶⁷ Holland, E., F. Dentener, B. Braswell and J. Sulzman. 1999. Contemporary and pre-industrial global reactive nitrogen budgets. *Biogeochemistry* 43: 7-43.
- ⁶⁸ Duce 1986 (see 39).
- Bergametti, G., E. Remoudaki, R. Losno, E. Steiner, B. Chatenet and P. Buat-Menard. 1992. Source, transport and deposition of atmospheric phosphorus over the northwestern Mediterranean. *J Atmos Chem* 14:502-513.
- Jassby, A.D., J.E. Reuter, R.P. Axler, C.R. Goldman and S. Hackley. 1994. Atmospheric deposition of nitrogen and phosphorus in the annual nutrient load of Lake Tahoe (California, U.S.A.). *Water Resources Research* 30:2207-2216.
- Prospero, J.M., K. Barrett, T. Church, F. Dentener, R.A. Duce, J.N. Galloway, H. Levy II, J. Moody, and P. Quinn. 1996. Atmospheric

- deposition of nutrients to the North Atlantic Basin. *Biogeochemistry* 35:27-73.
- Herut, B., M.D. Krom, G. Pan and R. Mortimer. 1999. Atmospheric input of nitrogen and phosphorus to the Southeast Mediterranean: Sources, fluxes and possible impact. *Limnol Oceanogr* 44:1683-1692.
- ⁶⁹ Herut et al. 1999 (see 52.)
- Kay, R.T., G.P. Johnson and D.L. Schrader. 2000. Hydrology, water quality, and nutrient loads to Lake Catherine and Channel Lake, near Antioch, Lake County, Illinois. U.S. geological Survey Water Resources Investigations Report 00-4088. USGS, Illinois District.
- ⁷⁰ Redfield, A.C. 1958. The biological control of chemical factors in the environment. *Am. Scientist* 46:205-222.
- ⁷¹ Kay et al. 2000 (see 53.)
- USEPA (US Environmental Protection Agency). 1999. Deposition of Air Pollutants to the Great Waters. Third Report to Congress. US Govt. Printing Office, Washington, D.C.
- ⁷² Koelliker, Y., L. A. Totten, C. L. Gigliotti, J. H. Offenber, J. R. Reinfelder, Y. Zhuang, and S. J. Eisenreich. 2004. Atmospheric Wet Deposition of Total Phosphorus in New Jersey. WASP.
- ⁷³ Eisenreich, S.J., Emmling, P.J., and Beeton, A.M. 1977. Atmospheric loading of phosphorus and other chemicals in Lake Michigan. *J Great Lakes Res* 3 (3-4):291-304.
- ⁷⁴ Church, T.M., J.M. Tramontanto, J.R. Scudlark, T.D. Jickells, J. J. Tokos, and A.H. Knapp. 1984. The wet deposition of trace metals to the western Atlantic Ocean at the mid-Atlantic coast and Bermuda. *Atmos Environ* 18:2657-2664.
- Scudlark, J.R. and T.M. Church. 1997. Atmospheric deposition of trace elements to the mid-Atlantic bight. pp. 195-208, in J. E. Baker [Ed], *Atmospheric Deposition of contaminants to the Great Lakes and Coastal Waters*. SETAC Press, Pensacola, Florida.
- Prospero et al., 1996 (see 68).
- ⁷⁵ Prospero et al. 1996 (see 68).
- ⁷⁶ Scudlark and Church 1997 (see 75).
- ⁷⁷ Zhuang, G., Z. Yi and G. T. Wallace. 1995. Iron (II) in rainwater, snow, and surface seawater from a coastal environment. *Mar Chem* 50:41-50.
- ⁷⁸ Simcik, M., H. Zhang, T. Franz, and S. J. Eisenreich. 1997. Urban contamination of the Chicago/Coastal Lake Michigan atmosphere by PCBs and PAHs during AEOLOS. *Environ Sci Technol* 31:2141-2147.
- Green, M. L., J. V. Depinto, C. Sweet, and K. C. Hornbuckle. 2000. Regional spatial and temporal interpolation of atmospheric PCBs: Interpretation of Lake Michigan mass balance data. *Environ Sci Technol* 34:1833-1841.
- ⁷⁹ Dachs, J., R. Lohmann, W.A. Ockenden, L. Méjanelle, S.J. Eisenreich and K.C. Jones. 2002. Oceanic biogeochemical controls on global dynamics of persistent organic pollutants. *Environ Sci Technol* 36 (20):4229-4237.
- Brunciak P. A., J. Dachs, T. P. Franz, C. L. Gigliotti, E. D. Nelson, B. J. Turpin, and S. J. Eisenreich. 2001. Polychlorinated Biphenyls and Particulate Organic/Elemental Carbon in the Chesapeake Bay Atmosphere. *Atmos Environ* 35:5663-5677.
- ⁸⁰ Franz, T.P.; S.J. Eisenreich; T.M. Holsen. 1998. Dry deposition of particulate PCBs and PAHs to Lake Michigan. *Environ Sci Technol* 32 (23):3681-3688.
- ⁸¹ Achman, D.R., K.C. Hornbuckle, and S.J. Eisenreich. 1993. Volatilization of polychlorobiphenyl congeners from Green Bay, Lake Michigan. *Environ Sci Technol* 27: 75-87.
- Totten, L. A., C. L. Gigliotti, J.H. Offenber, J. Baker, and S. J. Eisenreich. 2003. Re-evaluation of air-water exchange fluxes of PCBs in Green Bay (GBMB) and Southern Lake Michigan (AEOLOS, LMMB). *Environ Sci Technol* 37(9): 1739-1743.
- ⁸² Totten, L.A., P. A. Brunciak, C. L. Gigliotti, J. Dachs, T. R. Glenn, E.D. Nelson, and S.J. Eisenreich. 2001. Dynamic air-water exchange of polychlorinated biphenyls in the New York-New Jersey Harbor Estuary. *Environ Sci Technol* 35(19):3834-3840.
- Gigliotti, C.L., L. Totten, P.A. Brunciak, J. Dachs, E.D. Nelson, R. Lohmann, S.J. Eisenreich. 2002. Air-Water Exchange of PAHs in the NY-NJ Harbor Estuary. *Environ Toxic Chem* 21(2):235-244.
- Lohmann, R., E.D Nelson, S.J. Eisenreich, and K.C. Jones. 2000. Evidence for dynamic air - water exchange of PCDD/Fs: A Study in the Raritan Bay/Hudson River Estuary, USA. *Environ Sci Technol* 34(15):3086-3093.
- Van Ry, D.A., J. Dachs, C.L. Gigliotti, P. Brunciak, E.D. Nelson, and S.J. Eisenreich. 2000. Atmospheric seasonal trends and environmental fate of alkylphenols in the lower Hudson River Estuary. *Environ Sci Technol* 34(12):2410-2417.
- ⁸³ Bamford, H., J.H. Offenber, R.K. Larsen, F.-C. Ko, and J.E. Baker. 1999. Diffusive exchange of Polycyclic Aromatic Hydrocarbons across the air-water Interface of the Patapsco River, an Urbanized Subestuary of the Chesapeake Bay, *Environ Sci Technol* 33:2138-2144.
- Bamford, H.A., F.-C. Ko, and J.E. Baker. 2002. Seasonal and annual air-water exchange of Polychlorinated Biphenyls across Baltimore Harbor and the Northern Chesapeake Bay. *Environ Sci Technol* 36:4245-4252.
- ⁸⁴ Jeremiason, J.D., K. Hornbuckle, and S.J. Eisenreich. 1994. Polychlorinated biphenyls (PCBs) in Lake Superior, 1978-1992: Decreases in water concentrations reflect loss by volatilization. *Environ Sci Technol* 28: 903-914.
- ⁸⁵ Swackhamer, D. L., Schottler, S., Pearson, R. F. 1999. Air-water exchange and mass balance of toxaphene in the Great Lakes. *Environ Sci Technol* 33(21):3864-3872.
- ⁸⁶ Jeremiason et al. 1994 (see 84).
- ⁸⁷ Schottler, S.P. and S.J. Eisenreich. 1997. A mass balance model to quantify atrazine sources, transformation rates, and trends in the Great Lakes. *Environ Sci Technol* 31(9):2616-2625.
- ⁸⁸ Macdonald, R., S.J. Eisenreich, T.F. Bidleman, J. Dachs, J. Pacyna, K.C. Jones, R. Bailey, D. Swackhamer, and D. Muir . 2000. Case studies on persistence and long range transport of persistent organic pollutants (Chapter 7). In Klecka, G.M., D. Mackay, J. Franklin, D. Graham, P. Howard, L. Grady, K. Kannan, R.J. Larson, D. Muir, and D. Van de Meent (Eds.) *Evaluation of Persistence and Long-range Transport of Organic Chemicals in the Environment* p.245-314. SETAC Press: Pensacola, FL. Figure reprinted with permission from SETAC, Pensacola, FL.
- ⁸⁹ Dachs, J., S.J. Eisenreich, J.E. Baker, F.-C. Ko, and J.D. Jeremiason. 1999. Coupling of phytoplankton uptake and air-water exchange of persistent organic pollutants. *Environ Sci Technol* 33(20), 3653-3660.
- Dachs et al. 2002 (see 9)
- Jeremiason et al. 1994 (see 84).
- Jeremiason, J.J., S.J. Eisenreich, M.J. Paterson, D. Findlay, H.J. Kling, K. Beaty, R. Hecky, and J. Elser. 1999. Biogeochemical cycling of atmospherically-derived PCBs in lakes of variable trophic status: a paired lake experiment. *Limnol Oceanogr* 44(3, Part 2):889-902.

- Totten et al. 2001 (see 82).
 Totten et al. 2003 (see 81).
 Gigliotti et al. 2002 (see 82).
 Lohmann et al. 2002 (see 82).
 Swackhamer et al. 1999 (see 85).
 MacDonald et al. 2000 (see 88).
- ⁹⁰Dachs et al. 1999 (see 89).
⁹¹Dachs et al. 1999 (see 89).
⁹²Dachs et al. 1999 (see 89).
 Dachs, J., S.J. Eisenreich, and R.M. Hoff. 2000. Influence of eutrophication on air-water exchange, vertical fluxes and phytoplankton concentrations of POPs. *Environ Sci Technol* 34(6):1095-1102.
 Dachs et al. 2002 (see 9).
- ⁹³Yan, S.; J. Dachs, C.W. Fan, J. Reinfelder, S. J. Eisenreich. 2004. Air-water exchange controls phytoplankton concentrations of polychlorinated biphenyls in impacted estuaries. *Environ Sci Technol* Submitted.
⁹⁴MacDonald et al., 2000 (see 88).
⁹⁵Kidd K.A., D.W. Schindler, D.C.G. Muir, W.L. Lockhart, and R.H. Hesselin. 1995. High concentrations of toxaphene in fishes from a subarctic lake. *Science* 269:240-241.
⁹⁶Cook, P., J.A. Robbins, D.D. Endicott, K.B. Lodge, P.D. Guiney, M.K. Walker, E.W. Zabelo and R.E. Peterson. 2003. Effects of aryl hydrocarbon receptor-mediated early life stage toxicity on lake trout populations in Lake Ontario during the 20th Century. *Environ Sci Technol* 37(17):3864-3877.
⁹⁷DeVault, D. S., R. Hesselberg, P.W. Rodgers, and T.J. Feist. 1996. Contaminant trends in Lake Trout and Walleye from the Laurentian Great Lakes. *J Great Lakes Res* 22(4):884-895.
⁹⁸Blais, J.M., D.W. Schindler, M. Sharp, E. Braekvelt, M. Laferniere, D.C.G. Muir, W.Strachan. 2001. Fluxes of semi-volatile organochlorines in Bow Lake, a remote high altitude, glacier-fed sub-alpine lake in the Canadian Rocky Mountains. *Limnol Oceanogr* 46: 2019-2031.
⁹⁹AMAP (Arctic Monitoring and Assessment Programme). 1998. Assessment report: Arctic Pollution Issues. Oslo.
¹⁰⁰AMAP 1998 (see 99).
¹⁰¹Donald, D.B., J. Syrgiannis, R.W. Crosley, G. Holdsworth, D.C.G. Muir, B. Rosenberg, A. Sole and D.W. Schindler. 1999. Delayed deposition of organochlorine pesticides at a temperate glacier. *Environ Sci Technol* 33: 1794-1798.
¹⁰²Holsen, T. M., K. E. Noll, S. P. Liu, and W. J. Lee. 1991. Dry Deposition of Polychlorinated-Biphenyls in Urban Areas. *Environ Sci Technol* 25:1075-1081.
 Cotham, W. E., and T. F. Bidleman. 1995. Polycyclic Aromatic-Hydrocarbons and Polychlorinated-Biphenyls in air at an urban and a rural site near Lake-Michigan. *Environ Sci Technol* 29:2782-2789.
 Zhang, H. X., S. J. Eisenreich, T. R. Franz, J. E. Baker, and J. H. Offenberg. 1999. Evidence for increased gaseous PCB fluxes to Lake Michigan from Chicago. *Environ Sci Technol* 33:2129-2137.
 Green et al. 2000 (see 78).
 Offenberg, J. H., and J. E. Baker. 2000. PCBs and PAHs in southern Lake Michigan in 1994 and 1995: Urban atmospheric influences and long-term declines. *J Great Lakes Res* 26:196-208.
 Miller, S. M., M. L. Green, J. V. Depinto, and K. C. Hornbuckle. 2001. Results from the Lake Michigan mass balance study: Concentrations and fluxes of atmospheric polychlorinated biphenyls and trans-nonachlor. *Environ Sci Technol* 35:278-285.
- Hsu, Y. K., T. M. Holsen, and P. K. Hopke. 2003. Locating and quantifying PCB sources in Chicago: Receptor modeling and field sampling. *Environ Sci Technol* 37:681-690.
 Buehler, S. S., I. Basu, and R. A. Hites. 2004. Causes of variability in pesticide and PCB concentrations in air near the Great Lakes. *Environ Sci Technol* 38:414-422.
- ¹⁰³Wania, F. and D. MacKay. 1993. Global fractionation and cold condensation of low volatility organo-chlorine compounds in polar regions. *Ambio* 22: 10-18.
¹⁰⁴Muir, D.C.G., Wagemann, R., Hargrave, B.T., Thomas, D.J., Peakall, D.B. and Norstrom, R.J. 1992. Arctic marine ecosystem contamination. *Sci Total Environ* 122:75-134.
 Norstrom R.J., S.E. Belikov, E.W. Born, G.W. Garner, B. Malone, S. Olpinski, et al. 1998. Chlorinated hydrocarbon contaminants in polar bears from eastern Russia, North America, Greenland, and Svalbard: biomonitoring of Arctic pollution. *Arch Environ Contam Toxicol* 35:354-367.
 Blais, J.M., D.W. Schindler, D.C.G. Muir, L.E. Kimpe, D.B. Donald, B. Rosenberg. 1998. Accumulation of persistent organochlorine compounds in mountains of western Canada. *Nature* 395: 585-588.
 Carrera, G., P. Fernández, J.O. Grimalt, M. Ventura, L. Camarero, J. Catalán, U. Nickus, H. Thies, and R. Psenner. 2002. *Environ Sci Technol* 36:2587.
- ¹⁰⁵AMAP 1998 (see 99).
 Donald et al. 1999 (see 101).
¹⁰⁶AMAP 1998 (see 99).
¹⁰⁷Schindler, D.W., K.A. Kidd, D.C.G. Muir, and W.L. Lockhart. 1995. The effects of ecosystem characteristics on contaminant distribution in northern freshwater lakes. *Sci Total Environ* 160(161):1-17.
¹⁰⁸AMAP 1998 (see 99).
 Muir et al. 1992 (see 104).
 Dewailly, E., P. Ayotte, S. Bruneau, C. Lalibert, D. Muir, and R. Norstrom. 1993. Inuit exposure to organochloride through the aquatic food chain in arctic Quebec. *Environ Health Persp* 101:618-20.
¹⁰⁹AMAP 1998 (see 99).
 Jacobson J.L and S.W. Jacobson. 1996. Intellectual impairment in children exposed to polychlorinated biphenyls in utero. *N Engl J Med* 335:783-789.
¹¹⁰Kidd et al. 1995 (see 95).
¹¹¹Modified from Hornbuckle, K. C., and M. L. Green. 2003. The impact of an urban-industrial region on the magnitude and variability of persistent organic pollutant deposition to Lake Michigan. *Ambio* 32:406-411.
¹¹²Donald, D., R. Bailey, R. Crosley, D. Muir, P. Shaw and J. Syrgiannis. 1993. Polychlorinated biphenyls and organochlorine pesticides in the aquatic environment along the continental divide region of Alberta and British Columbia, Inland Waters Directorate, Regina, Sask.
¹¹³Blais et al. 1998 (see 104).
 Donald et al. 1999 (see 101).
 Blais, J.M., K. Duff, D.W. Schindler, J.P. Smol, P.R. Leavitt, M. Agbeti. 2000. Assessment of recent eutrophication histories in Lac Ste. Anne and Lake Isle, Alberta, Canada, using paleolimnological methods. *Lake and Reservoir Management* 16: 292-304.
¹¹⁴Campbell, L.M., D.W. Schindler, D.C.G. Muir, D.B. Donald, and K.A. Kidd. 2000. Organochlorine transfer in the food web of

- subalpine Bow Lake, Banff National Park. *Canadian Journal of Fisheries and Aquatic Science*, 57: 1258-1269.
- Blais et al. 2000 (see 113)
- Braekvelt, E., G.T. Tomy, and G.A. Stern. 2001. Comparison of an individual congener standard and a technical mixture for the qualification of toxaphene in environmental matrices by HRGC/ECNI-HRMS. *Environ Sci* 35(17):3513-3518.
- ¹¹⁵Grimalt, J.O., P. Fernandez and R.M. Vilanova. 2001. Trapping of organochlorine compounds in high mountain lakes. *The Scientific World* 1:609-611.
- Carrera et al. 2002 (see 104).
- ¹¹⁶Skaare, J.U., Ø. Wiig and A. Bernhoft, 1994. Klorerte organiske miljøgifter; nivåer og effekter på isbjørn. Norsk Polarinstituttets Rapportserie 86, 27 pp.
- ¹¹⁷AMAP 1998 (see 99).
- ¹¹⁸Ahlborg, V.G., G.C. Becking, L.S. Birnbaum, A. Brower, H.J.G.M. Derks, M. Feeley, C. Golor, A. Hanberg, J.C. Larsen, A.K.D. Liem, S.H. Safe, C. Schlatter, F. Waern, M. Younes, and E. Yrkanekki, E. 1994. Toxic equivalency factors for dioxin-like PCBs. *Chemosphere*. 28(6):1049-1067.
- ¹¹⁹D. Muir, pers. comm
- ¹²⁰Ikonomou et al. 2002 (see 22).
- Boon, J. P., J.J. van Zanden, W.E. Lewis, B.N. Zegers, A. Goksoyr, and A. Arukwe. 2002. *Marine Environmental Research*, 54(3-5):719-724.
- ¹²¹Norstrom R.J., M. Simon, J. Moisey, B. Wakeford, and D.V.C. Weseloh. 2002. Geographical distribution (2000) and temporal trends (1981 to 2000) of brominated diphenyl ethers in Great Lakes herring gull eggs.
- ¹²²Mierle, G. and R. Ingram. 1991. The role of humic substances in the mobilization of mercury from watersheds. *Water Air and Soil Pollution* 56: 349-357.
- ¹²³Wiener, J. G., D. P. Krabbenhoft, G. H. Heinz, and A. M. Scheuhammer. 2003. Ecotoxicology of mercury. Pages 409-463 in J. J. Cairns, editor. *Handbook of Ecotoxicology*. Lewis Publishers, Boca Raton, FL.
- ¹²⁴Scheuhammer, A. M. 1991. Effects of Acidification on the Availability of Toxic Metals and Calcium to Wild Birds and Mammals. *Environ Poll* 71:329-375.
- Hammerschmidt, C. R., M. B. Sandheinrich, J. G. Wiener, and R. G. Rada. 2002. Effects of dietary methylmercury on reproduction of fathead minnows. *Environ Sci Technol* 36:877-883.
- ¹²⁵Grandjean, P., P. Weihe, R. F. White, F. Debes, S. Araki, K. Yokoyama, K. Murata, N. Sorensen, R. Dahl, and P. J. Jorgensen. 1997. Cognitive deficit in 7-year-old children with prenatal exposure to methylmercury. *Neurotoxicol Teratol* 19:417-428.
- Davidson, P.W., D. Palumbo, G.J. Myers, C. Cox, C. F. Shamlaye, J. Sloane-Reeves, E. Cernichiari, G. E. Wilding, and T. W. Clarkson. 2000. Neurodevelopmental outcomes of seychellois children from the pilot cohort at 108 months following prenatal exposure to methylmercury from a maternal fish diet. *Environ Res* 84:1-11.
- NRC (National Research Council). 2000. *Clean Coastal Waters: Understanding and Reducing the Effects of Nutrient Pollution*. National Academy Press, Washington, DC.
- ¹²⁶NRC 2000 (see 125).
- ¹²⁷Stolte, W., T. McCollin, A. Noordeloos, and R. Riegman. 1994. Effects of nitrogen source on the size distribution within marine phytoplankton populations. *J Exper Mar Biol Ecol* 184:83-97.
- Pinckney, J.L., H.W. Paerl, and M.B. Harrington. 1999. Responses of the phytoplankton community growth rate to nutrient pulses in variable estuarine environments. *J Phycol* 35:1455-1463.
- ¹²⁸Cornell, S., A. Rendell, and Jickells. 1995. Atmospheric inputs of dissolved organic nitrogen to the oceans. *Nature* 376:243-246.
- Peierls, B.L. and H.W. Paerl. 1997. The bioavailability of atmospheric organic nitrogen deposition to coastal phytoplankton. *Limnol Oceanogr* 42:1819-1880.
- Seitzinger, S. P. and R. W. Sanders. 1999. Atmospheric inputs of organic nitrogen stimulate estuarine bacteria and phytoplankton. *Limnol Oceanogr* 44:721-730.
- Antia N., P. Harrison, and L. Oliveira. 1991. The role of dissolved organic nitrogen in phytoplankton nutrition, cell biology and ecology. *Phycologia* 30:1-89.
- ¹²⁹Prospero et al. 1996 (see 68).
- Takeda et al. 1995 (see 40).
- Paerl et al. 1999 (see 41).
- ¹³⁰Tyrell, T. 1999. The relative influences of nitrogen and phosphorus on oceanic primary production. *Nature* 400:525-531.
- ¹³¹Boesch, D. F., E. Burreson, W. Dennison, E. Houde, M. Kemp, V. Kennedy, R. Newell, K. Paynter, R. Orth and R. Ulanowicz. 2001. Factors in the decline of coastal ecosystems. *Science* 293:629-638.
- Elmgren and Larsson 2001 (see 44).
- ¹³²Duce 1986 (see 39).
- GESAMP 1989 (see 51).
- Herut et al. 1999 (see 69).
- ¹³³Martin et al. 1994 (see 40).
- Takeda et al. 1995 (see 40)
- Paerl, H. W., L. Prufert Bebout and C. Guo. 1994. Iron stimulated N₂ fixation and growth in natural and cultured populations of the planktonic marine cyanobacterium *Trichodesmium*. *Appl Environ Microbiol* 60:1044-1047.
- Paerl, H.W., J. Pinckney, J. Fear and B. Peierls. 1998. Ecosystem responses to internal and watershed organic matter loading: Consequences for hypoxia in the eutrophying Neuse River Estuary, NC, USA. *Mar Ecol Progr Ser* 166:17-25.
- ¹³⁴Zhang, J. 1994. Atmospheric wet deposition of nutrient elements: Correlation with harmful biological blooms in Northwest Pacific coastal zones. *Ambio* 23:464-468.
- Walsh, J. J. and K.A. Steidinger. 2001. Saharan Dust and Florida red tides: The cyanophyte connection. *J Geophys Res* 106:11597-11612.
- ¹³⁵Scudlark and Church 1997 (see 74).
- ¹³⁶Takeda et al. 1995 (see 40).
- Paerl et al. 1999 (see 41).
- ¹³⁷Stolte et al. 1994 (see 127).
- ¹³⁸Eppley, R.W., J. N. Rogers, and J.J. McCarthy. 1969. Half saturation constants for uptake of nitrate and ammonia by marine phytoplankton. *Limnol Oceanogr* 14:912-920.
- ¹³⁹Syrett, P.J. 1981. Nitrogen metabolism of microalgae. *Canadian Bulletin of Fisheries and Aquatic Sciences* 210:182-210.
- Molloy, C., and P. Syrett. 1988. Interrelationships between uptake of urea and uptake of ammonium by microalgae. *J Exper Mar Biol Ecol* 118:85-95.
- Dortch, Q. 1990. The interaction between ammonium and nitrate uptake in phytoplankton. *Mar Ecol Progr Ser* 61:183-201.
- ¹⁴⁰Eppley et al. 1969 (see 138).
- Stolte et al. 1994 (see 127).

- ¹⁴¹Harrington, M.B. 1999. Responses of natural phytoplankton communities from the Neuse River Estuary, NC to changes in nitrogen supply and incident irradiance MSc. Thesis. University of North Carolina at Chapel Hill.
- ¹⁴²Cloern, J.E. 1999. The relative importance of light and nutrient limitation of phytoplankton growth: a simple index of coastal ecosystems sensitivity to nutrient enrichment. *Aquatic Ecology* 33:3-16.
- ¹⁴³Peierls and Paerl 1997 (see 128).
- ¹⁴⁴Antia et al. 1991 (see 128).
- Paerl, H.W. 1991. Ecophysiological and trophic implications of light stimulated amino acid utilization in marine picoplankton. *Appl Environ Microbiol* 57:473-479.
- ¹⁴⁵Skoglund, R.S. and D.L. Swackhamer. 1994. Fate of hydrophobic organic contaminants: processes affecting uptake by phytoplankton, p. 559-574. In Baker, L.A. (ed.), *Environmental chemistry of lakes and reservoirs*. American Chemical Society, Washington, D.C.
- Skoglund RS, K. Stange, and D.L. Swackhamer. 1996. A kinetics model for predicting the accumulation of PCBs in phytoplankton. *Environ Sci Technol* 30:2113-2120.
- Swackhamer D.L. and R.S. Skoglund. 1991. The role of phytoplankton in the partitioning of hydrophobic organic contaminants in water. In Baker, R.A. (Ed). *Organic Substances and Sediments in Water*. Lewis, Boca Raton FL, USA, pp 91-105.
- Swackhamer D.L. and R.S. Skoglund. 1993 Bioaccumulation of PCBs by phytoplankton: kinetics vs. equilibrium. *Environ Toxicol Chem* 12:831-838.
- ¹⁴⁶Hudson, M. 2004. Microbial Facilitation of Contaminant Transfer in Lake Superior Foodwebs. MS Thesis, Water Resources Science, University of Minnesota.

ABOUT THE PANEL OF SCIENTISTS

- Deborah L. Swackhamer, Co-Chair, School of Public Health, University of Minnesota, Minneapolis, MN 55455
- Hans W. Paerl, Co-Chair, Institute of Marine Sciences, The University of North Carolina at Chapel Hill, Morehead City, NC 28557
- Steven J. Eisenreich, Joint Research Centre of the European Commission, Institute for Environment and Sustainability, Ispra, Italy
- James Hurley, Aquatic Sciences Center, University of Wisconsin, Madison, WI 53706
- Keri C. Hornbuckle, Seamans Center for the Engineering Arts and Sciences, University of Iowa, Iowa City, Iowa 52242
- Michael McLachlan, The Institute of Applied Environmental Research, Stockholm University, Stockholm Sweden
- David Mount, Ecotoxicology Analysis Research Branch, Mid-Continent Ecology Division, U.S. Environmental Protection Agency, Duluth, MN 55804 USA
- Derek Muir, National Water Research Institute, Environment Canada, Burlington, Ontario L7R 4A6 Canada
- David Schindler, Biological Sciences, University of Alberta, Edmonton, Alberta T6G 2E9 Canada

About the Science Writer

Yvonne Baskin, a science writer, edited the report of the panel of scientists to allow it to more effectively communicate its findings with non-scientists.

About Issues in Ecology

Issues in Ecology is designed to report, in language understandable by non-scientists, the consensus of a panel of scientific experts on issues relevant to the environment. *Issues in Ecology* is supported by a Pew Scholars in Conservation Biology grant to David Tilman and by the Ecological Society of America. All reports undergo peer review and must be approved by the editorial board before publication. No responsibility for the views expressed by authors in ESA publications is assumed by the editors or the publisher, the Ecological Society of America.

Editorial Board of Issues in Ecology

Dr. William Murdoch, Editor-in-Chief, Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, CA 93106. Email: murdoch@lifesci.ucsb.edu.

Board members

- Dr. Peter Kareiva, The Nature Conservancy, Seattle, WA 98105.
- Dr. Ann Kinzig, Department of Biology, Arizona State University, Tempe, AZ 85287-1501.
- Dr. Jane Lubchenco, Department of Zoology, Oregon State University, Corvallis, OR 97331-2914.
- Dr. Judy L. Meyer, Institute of Ecology, University of Georgia, Athens, GA 30602-2202.
- Dr. Gordon Orians, Department of Zoology, University of Washington, Seattle, WA 98195.
- Dr. Lou Pitelka, Appalachian Environmental Laboratory, Gunter Hall, Frostburg, MD 21532.
- David Schimel, National Center for Atmospheric Research, Editor, *Ecological Applications*, Boulder, CO 80305.
- Dr. William Schlesinger, Departments of Botany and Geology, Duke University, Durham, NC 27708-0340.
- Dr. David Wilcove, Woodrow Wilson School of Public and International Affairs, Princeton University, Princeton, NJ 08544-1013.

Previous Reports

Previous *Issues in Ecology* reports available from the Ecological Society of America include:

- Vitousek, P.M., J. Aber, R.W. Howarth, G.E. Likens, P.A. Matson, D.W. Schindler, W.H. Schlesinger, and G.D. Tilman. 1997. Human Alteration of the Global Nitrogen Cycle: Causes and Consequences, *Issues in Ecology* No. 1.
- Daily, G.C., S. Alexander, P.R. Ehrlich, L. Goulder, J. Lubchenco, P.A. Matson, H.A. Mooney, S. Postel, S.H. Schneider, D. Tilman, and G.M. Woodwell. 1997. Ecosystem Services: Benefits Supplied to Human Societies by Natural Ecosystems, *Issues in Ecology* No. 2.
- Carpenter, S., N. Caraco, D. L. Correll, R. W. Howarth, A. N. Sharpley, and V. H. Smith. 1998. Nonpoint Pollution of

- Surface Waters with Phosphorus and Nitrogen, *Issues in Ecology* No. 3.
- Naeem, S., F.S. Chapin III, R. Costanza, P.R. Ehrlich, F.B. Golley, D.U. Hooper, J.H. Lawton, R.V. O'Neill, H.A. Mooney, O.E. Sala, A.J. Symstad, and D. Tilman. 1999. Biodiversity and Ecosystem Functioning: Maintaining Natural Life Support Processes, *Issues in Ecology* No. 4.
- Mack, R., D. Simberloff, W.M. Lonsdale, H. Evans, M. Clout, and F. Bazzaz. 2000. Biotic Invasions: Causes, Epidemiology, Global Consequences and Control, *Issues in Ecology* No. 5.
- Aber, J., N. Christensen, I. Fernandez, J. Franklin, L. Hiding, M. Hunter, J. MacMahon, D. Mladenoff, J. Pastor, D. Perry, R. Slangen, H. van Miegroet. 2000. Applying Ecological Principles to Management of the U.S. National Forests, *Issues in Ecology* No. 6.
- Howarth, R., D. Anderson, J. Cloern, C. Elfring, C. Hopkinson, B. LaPointe, T. Malone, N. Marcus, K. McGlathery, A. Sharpley, and D. Walker. Nutrient Pollution of Coastal Rivers, Bays, and Seas, *Issues in Ecology* No. 7.
- Naylor, R., R. Goldburg, J. Primavera, N. Kautsky, M. Beveridge, J. Clay, C. Folke, J. Lubchenco, H. Mooney, and M. Troell. 2001. Effects of Aquaculture on World Fish Supplies, *Issues in Ecology* No. 8.
- Jackson, R., S. Carpenter, C. Dahm, D. McKnight, R. Naiman, S. Postel, and S. Running. 2001. Water in a Changing World, *Issues in Ecology* No. 9.
- Baron, J.S., N.L. Poff, P.L. Angermeier, C.N. Dahm, P.H. Gleick, N.G. Hairston, Jr., R.B. Jackson, C.A. Johnston, B.D. Richter, and A.D. Steinman. 2003. Sustaining Healthy Freshwater Ecosystems, *Issues in Ecology* No. 10.
- Beck, M.W., K.L. Heck, Jr., K. W. Able, D.L. Childers, D.B. Eggleston, B.M. Gillanders, B.S. Halpern, C.G. Hays, K. Hoshino, T.J. Minello, R.J. Orth, P.F. Sheridan, and M. P. Weinstein. 2003. The Role of Nearshore Ecosystems as Fish and Shellfish Nurseries, *Issues in Ecology* No. 11.

Additional Copies

To receive additional copies of this report (\$3 each) or previous *Issues in Ecology*, please contact:

Ecological Society of America
1707 H Street, NW, Suite 400
Washington, DC 20006
(202) 833-8773, esahq@esa.org



The *Issues in Ecology* series is also available electronically at http://www.esa.org/sbi/sbi_issues/